arXiv:1106.4977v1 [math.AG] 24 Jun 2011

MIRROR SYMMETRY FOR LOG CALABI-YAU SURFACES I

MARK GROSS, PAUL HACKING, AND SEAN KEEL

Abstract. We give a canonical synthetic construction of the mirror family to a pair (Y, D) of a smooth projective surface with an anti-canonical cycle of rational curves, as the spectrum of an explicit algebra defined in terms of counts of rational curves on Y meeting D in a single point. In the case D is contractible, the family gives a smoothing of the dual cusp, and thus a proof of Looijenga's 1981 cusp conjecture.

Contents

Introduction 2
0.1. The main theorems 2
0.2. Consequences of the main theorem 5
0.3. Description of the coordinate ring 8
0.4. Frobenius structure conjecture 11
0.5. Mirror symmetry conjectures 13
0.6. Overview of the proof 18
0.7. Acknowledgements. 27
1. Basics 28
1.1. Tropical Looijenga pairs 28
1.2. The Mumford degeneration 37
1.3. Some toric constructions 40
2. Modified Mumford deformations 42
2.1. Generalities on integral linear manifolds and multi-valued functions 42
2.2. Scattering diagrams on B 50
2.3. Broken lines 55
2.4. The algebra structure 62
3. The canonical scattering diagram 63
3.1. Reduction to the Gross–Siebert locus 68
3.2. The proof of Theorem 3.32 83
3.3. The proof of Proposition 3.26: The connection with [GPS09] 88
3.4. Smoothness: Around the Gross–Siebert locus 92
3.5. The relative torus 97
4. Looijenga's conjecture 98

4.1. Cusp family 98
4.2. Thickening of cusp family 104
4.3. Smoothness 113
5. Extending the family over boundary strata 118
5.1. Proof of Theorems 0.1 and 0.2 for $n < 3$ 122
5.2. D positive 123
5.3. Relation to cluster varieties 126
6. Deformations of cyclic quotient singularities 128
6.1. Construction of the deformation 128
6.2. P-resolutions and deformations 133
6.3. Main Theorem 134
6.4. Deformations of pairs 136
References 140

INTRODUCTION

0.1. **The main theorems.**

Let $\mathbb{V}_n$ for $n \geq 3$ be the n-cycle of coordinate planes in $\mathbb{A}^n$,

$$\mathbb{V}_n := \mathbb{A}^2_{x_1,x_2} \cup \mathbb{A}^2_{x_2,x_3} \cup \cdots \cup \mathbb{A}^2_{x_n,x_1} \subset \mathbb{A}^n_{x_1,\ldots,x_n}.$$

We refer to $\mathbb{V}_n$ as the *vertex of degree n*. (See (1.5) and (1.6) for the definition of $\mathbb{V}_1$ and $\mathbb{V}_2$.) The surfaces $\mathbb{V}_n$ are the most important singularities for moduli of smooth surfaces. Recall that a functorial compactification of moduli of canonically polarized manifolds is obtained by allowing so-called stable varieties — varieties with semi-log canonical (SLC) singularities and ample dualizing sheaf [K11]. One-dimensional SLC singularites are ordinary nodes. Every SLC germ is (in a canonical way) a quotient of a Gorenstein SLC germ by a finite cyclic group, and every Gorenstein SLC surface germ is obtained as a deformation of $\mathbb{V}_n$ (for some n), which is itself Gorenstein and SLC. Here we use mirror symmetry to construct canonical embedded smoothings, jam-packed with geometric information.

Throughout the paper $(Y, D = D_1 + \cdots + D_n)$ will denote a smooth projective (necessarily rational) surface, with $D \in |-K_Y|$ a singular nodal curve (either an irreducible rational nodal curve, or a cycle of $n \geq 2$ smooth rational curves) over an algebraically closed field $\Bbbk$ of characteristic zero. We call (Y, D) a *Looijenga pair* for, as far as we know, their rich geometry was first investigated in [L81]. We cyclically order the components of D and take indices modulo n. By assumption there is a holomorphic symplectic 2-form Ω, unique up to scaling, on $Y \setminus D$, with simple poles along D, and thus $U := Y \setminus D$ is a log Calabi-Yau surface. Our main result is a synthetic construction of the mirror family, which gives a canonical (and in many cases modular) embedded smoothing of $\mathbb{V}_n \subset \mathbb{A}^n$ parameterized roughly by the formal completion of the affine toric variety $\operatorname{Spec} \Bbbk[\mathrm{NE}(Y)]$ along the union of toric boundary strata corresponding to contractions $f\colon Y \to \bar{Y}$. Here $\mathrm{NE}(Y)$ denotes the monoid $\mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}} \cap A_1(Y, \mathbb{Z})$ where $\mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}} \subset A_1(Y, \mathbb{R})$ is the cone generated by curve classes. This is just an approximate statement of our result, as $\mathrm{NE}(Y)$ is not in general finitely generated, and in some cases we restrict to analytic neighbourhoods of the origin. By restricting the family to slices of the base transverse to various boundary strata, we recover most of what is known about deformations of log canonical surface singularities, unifying the results in a single Mori and mirror theoretic framework, and settle the main open question in the subject, Looijenga's 1981 conjectural criterion for smoothability of cusp singularities.

More precisely, let $B_0(\mathbb{Z})$ be the set of pairs (E, n) where E is a divisor on some blowup of Y along which Ω has a pole and n is a positive integer. Set $B(\mathbb{Z}) := B_0(\mathbb{Z}) \cup \{0\}$. See §0.3.1 below for another description of this set (as integer points in a natural integral affine manifold). Let $v_i \in B(\mathbb{Z})$ be the pair $(D_i, 1)$. Choose $\sigma_P \subset A_1(Y, \mathbb{R})$ a strictly convex rational polyhedral cone containing $\mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}}$, and let $P := \sigma_P \cap A_1(Y, \mathbb{Z})$ be the associated monoid and $R = \Bbbk[P]$ the associated $\Bbbk$-algebra.

For each monomial ideal $I \subset R$, consider the free $R_I := R/I$-module

$$A_I := \bigoplus_{q \in B(\mathbb{Z})} R_I \cdot \vartheta_q. \tag{0.1}$$

Let $\mathfrak{m} \subset R$ denote the maximal monomial ideal. Let $T^D = \mathbb{G}_m^n$ be the torus with character group $\chi(T^D)$ having basis e_{D_i} indexed by the components $D_i \subset D$. There is a homomorphism $T^D \to \operatorname{Spec} \Bbbk[P^{\mathrm{gp}}]$ induced by $C \mapsto \sum (C \cdot D_i) e_{D_i}$, so T^D acts on $\operatorname{Spec} R_I$. In §3.5, an action of T^D on A_I is also constructed.

Theorem 0.1. *Let $I \subset R$ be a monomial ideal with $\sqrt{I} = \mathfrak{m}$. There is a canonical T^D-equivariant finitely generated R_I-algebra structure on A_I, determined by relative Gromov-Witten invariants of (Y, D) counting rational curves meeting D in a single point. This induces a flat T^D-equivariant map*

$$f : X_I := \operatorname{Spec} A_I \to \operatorname{Spec} R_I.$$

The precise description of the multiplication rule is slightly involved, so we defer the details of this multiplication until after we have explained the applications, see §0.3.2. We also have a simple and elegant alternative that we conjecture gives the mirror in all dimensions directly in terms of counts of rational curves. See §0.4.

We use the notation ϑ_q for generators of our algebra, since for example, as we explain below, theta functions for polarized elliptic curves occur as a special case. Tyurin conjectured the existence of canonical theta functions (i.e., a basis of global sections) for polarized K3 surfaces and Kontsevich and Soibelman made similar speculations. Our results are log analogs of Tyurin's conjecture. In work in progress we apply similar ideas to obtain Tyurin's conjecture in the K3 case as well, see [K3].

Theorem 0.2. *There is a unique smallest radical monomial ideal* $J \subset R$ *with the following properties:*

1. *For every monomial ideal* I *with* $J \subset \sqrt{I}$ *there is a finitely generated* R_I*-algebra structure on* A_I *compatible with the* $R_{I+\mathfrak{m}^N}$*-algebra structure on* $A_{I+\mathfrak{m}^N}$ *of Theorem 0.1 for all* $N > 0$*.*
2. *If the intersection matrix* $(D_i \cdot D_j)$ *is not negative semi-definite then* $J = 0$*. In general, the zero locus* $V(J) \subset \operatorname{Spec} R$ *contains the union of the closed toric strata corresponding to faces* F *of* σ_P *such that* F *does not contain the class of some component of* D*.*
3. *Let* $\hat{R}$ *denote the* J*-adic completion of* R *and* $\operatorname{Spf} \hat{R}$ *the associated formal scheme. The algebras* A_I *determine a canonical* T^D*-equivariant formal flat family of affine surfaces*
$$f : \mathcal{X} \to \operatorname{Spf} \hat{R}$$
with fibre $\mathbb{V}_n$ *over* 0*, and with scheme-theoretic singular locus of* f *not surjecting scheme-theoretically onto the base.*
4. *The* ϑ_q *determine a canonical embedding* $\mathcal{X} \subset \mathbb{A}^{\max(n,3)} \times \operatorname{Spf} \hat{R}$*.*

We note the statement about the scheme-theoretic singular locus implies in particular that whenever the family is the formal completion of an algebraic family, the generic fibre of the algebraic family is smooth. As it is defined using Gromov-Witten invariants, the family depends only on the deformation type of (Y, D).

Remark 0.3. When $\mathrm{NE}(Y) \subset P' \subset P \subset A_1(Y)$, then $J' \subset J$ and the formal family $\mathcal{X}$ for P comes from the family for P' by base-change. In this sense the family is independent of the choice of P.

We conjecture the families constructed by the above theorems are mirror to $U = Y \setminus D$ in the sense of homological mirror symmetry in the case $\Bbbk = \mathbb{C}$. In Part II of this paper, we shall prove some cases of this by relating our construction to previously known mirror constructions. In addition we conjecture the algebra A is the symplectic cohomology ring $SH^0(U)$. See §0.5.1 for precise statements.

Next we explain the myriad applications independent of mirror symmetry.

0.2. **Consequences of the main theorem.** There are three broad classes of behaviour for our construction, depending on the properties of the intersection matrix $(D_i \cdot D_j)$.

The intersection matrix is not negative semi-definite. We call this the *positive* case. It holds iff U is the minimal resolution of an affine surface, see Lemma 5.9. In this case, the cone $\mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}}$ is rational polyhedral, so we may take $P = \mathrm{NE}(Y)$, and the ideal J of Theorem 0.2 equals 0. Thus our construction defines an algebraic family over $\operatorname{Spec} \Bbbk[\mathrm{NE}(Y)]$, with smooth generic fibre. We will show in Part II that the restriction of this family to the structure torus

$$\mathcal{X} \to T_Y := \mathrm{Pic}(Y) \otimes \mathbb{G}_m = \operatorname{Spec} \Bbbk[A_1(Y)] \subset \operatorname{Spec} R$$

is the universal family of deformations of $U = Y \setminus D$.

More precisely, we will show independently of the positivity of the intersection matrix that our formal family has a simple and canonical (fibrewise) compactification to a formal family $(\mathcal{Z}, \mathcal{D})$ of Looijenga pairs (with $\mathcal{X} = \mathcal{Z} \setminus \mathcal{D}$). In the positive case this extends naturally over all of $\operatorname{Spec} R$, and its restriction $(\mathcal{Z}, \mathcal{D}) \to T_Y$ comes with a trivialization of the boundary $\mathcal{D} = D_* \times T_Y$ realizing it as the universal family of Looijenga pairs (Z, D_Z) deformation equivalent to (Y, D) together with an isomorphism $D_Z \xrightarrow{\sim} D_*$. In particular, choosing such an isomorphism $D \xrightarrow{\sim} D_*$ for our original pair (Y, D) canonically identifies it with a fibre of the family $(\mathcal{Z}, \mathcal{D})/T_Y$. The fact that (Y, D) appears as a fibre is perhaps a bit surprising as, after all, we set out to construct the mirror and have obtained the original surface back. (Note however that dual Lagrangian torus fibrations in dimension 2 are topologically equivalent, so this is consistent with the SYZ formulation of mirror symmetry.) More importantly, the restrictions of the theta functions ϑ_q to $U \subset \mathcal{X}$ endow the affine surface $U = Y \setminus D$ with canonical functions.

We believe the existence of these canonical functions on such familiar objects (for example, the complement to a nodal plane cubic) represents the deepest reach to date of mirror symmetry into classical mainstream algebraic geometry.

To illustrate, in Example 5.11 we explicitly compute the theta functions in the case (Y, D) is the del Pezzo of degree 5 together with a cycle of 5 (-1)-curves. In Example 5.12, we give the expression in the case of a triangle of lines on a cubic surface, deferring in this case the proof until Part II. In each of these cases there is a characterisation of the ϑ_{v_i} in terms of classical geometry.

Both of these are examples of cluster varieties, and in each instance our theta functions agree with canonical Fock-Goncharov bases of universally positive Laurent polynomials, see [FG09]. Indeed we observe that all cluster algebras are log Calabi-Yau, and believe all Fock-Goncharov cluster ensembles (with their canonical bases) are instances of our Conjecture 0.8. We note that a cluster algebra comes with a great deal of auxiliary structure, but we believe that this structure is not necessary for the existence of Fock-Goncharov bases, it exists for any affine log Calabi Yau (with sufficiently degenerate boundary), and moreover that the many beautiful combinatorial rules for the multiplication rule of Fock-Goncharov bases admit a uniform expression in terms of counts of rational curves. For more details see §5.3.

We now discuss refinements of the general result of Theorem 0.2 in various special cases. For simplicity of exposition, assume that $\sigma_P = \mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}}$ is a rational polyhedral cone, so that $S := \operatorname{Spec} \Bbbk[P] = \operatorname{Spec} R$ is an affine toric variety (when this fails we approximate $\mathrm{NE}(Y)$ by P as above). Associated to a contraction $f : Y \to \bar{Y}$ is a closed boundary stratum of $S_f \subset S$, with monomial ideal I_f generated by monomials z^C for curves *not* contracted to a point by f. The most interesting cases for deformation theory turn out to be when D itself is a fibre of f. This occurs if D is negative semi-definite. In this case Theorem 0.2 does not apply directly; however, the same methods, with $\Bbbk = \mathbb{C}$, produce a canonical extension of our family over the formal completion $\widehat{S'}$ of an analytic open neighborhood S' of the zero-dimensional stratum of S along $S_f \cap S'$. See Theorem 4.1 for the precise statement in the case that D is negative definite, and Part II for the negative semi-definite case.

We denote by X_f the fibre over a general point of $S_f \cap S'$. Then X_f is a Gorenstein SLC surface, and the family $\mathcal{X}|_{S_f}$ is a partial smoothing of the vertex $\mathbb{V}_n$ to X_f. Furthermore, the family $\mathcal{X}$, restricted to a slice of S transverse to S_f, yields a formal smoothing of X_f. As remarked at the outset, every Gorenstein SLC singularity is a partial smoothing of some $\mathbb{V}_n$, and in fact every such singularity which admits a smoothing occurs as X_f for some f. We now discuss the two main examples.

The intersection matrix is negative semi-definite but not definite. In this case, after inductively contracting (-1)-curves contained in D, we may assume that D is a cycle of (-2)-curves or an irreducible rational nodal curve with $D^2 = 0$. We can choose the complex structure on Y so that D is a fibre of an elliptic fibration $f : Y \to \bar{Y}$ (recall our family depends only on the deformation type of (Y, D)). Then X_f has a unique singular point, a cone over an elliptic curve of degree n. The restriction of our family to the boundary stratum $\mathcal{X} \to S_f \cap S'$ is the cone over Mumford's construction of the Tate curve, and our theta functions restrict to his toric realization of classical theta functions for elliptic curves. Moreover our family $\mathcal{X} \to S$ is defined over the analytic open set $S' = \{x \in \operatorname{Spec} \Bbbk[P] \mid |z^D(x)| < 1\} \subset S$, and the modular parameter of the elliptic curve is $q = \exp(2\pi i \tau) = z^D \in \Bbbk[\mathrm{NE}(Y)]$. This in particular shows the family cannot be algebraic. Thus $\mathcal{X}$ gives a smoothing of the cone over an elliptic curve of degree n. We recover in this way a result of Pinkham:

Corollary 0.4. [P74, §9] *A cone over an elliptic curve of degree at most* 9 *is smoothable.*

More importantly (as there are much easier ways to smooth this singularity) our construction now gives a vast generalisation of Mumford's toric construction of theta functions for elliptic curves. Indeed, we will show in Part II that the fibrewise compactification of our family $(\mathcal{Z}, \mathcal{D}) \to \operatorname{Spf} \widehat{R}$, together with the theta functions (on $\mathcal{X} = \mathcal{Z} \setminus \mathcal{D}$) extends canonically over S'. The restriction to $S' \cap T_Y$ gives the universal family of pairs (Z, D) of a smooth anti-canonical elliptic curve on a del Pezzo surface, constructed in [L76] and [M82], but now endowed with canonical theta functions not previously observed.

The intersection matrix is negative definite. In this case, there is a birational morphism $f : Y \to \bar{Y}$ contracting D to a cusp singularity $p \in \bar{Y}$. See Example 1.9 for background on cusp singularities. Cusps come in natural dual pairs, as observed in [N80]. In fact, as we'll see in Example 1.9, this duality can be viewed as an early instance of mirror symmetry. We prove that the fibre X_f has a unique singularity, the dual cusp to $p \in \bar{Y}$, and we thus obtain (in Theorem 4.13) the following:

Corollary 0.5 (Looijenga's conjecture)**.** *A* 2*-dimensional cusp singularity is smoothable if and only if the exceptional cycle of the dual cusp occurs as an anti-canonical cycle on a smooth projective rational surface.*

This was conjectured by Looijenga in [L81], where he also proved the forward implication. Partial results were obtained in [FM83] and [FP84]. The above procedure produces a (formal) smoothing of the cusp singularity X_f, and we conjecture this dominates an irreducible component of the versal deformation space (the space is known to be equidimensional, and our base at least has the correct dimension).

A simpler version of our construction can be used to describe deformations of cyclic quotient singularities, see §6. In this case there is a complete theory due to Kollár–Shepherd-Barron [KSB88] and Stevens [S91]. However our approach gives a new perspective and in particular clarifies the connection with symplectic geometry discovered by Lisca [L08]. This is a baby version of the general construction, a nice introduction to the ideas of the paper. Our families in this case have natural modular meaning, see Theorem 6.6.

Our construction in general is based on ideas of Gross and Siebert in [GS07]. The relation is as follows. Suppose that $f : Y \to \bar{Y}$ is a *toric model*, i.e., f is the blowup of a smooth projective toric surface $\bar{Y}$ at a collection of smooth points of the toric boundary $\bar{D} \subset \bar{Y}$, and D is the strict transform of $\bar{D}$ (essentially all Looijenga pairs occur this way, see Proposition 1.19). In this case the restriction of $\mathcal{X}/S$ to S_f is the trivial family $\mathbb{V}_n \times S_f$. If $A \in \mathrm{Pic}(\bar{Y})$ is an ample divisor, then after pulling it back to Y, it determines a one-parameter subgroup of the torus $T_Y := \mathrm{Spec}\, \Bbbk[A_1(Y)]$. The closure of an orbit of this subgroup in $S = \mathrm{Spec}\, \Bbbk[P]$ intersects the open torus orbit of the stratum S_f. If we restrict the family $\mathcal{X}$ to this one-dimensional scheme, then we obtain the kind of smoothing constructed by Gross and Siebert in [GS07] in the two-dimensional case (closely related to the construction of [KS06]). We improve on the Gross–Siebert construction is several ways. The Gross–Siebert construction includes a number of choices. Incorporating those choices gives essentially the family over the formal neighbourhood of the interior of S_f. The main work for us is to extend the family over the closure. The base we obtain is a formal version of the Kähler moduli space of the pair (Y, D), the natural parameterizing space from the mirror symmetry perspective.

Most importantly, our construction comes with theta functions. In this paper they play a crucial technical role, in that they allow us to extend our deformations over parts of $\mathrm{Spec}\, \Bbbk[\mathrm{NE}(Y)]$ which the construction of Gross and Siebert cannot reach (which includes all the boundary strata that are interesting from the singularity point of view). Without the theta functions, the Gross–Siebert families do not carry very much information. We will see in Part II that they can in fact be constructed abstractly via a birational modification of the standard Mumford degenerations (in particular the Gross–Siebert families are formal completions of holomorphic families — a fact that is not at all apparent from their construction, which is purely formal, carried out order by order). We stress it is the abstract deformations that admit a simple description; the theta functions themselves are highly non-trivial. They are not at all apparent from the abstract description. For example, this birational description depends on a choice of toric model $Y \to \bar{Y}$, and a given Looijenga pair can have an infinite number of such models. On the other hand, the theta functions are independent of all choices, and thus are completely canonical.

We will also show in Part II that, when D is not negative definite, our family satisfies a natural universal property, and moreover we expect this holds even in the negative cases, generalizing Looijenga's family of good pairs [L81], II.2.7.

0.3. **Description of the coordinate ring.** We will now give a precise description of the algebra structure on A, and thus of our mirror family to (Y, D).

0.3.1. *The integral linear manifold* $B_{(Y,D)}$. If Y is a smooth projective toric surface, and D is the toric boundary, then Y is determined by a fan Σ in $B = \mathbb{R}^2$. Note the fan is determined (up to $\mathrm{SL}_2(\mathbb{Z})$) by the self-intersection numbers D_i^2. This allows us to use this same data to define an analogous object associated to any Looijenga pair.

In general, let $B = B_{(Y,D)}$ be the *dual complex* of (Y, D), i.e., the piecewise-linear surface obtained by gluing maximal cones $\sigma_{i,i+1} := (\mathbb{R}_{\geq 0})^2$, one for each singular point $D_i \cap D_{i+1}$ of D. The two edges of the cone $\sigma_{i,i+1}$ correspond to the divisors D_i and D_{i+1} respectively, and the cones $\sigma_{i-1,i}$ and $\sigma_{i,i+1}$ are glued along the edges corresponding to the divisor D_i. The resulting topological space B is homeomorphic to $\mathbb{R}^2$, and it comes along with a "fan" Σ consisting of the cones $\sigma_{i,i+1}$ and their faces. By construction B comes with a set of natural integral points $B(\mathbb{Z})$. As in the toric case, the rays of Σ correspond to the D_i, and each D_i determines an integral point $v_i \in B(\mathbb{Z})$, the first lattice point along its ray.

Furthermore, $B_0 := B \setminus \{0\}$ carries an integral affine structure — it has charts with transition functions in $\mathrm{SL}_2(\mathbb{Z})$. This structure is determined by declaring that the unique continuous piecewise linear function on $\sigma_{i-1,i} \cup \sigma_{i,i+1}$ with value a_i at v_i is linear if and only if

$$(a_{i-1}D_{i-1} + a_iD_i + a_{i+1}D_{i+1}) \cdot D_i = 0. \tag{0.2}$$

Let Λ denote the locally constant sheaf of integral tangent vectors, a locally constant sheaf of rank 2 lattices on B_0. For any convex set $\tau \subset B_0$ let Λ_τ indicate the stalk at a point in τ (any two are canonically identified by parallel transport). Now an integer point $q \in B_0(\mathbb{Z})$ is the same data as the ray $0 \in \mathbb{R}_{\geq 0} \cdot q \subset B$ with rational slope, together with a positive integer (expressing q as a multiple of the primitive integer vector on this ray). Just as in the toric case, a ray with rational slope is the same information as a choice of exceptional divisor on a *toric* blowup $Y' \to Y$, obtained by a weighted blowup of a node of D. Such exceptional divisors E are characterized by the property that Ω has a pole along E. Thus $B(\mathbb{Z})$ agrees with the earlier definition. For full details and examples see §1.1.

0.3.2. *Broken lines and multiplication.* Now we sketch a description of the canonical ring structure of Theorem 0.1. For the motivation behind the definition see §0.6.2. We continue with (Y, D) and (B, Σ) as above.

Let $C \subset V$ be a cone spanning the real vector space V. Say $a \in V$ is *above* $b \in V$ if $a = b + c$ for some $c \in C$. Now we say a continuous Σ-piecewise linear function $\varphi : B \to V$ is *C-convex* if everywhere locally the graph lies above the graphs of the linear extensions, as in the usual definition for $\mathbb{R}$-valued functions. Equivalently, a piecewise linear φ has a canonically defined *change of slope*, or bending parameter,

$p_{\rho_i,\varphi} \in V$, as we cross a ray $\rho_i \in \Sigma$, and φ is C-convex if and only if all bending parameters $p_{\rho_i,\varphi}$ lie in the cone C. The bending parameters determine φ uniquely, up to linear functions. For the precise definitions, see §1.2.

Now take $C = \mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}}$. We assume for simplicity of exposition that there exists a $\varphi : B \to A_1(Y,\mathbb{R})$ with bending parameters $p_{\rho_i,\varphi_i} = [D_i] \in P = \mathrm{NE}(Y)$. This is not the case in general, and instead one uses a *multi-valued piecewise linear function*, see §2.1, but the difference is largely notational.

Let $0 \neq v \in B(\mathbb{Z})$ be a primitive integer vector, $v = av_i + bv_{i+1}$, with $a \geq 0$, $b > 0$. As previously noted, the ray generated by v determines a weighted blowup $\pi\colon \tilde{Y} \to Y$ with irreducible exceptional divisor E mapping to $D_i \cap D_{i+1}$ (unless $a = 0$ when there is no blowup and $E = D_{i+1}$). We attach to the ray $\rho_v := \mathbb{R}_{\geq 0} \cdot v \subset B$ the formal sum

$$f_v := \exp\left(\sum_{\beta} k_\beta N_\beta z^{\overline{\beta}} \cdot (z^{(-v,-\varphi(v))})^{k_\beta}\right) \tag{0.3}$$

of monomials in $\Bbbk[\Lambda_v \oplus P^{\mathrm{gp}}]$. Here $\beta \in \mathrm{NE}(\tilde{Y})$, and N_β is a relative Gromov-Witten invariant roughly counting genus 0 curves in $\tilde{Y}$ of class β meeting E at a single point with contact order $k_\beta = \tilde{D} \cdot \beta$. Finally, $\pi_*\beta =: \overline{\beta} \in \mathrm{NE}(Y)$. We say $\overline{\beta}$ is an $\mathbb{A}^1$-*class* if $N_\beta \neq 0$, and a relative stable map contributing to N_β is called an $\mathbb{A}^1$-*curve*. The sum converges in the $\mathfrak{m}$-adic topology (recall $\mathfrak{m} \subset \Bbbk[P]$ denotes the torus invariant maximal ideal). We call

$$\mathfrak{D} := \{(\rho_v, f_v) \mid v \in B_0(\mathbb{Z}) \text{ primitive}\} \tag{0.4}$$

the *canonical scattering diagram.*

Definition 0.6 (Broken Line)**.** A *broken line* γ is a closed piecewise linear directed path in B_0 composed of finitely many straight line segments $L_1, L_2, \ldots, L_N$, with no segment contained in a ray of $\mathfrak{D}$ or Σ, each decorated by a monomial $m_i := c_i z^{(v_i,p_i)} \in \mathbb{Q}[\Lambda_{L_i} \oplus P^{\mathrm{gp}}]$ satisfying the following: γ can bend only where it crosses a ray of $\mathfrak{D}$. Each L_i is compact except for L_1, which is a ray that goes to infinity inside some two-dimensional cone $\sigma \in \Sigma$. Each v_i is non-zero and parallel to L_i, and points in the *opposite* direction to L_i. Also $m_1 = z^{(v_1,\varphi(v_1))}$ and (necessarily) $v_1 \in \sigma$. Subsequent decorations are determined inductively as follows: By assumption $L_i \cap L_{i+1}$ is on a ray $(\rho, f) \in \mathfrak{D}$. The monomial m_{i+1} is required to be one of the monomial terms in $m_i \cdot f^{\langle n,v_i\rangle}$ where $n \in \Lambda_v^*$ is the unique primitive element vanishing on the tangent space to ρ and positive on v_i.

One can show that $p_i - \varphi(v_i) \in P$ for all i. Let $c(\gamma), p(\gamma), v(\gamma)$ be c_N, p_N, v_N. Let $\mathrm{Limits}(\gamma) = (v_1, s)$, where $s \in B$ is the endpoint of the final segment L_N. Note

$s \in B_0(\mathbb{Z})$ determines an integer vector $v_s \in \Lambda_s$ (given in an affine chart taking $0 \in B$ to $0 \in \mathbb{R}^2$ by the vector from 0 to s).

We then have the following theorem, a summary of (parts of) Theorems 2.33, 2.38, and 3.8:

Theorem 0.7. *Let $I \subset \Bbbk[P]$ be a monomial ideal such that $\sqrt{I} = \mathfrak{m}$. For $p_1, p_2, s \in B(\mathbb{Z})$ define the formal sum*

$$\alpha_s(p_1,p_2) = \sum_{\substack{(\gamma_1,\gamma_2)\\ \mathrm{Limits}(\gamma_i)=(p_i,s)\\ v(\gamma_1)+v(\gamma_2)=s}} c(\gamma_1)c(\gamma_2)z^{p(\gamma_1)+p(\gamma_2)-\varphi(s)}.$$

For any pair (γ_1,γ_2) in the sum the exponent $p := p(\gamma_1)+p(\gamma_2)-\varphi(s)$ lies in P, and $z^p \in I$ for all but finitely many pairs, so the sum defines an element of $R_I := \Bbbk[P]/I$. The multiplication on generators of $A_I = \bigoplus_{s\in B(\mathbb{Z})} R_I \cdot \vartheta_s$ defined by

$$\vartheta_{p_1}\vartheta_{p_2} = \sum_{s\in B(\mathbb{Z})} \alpha_s(p_1,p_2)\vartheta_s$$

yields a well-defined R_I-algebra structure on A_I. The induced map $f_I : \mathrm{Spec}(A_I) \to \mathrm{Spec}(R_I)$ is the flat deformation of $\mathbb{V}_n$ of Theorem 0.1.

0.4. **Frobenius structure conjecture.** In this subsection we let (Y,D) be a smooth projective variety Y of arbitrary dimension together with an anti-canonical normal crossing divisor D (whereas elsewhere in the paper Y is a surface). Write $U := Y \setminus D$. Let Ω denote a nowhere zero top-dimensional holomorphic form on U with simple poles along D (uniquely determined up to scaling). We assume that D is connected and has a zero-dimensional stratum, that is, a point cut out by $\dim Y$ components of D.

Let S denote the dual complex of D. That is, let $D_1,\dots,D_n$ be the irreducible components of D. Assume for simplicity that each D_i is smooth and for each $1 \le i_1 < \cdots < i_p \le n$ the intersection $D_{i_1}\cap\cdots\cap D_{i_p}$ is connected (possibly empty). We can always reduce to this case by blowing up along boundary strata. Then S is the simplicial complex with vertices $v_1,\dots,v_n$ and simplices $\langle v_{i_1},\dots,v_{i_p}\rangle$ corresponding to non-empty intersections $D_{i_1}\cap\cdots\cap D_{i_p}$. Let B denote the cone over S and Σ the induced subdivision of B into simplicial cones. Let $B_0 := B\setminus\{0\}$ denote the complement of the vertex.

Now suppose we are given points $q_1,\dots,q_s \in B_0(\mathbb{Z})$. Each q_i can be written as a linear combination $q_i = \sum_j m_{ij}v_{ij}$ for primitive generators v_{ij} of rays in Σ, with the ray generated by v_{ij} corresponding to a divisor D_{ij}; necessarily $\bigcap_j D_{ij} \neq \emptyset$. Suppose also given a class $\beta \in A_1(Y)$ with the property that for a component F of D,

$$\beta \cdot F = \sum_{\substack{i,j \\ E = D_{ij}}} m_{ij}. \tag{0.5}$$

Using the results of [GS11] or [AC11], one can define a moduli stack $\overline{\mathcal{M}}_{0,s+1}(Y/D,\beta)$ of logarithmic stable maps, using the divisorial log structure on Y induced by D, which intuitively provides a compactification for the space of stable maps

$$g\colon (C, p_0, p_1, \ldots, p_s) \to Y$$

such that $g_*[C] = \beta$ and C meets D_{ij} at p_i with contact order m_{ij} for each i, j, $1 \le i \le s$, and order zero contact with D at p_0. (We note that we cannot use the more classical notion of relative stable map [LR01], [Li00] here, as D is not a smooth divisor). We observe that the stack $\overline{\mathcal{M}}_{0,s+1}(Y/D,\beta)$ has expected dimension $\dim Y + s - 2$. For $s \ge 2$ define the associated relative Gromov–Witten invariant

$$N_\beta(q_1, \ldots, q_s) := \int_{[\overline{\mathcal{M}}_{0,s+1}(Y/D,\beta)]^{vir}} \mathrm{ev}_0^*[\mathrm{pt}] \cdot \psi_0^{s-2}.$$

Informally, $N_\beta(q_1, \ldots, q_s)$ counts maps $g\colon (C, p_0, p_1, \ldots, p_s) \to Y$ such that C is a smooth rational curve, $g_*[C] = \beta$, C meets D_{ij} at p_i with contact order m_{ij} for $i = 1, \ldots, s$ and is otherwise disjoint from D, $g(p_0)$ is a fixed general point of Y, and the isomorphism type of the pointed curve $(C, p_0, p_1, \ldots, p_s)$ is fixed.

Assume that D supports an ample divisor. Then it follows from the Cone Theorem [KM98, 3.7] that the cone of curves $\mathrm{NE}(Y)_{\mathbb{R}_{\ge 0}} \subset A_1(Y, \mathbb{R})$ is rational polyhedral. Write $\mathrm{NE}(Y) := \mathrm{NE}(Y)_{\mathbb{R}_{\ge 0}} \cap A_1(Y, \mathbb{Z})$ for the associated monoid and $R := \Bbbk[\mathrm{NE}(Y)]$ for the associated finitely generated $\Bbbk$-algebra. For $\beta \in \mathrm{NE}(Y)$ we write $z^\beta \in R$ for the corresponding element. Let $A := \bigoplus_{q \in B(\mathbb{Z})} R \cdot \vartheta_q$ be the free R-module with basis $\{\vartheta_q\}$ for $q \in B(\mathbb{Z})$.

Define the R-multilinear symmetric *s-point functions*

$$\langle \bullet \rangle : (A)^{\times s} \to R,$$

for $s \ge 1$ as follows. For $s \ge 2$ and $q_1, \ldots, q_s \in B_0(\mathbb{Z})$, we define

$$\langle \vartheta_{q_1}, \ldots, \vartheta_{q_s} \rangle := \sum_{\beta \in \mathrm{NE}(Y)} N_\beta(q_1, \ldots, q_s) z^\beta \in R$$

where the sum is over classes β satisfying the condition (0.5). (Note that the ampleness assumption implies that the sum is finite.) For $s = 1$ we define $\langle \vartheta_0 \rangle = 1$ and $\langle \vartheta_q \rangle = 0$ for $q \in B_0(\mathbb{Z})$. Finally we define $\langle \vartheta_0, \vartheta_{q_1}, \ldots, \vartheta_{q_s} \rangle := \langle \vartheta_{q_1}, \ldots, \vartheta_{q_s} \rangle$ for $s \ge 1$.

Let $T_Y := \mathrm{Hom}(A_1(Y), \mathbb{G}_m)$ denote the big torus of the affine toric variety $\operatorname{Spec} R$.

Conjecture 0.8. *There is a unique finitely generated commutative and associative R-algebra structure on A with $\vartheta_0 = 1 \in A$ such that*

$$\langle a_1, a_2, \ldots, a_s \rangle = \langle a_1 \cdot a_2 \cdots a_s \rangle \in R.$$

That is, the s-point function $\langle a_1, \ldots, a_s \rangle$ is the coefficient of $\vartheta_0 = 1$ in the unique expansion of the product $a_1 \cdot a_2 \cdot \cdots \cdot a_s$ in terms of the R-module basis ϑ_q, $q \in B(\mathbb{Z})$.

The induced map $X := \operatorname{Spec}(A) \to S := \operatorname{Spec}(R)$ is a flat family of affine varieties with trivial relative dualizing sheaf $\omega_{X/S}$ and semi log canonical singularities. The fibers over $T_Y \subset S$ are irreducible and have log canonical singularities.

If $\Bbbk = \mathbb{C}$ the algebra A is the symplectic cohomology ring $SH^0(U)$.

Let $\mathbf{B} + i\omega$ be a complexified Kähler form on Y and $s = \exp(2\pi i(\mathbf{B} + i\omega)) \in T_Y \subset S$. Suppose $\Bbbk = \mathbb{C}$ and the fiber X_s is smooth. Then X_s is mirror to $(U = Y \setminus D, (\mathbf{B}+i\omega)|_U)$ in the sense of homological mirror symmetry. That is, the bounded derived category of X_s is equivalent to the wrapped Fukaya category of $(U, (\mathbf{B} + i\omega)|_U)$.

We can similarly formulate the conjecture without any ampleness assumption on D, using a polyhedral approximation σ_P to $\operatorname{NE}(Y)_{\mathbb{R}_{\geq 0}}$ and a monomial ideal I with $\sqrt{I} = \mathfrak{m}$ as in Theorem 0.1. We believe that in dimension 2 it would be relatively straightforward to prove the conjecture using Theorem 0.7 and tropical techniques. On the other hand, it would be far more interesting to prove that the prescription defines an associative R-algebra directly, without tropical mediation.

0.5. **Mirror symmetry conjectures.** We now discuss our expectation that our construction yields the mirror family to (Y, D) from a number of points of view. Assume $\Bbbk = \mathbb{C}$.

0.5.1. *Homological mirror symmetry.* Suppose that (Y, D) is not negative semi-definite. As discussed in §0.2, our construction gives a family $\mathcal{X}$ defined over the full toric base $S = \operatorname{Spec} \Bbbk[\operatorname{NE}(Y)]$.

Let $\mathbf{B} + i\omega$ be a complexified Kähler form on Y. Write $U = Y \setminus D$. Let $D^{\pi}\mathcal{F}_{\mathrm{wr}}(U)$ denote the wrapped Fukaya category of $(U, (\mathbf{B}+i\omega)|_U)$, [AS10] and [A09], §5.2. Let $s = \exp(2\pi i(\mathbf{B}+i\omega)) \in S$ denote the point corresponding to $[\mathbf{B}+i\omega] \in H^2(Y, \mathbb{C})/H^2(Y, \mathbb{Z})$. Let $X := \mathcal{X}_s$ denote the fiber of $\mathcal{X}/S$ over $s \in S$ and $D(X)$ the bounded derived category of coherent sheaves on X. Then, as an instance of homological mirror symmetry, we expect:

Conjecture 0.9. *There is an equivalence of triangulated categories*

$$D^{\pi}\mathcal{F}_{\mathrm{wr}}(U) \simeq D(X).$$

Remark 0.10. In [S08a] and [AS10] symplectic cohomology and the wrapped Fukaya category are defined only in the exact case. Also, they do not consider the **B**-field. Let $N \subset D$ be a tubular neighbourhood of D and $U' := Y \setminus D$. By our positivity assumption on D there is a vector field V defined in a neighbourhood of the boundary $\partial U'$ which points outwards along $\partial U'$ and satisfies $\mathcal{L}_V \omega = \omega$, or equivalently $d(\iota_V \omega) = \omega$. The vector field V is called a *Liouville vector field*. The existence of V suffices to define symplectic cohomology, cf. [Oa04], and [R09], §3. The same is expected for the wrapped Fukaya category.

The operations in symplectic cohomology and the morphisms in the Fukaya category are defined using counts of pseudoholomorphic curves $f \colon C \to U$ weighted by $\exp(2\pi i \int_C f^*(\mathbf{B} + i\omega))$. (In general one must work over a Novikov ring to avoid convergence issues, however we expect this is not necessary in our case because U is the minimal resolution of an affine surface, see Lemma 5.9.)

In the presence of a **B**-field the objects of the Fukaya category are generated by Lagrangians together with a complex vector bundle with unitary connection having curvature $-2\pi i \mathbf{B}$. See [AKO06], §4.

Remark 0.11. In general the fiber $X = \mathcal{X}_s$ has Du Val singularities (that is, quotient singularities $\mathbb{C}^2/G$ for $G \subset \mathrm{SL}(2, \mathbb{C})$ a finite subgroup). In this case we regard X as an orbifold and define $D(X)$ as the bounded derived category of coherent sheaves on the orbifold. Equivalently, we can replace X by its minimal resolution [BKR01].

Let $SH^*(U)$ denote the symplectic cohomology of $(U, \mathbf{B} + i\omega)$ [S08a],[R10]. There is a $\mathbb{C}$-algebra homomorphism (the "open-closed string map")

$$SH^*(U) \to HH^*(D^\pi \mathcal{F}_{\mathrm{wr}}(U))$$

from the symplectic cohomology of U to the Hochschild cohomology of its wrapped Fukaya category, which is conjectured to be an isomorphism (under slightly more restrictive conditions than considered here), see [S02], Conjecture 4. Since $HH^*(D(X)) = HH^*(X)$, $HH^0(X) = H^0(X, \mathcal{O}_X)$, and "Hochschild cohomology is an invariant of triangulated categories", if the open-closed string map is an isomorphism and Conjecture 0.9 holds, we find

Conjecture 0.12. *There is an isomorphism of* $\mathbb{C}$*-algebras*

$$SH^0(U) \simeq H^0(X, \mathcal{O}_X).$$

That is, the mirror X *of* $(U, \mathbf{B} + i\omega)$ *may be constructed as*

$$X := \operatorname{Spec} SH^0(U).$$

0.5.2. *The negative semi-definite case.* Suppose that $D \subset Y$ is a cycle of (-2)-curves. Replacing (Y, D) by a deformation equivalent pair, we may assume that the normal bundle of $D \subset Y$ is trivial. (Such pairs form a divisor in the moduli space.) Then Y admits a holomorphic elliptic fibration $w\colon Y \to \mathbb{P}^1$ such that $D = w^{-1}(\infty)$. Let $\mathbf{B}+i\omega$ be a complexified Kähler form on Y. Write $U = Y \setminus D = w^{-1}(\mathbb{C})$. Let $\mathcal{F}S(U, w)$ denote the Fukaya–Seidel category of vanishing cycles for the proper symplectic fibration $w\colon (U, (\mathbf{B} + i\omega)|_U) \to \mathbb{C}$, see [AKO06], [S09b].

Let $\mathcal{X}/S'$ be the mirror family constructed from (Y, D) as described in §0.2, with fibrewise compactification $(\mathcal{Z}, \mathcal{D})/S'$. Here $S' = \{|z^D| < 1\} \subset S$ is an analytic neighbourhood of the zero-dimensional torus orbit in $S = \operatorname{Spec} \Bbbk[P]$. Let $s = \exp(2\pi i(\mathbf{B} + i\omega)) \in S'$, and let $X := \mathcal{X}_s$, $(Z, E) := (\mathcal{Z}_s, \mathcal{D}_s)$ denote the fibers over s. Then Z is a del Pezzo surface, $E \subset Z$ is an anticanonical smooth elliptic curve, and $X = Z \setminus E$.

Conjecture 0.13. *There is an equivalence of triangulated categories*

$$\mathcal{F}S(U, w) \simeq D(Z).$$

Assuming Conjecture 0.20 below concerning the periods of the family $\mathcal{X}/S'$, Conjecture 0.13 is an easy consequence of the main theorem of [AKO06]. (One just has to check that the mirror map in [AKO06] agrees with the map $s = \exp(2\pi i(\mathbf{B}+i\omega))$ used here.)

Remark 0.14. In [AKO06] an equivalence of categories was established between the Fukaya–Seidel category of $(U, w, \mathbf{B} + i\omega)$ for $\mathbf{B} + i\omega$ a complexified Kähler form on U and the derived category of a noncommutative del Pezzo surface. In our approach the class $[\mathbf{B}+i\omega]$ lies in the image of $H^2(Y, \mathbb{C})$. These are precisely the cases for which the mirror is commutative.

One can define the wrapped Fukaya category $D^\pi\mathcal{F}_{\mathrm{wr}}(U, w)$ for $(U, w, \mathbf{B} + i\omega)$. Here to define the morphisms we use the Hamiltonian vector field associated to $H = h(|w|)$ where $h(t)$ grows sufficiently fast as $t \to \infty$. See [A09], §5.2 and [KKP08], §4.5. The wrapped Fukaya category $D^\pi\mathcal{F}_{\mathrm{wr}}(U, w)$ can be constructed from the Fukaya–Seidel category $\mathcal{F}S(U, w)$ by localization of a natural transformation

$$\sigma \to \mathrm{id},$$

where σ is determined by the monodromy of the fibration w at infinity and, according to a conjecture of Kontsevich, $\sigma[2]$ is the Serre functor of $\mathcal{F}S(U, w)$, see [S09a], §4. This is mirror to the construction of the derived category $D(X)$ from $D(Z)$ by localization of the natural transformation

$$(\cdot) \otimes K_Z \to \mathrm{id}$$

given by a defining equation for the boundary divisor $E \subset Z$. See [S08b], §1 and §6. So Conjecture 0.13 suggests the analogue of Conjecture 0.9:

Conjecture 0.15. *There is an equivalence of triangulated categories*

$$D^\pi \mathcal{F}_{\mathrm{wr}}(U, w) \simeq D(X)$$

0.5.3. *Y is a del Pezzo surface.* Now assume that Y is a del Pezzo surface, that is, $-K_Y = D$ is ample. Let $\vartheta_i := \vartheta_{v_i}$, $i = 1, \ldots, n$ be the global theta functions on $\mathcal{X}$ given by Theorem 0.1. Let $W := \vartheta_1 + \cdots + \vartheta_n \colon \mathcal{X} \to \mathbb{C}$, the *Landau–Ginzburg potential.*

Let $\mathbf{B} + i\omega$ be a complexified Kähler form on Y and $D^\pi \mathcal{F}(Y)$ the Fukaya category of $(Y, \mathbf{B} + i\omega)$. Let $s = \exp(2\pi i(\mathbf{B} + i\omega))$ and $X = \mathcal{X}_s$. Let $D_{\mathrm{Sing}}(X, W)$ be the derived category of singularities of $W \colon X \to \mathbb{C}$ [Or04].

Conjecture 0.16. *Assume $-K_Y$ is ample. Then $D^\pi \mathcal{F}(Y) \simeq D_{\mathrm{Sing}}(X, W)$.*

Remark 0.17. There is ongoing work of Auroux, Katzarkov, Orlov, and Pantev related to Conjecture 0.16 [AKO06],[P09].

Remark 0.18. If $-K_Y$ is not ample (more specifically, if there exist components D_i of the boundary D such that $-K_Y \cdot D_i \leq 0$) then the correct superpotential includes terms corresponding to stable maps of holomorphic disks containing components of the boundary in their image. See e.g. [A09], §3.2 for the toric examples $Y = \mathbb{F}_2, \mathbb{F}_3$. Our construction does not account for such maps.

Let $\mathrm{Sing}(W) \subset X$ denote the scheme theoretic singular locus of $W \colon X \to \mathbb{C}$, that is, the closed subscheme defined by $(dW = 0)$. Let $J(X, W) = \Gamma(\mathcal{O}_{\mathrm{Sing}\, W})$, the *Jacobian ring* of $W \colon X \to \mathbb{C}$. Assuming Conjecture 0.16, passing to Hochschild cohomology, assuming Kontsevich's conjecture [K95] identifying the Hochschild cohomology of the Fukaya category with the small quantum cohomology, and using the identification of the Hochschild cohomology of $D_{\mathrm{Sing}}(X, W)$ with the Jacobian ring [D09], we obtain

Conjecture 0.19. *Assume $-K_Y$ is ample. Then there is an isomorphism of $\mathbb{C}$-algebras*

$$J(X, W) \to QH^*(Y, \mathbf{B} + i\omega)$$

from the Jacobian ring of $W \colon X \to \mathbb{C}$ to the small quantum cohomology ring of $(Y, \mathbf{B} + i\omega)$, given by

$$\vartheta_i \mapsto [D_i].$$

In particular,

$$W \mapsto -[K_Y].$$

Conjecture 0.19 holds if Y is toric by [B93]. We have verified Conjecture 0.19 in the case (Y, D) is a cubic surface together with a triangle of (-1)-curves and $\mathbf{B} + i\omega$ is generic (joint work with A. Oblomkov).

0.5.4. *Periods.* We expect that the natural toric monomials on our base $S = \operatorname{Spec} \Bbbk[\mathrm{NE}(Y)]$ are *canonical coordinates* in the sense of mirror symmetry, i.e., given by periods of the natural holomorphic 2-form on the mirror family. In Part II we will prove this when D is not negative semi-definite.

Write

$$Q := \langle D_1, \ldots, D_n \rangle^{\perp} \subset H_2(Y, \mathbb{Z}).$$

We have an exact sequence

$$0 \to \mathbb{Z} \to H_2(U, \mathbb{Z}) \to Q \to 0.$$

The kernel is generated by the class γ of a real 2-torus in U which may be described explicitly as follows. Let $p \in D$ be a node, and identify the germ $(p \in Y, D)$ with $(0 \in \mathbb{C}^2_{z_1,z_2}, V(z_1 z_2))$. Then γ is the class of the compact torus $(|z_1| = |z_2| = \epsilon) \subset (\mathbb{C}^*)^2$ (the orientation is determined by the choice of cyclic ordering of the components of D).

The surface U admits a C^∞ real 2-torus fibration $f\colon U \to B$ with ordinary (Lefschetz) singularities over a disk B, with fiber class γ. (If $p\colon (Y, D) \to (\bar{Y}, \bar{D})$ is a blowdown to a toric surface together with its toric boundary then f can be obtained by modifying the smooth fibration of $\bar{Y} \setminus \bar{D} = (\mathbb{C}^*)^2$ given by the quotient by the compact torus $(S^1)^2 \subset (\mathbb{C}^*)^2$, cf. [S03]. We can always reduce to this case by Proposition 1.19.)

Assume that D is not negative definite, so that our formal family extends to a family $\mathcal{X}/S'$ over an open analytic subset S' of $S = \operatorname{Spec} \Bbbk[P]$. Let $S'^{o} \subset S'$ denote the locus of smooth fibers. One can show that a smooth fiber $\mathcal{X}_s$ of $\mathcal{X}/S'$ admits a dual torus fibration $\check{f}\colon \mathcal{X}_s \to B$. (This is a topological version of the SYZ description of mirror symmetry.) The torus fibration of $\mathcal{X}_s$ is obtained as a deformation of the (singular) torus fibration of the vertex $\mathbb{V}_n$ given by the quotient by the compact torus $(S^1)^2 \subset (\mathbb{C}^*)^2$ on each component $\mathbb{C}^2$. We let $\gamma_s \in H_2(\mathcal{X}_s, \mathbb{Z})$ denote the class of the torus fiber. (One can use the T^D action to show that the class γ_s is monodromy invariant.)

Since we are in dimension 2, we have

$$R^1 \check{f}_* \mathbb{Z} = (R^1 f_* \mathbb{Z})^* = R^1 f_* \mathbb{Z}.$$

using Poincaré duality on fibers of f. This gives an identification

$$H_2(\mathcal{X}_s, \mathbb{Z})/\langle \gamma_s \rangle = H_2(U, \mathbb{Z})/\langle \gamma \rangle = Q.$$

The dualizing sheaf $\omega_{\mathcal{X}/S}$ of the family $\mathcal{X}/S'$ is trivial, see Proposition 2.36. Moreover, as discussed in §0.2, the family has a fibrewise compactification $(\mathcal{Z}, \mathcal{D})/S'$ such that $\omega_{\mathcal{Z}/S}(\mathcal{D})$ is trivial. Let Ω be the global section of $\omega_{\mathcal{Z}/S}(\mathcal{D})$ such that $\int_{\gamma_s} \Omega = 1$

for each $s \in S'^o$. Thus the restriction of Ω to each smooth fibre $\mathcal{X}_s$ is a nowhere zero holomorphic 2-form with simple poles along $\mathcal{D}_s$, normalized so that $\int_{\gamma_s} \Omega = 1$. For $\beta \in H_2(Y, \mathbb{Z})$ we write $z^\beta \in \Bbbk[\mathrm{NE}\, Y]$ for the corresponding monomial.

Conjecture 0.20. *The local system*

$$S'^o \ni s \mapsto H_2(\mathcal{X}_s, \mathbb{Z})/\langle \gamma_s \rangle$$

has trivial monodromy.

Identify $H_2(\mathcal{X}_s, \mathbb{Z})/\langle \gamma_s \rangle = Q$ *as above. Then, for each* $\beta \in Q$*, writing* $\tilde{\beta} \in H_2(\mathcal{X}_s, \mathbb{Z})$ *for a lift of* $\beta \in Q$*, we have* $z^\beta = \exp(2\pi i \int_{\tilde{\beta}} \Omega)$.

The functions $\exp(2\pi i \int_{\tilde{\beta}} \Omega)$, $\beta \in Q$ are the so-called *canonical coordinates* on the complex moduli space of the surfaces $\mathcal{X}_s$ associated to the large complex structure limit $s \to 0 \in S'$.

Recall that the family $\mathcal{X}/S'$ is equivariant for the action of the torus T^D. We have the exact sequence

$$0 \to Q \to H_2(Y, \mathbb{Z}) \to \mathbb{Z}^n, \quad H_2(Y, \mathbb{Z}) \ni \beta \mapsto (\beta \cdot D_i)_{i=1}^n$$

and so, applying $\mathrm{Hom}(\cdot, \mathbb{C}^*)$, the exact sequence of tori

$$T^D \to T_Y \to \mathrm{Hom}(Q, \mathbb{C}^*) \to 0.$$

The torus $\mathrm{Hom}(Q, \mathbb{C}^*)$ is the period domain for the variation of mixed Hodge structure given by $H^2(\mathcal{X}_s)$, and local Torelli holds. See [L81], II.2.5 and [F83b]. So (assuming Conjecture 0.20) the family $\mathcal{X}/S'$ induces a versal deformation of each smooth fiber.

0.6. **Overview of the proof.**

We will now give a fairly detailed sketch of the proofs of Theorem 0.1 and Corollary 0.5. It may be helpful to read this section in conjunction with the body of the paper: we shall occasionally use notation introduced there.

0.6.1. *The SYZ picture of Mirror Symmetry.* Much of what we do in this paper, following the philosophy of the Gross–Siebert program, is to tropicalize the SYZ picture [SYZ96]. Thus it is helpful to review informally this picture in the context of mirrors to pairs (Y, D). The SYZ picture will be a heuristic philosophical guide, and hence we make no effort to be rigorous. Here we follow the exposition from [A07] concerning SYZ on the complement of an anti-canonical divisor, itself a generalization of ideas of Cho and Oh for interpreting the Landau-Ginzburg mirror of a toric variety in terms of counting holomorphic Maslov index two disks [CO06]. For the most part we follow Auroux's notation, except that we use Y instead of his X, and our X is his M.

Recall that (Y, D) is a rational surface together with D an anti-canonical cycle of rational curves of length n. We fix a symplectic form ω on Y, and a nowhere vanishing holomorphic 2-form Ω on $U := Y \setminus D$. Now suppose we have a fibration $f\colon U \to B$ by special Lagrangian 2-tori (i.e., a general fibre L of f satisfies $\operatorname{Im}\Omega|_L = \omega|_L = 0$). Then the SYZ mirror X of (U, ω) is the dual torus fibration $\check{f}\colon X \to B$. This can be thought of as a moduli space of pairs (L, ∇) consisting of a special Lagrangian torus L in U (a fibre of f) equipped with a unitary connection ∇ modulo gauge equivalence (or equivalently a holonomy map $\mathrm{hol}_\nabla : H_1(L, \mathbb{Z}) \to U(1) \subset \mathbb{C}^*$). The complex structure on X is subtle, specified by so-called instanton corrections.

In this picture we can define local holomorphic functions on X associated to a basis of $H_2(Y, L, \mathbb{Z})$ (in a neighbourhood of a fibre of $\check{f}$ corresponding to a non-singular fibre L of f) as follows. For $A \in H_2(Y, L, \mathbb{Z})$ define

$$z^A := \exp\left(-2\pi \int_A \omega\right) \mathrm{hol}_\nabla(\partial A) : X \to \mathbb{C}^*. \tag{0.6}$$

By choosing a splitting of $H_2(Y, L, \mathbb{Z}) \twoheadrightarrow H_1(L, \mathbb{Z})$ we can pick out local coordinates on X which define a complex structure. See [A07], Lemma 2.7. Note that as the fibre L varies, the relative homology group $H_2(Y, L, \mathbb{Z})$ forms a local system over $B_0 \subset B$, where B_0 is the subset of points with non-singular fibres. This local system has monodromy, and as a consequence, the functions z^A are only well defined locally.

However, there are also well defined global functions, $\vartheta_1, \ldots, \vartheta_n$ on X. These are defined locally in neighbourhoods of fibres of $\check{f}$ corresponding to fibres of f not bounding holomorphic disks in U, via a (rough) expression

$$\vartheta_i = \sum_{\beta \in H_2(Y,L,\mathbb{Z})} n_\beta z^\beta, \tag{0.7}$$

where n_β is a count of so-called Maslov index two disks with boundary on L representing the class β and intersecting D in one reduced point lying in D_i. (We note that in our setting the Maslov index μ of a holomorphic disk $f\colon \Delta \to Y$ with boundary lying on a special Lagrangian torus $L \subset Y$ is given by $\mu = 2 \deg f^* D$. See [A07], Lemma 3.1.) In the case that D is ample, there are, for generic L, only finitely many such disks; it is not known how to treat the general case in this symplectic setting.

For ϑ_i to make sense the moduli space of Maslov index 2 disks with boundary on L must deform smoothly with the Lagrangian L. This fails for Lagrangians that bound holomorphic disks contained in U (Maslov index zero disks). This is a codimension one condition on L, and thus defines canonical *walls* in the affine manifold B. When we cross the wall the ϑ_i are discontinuous. But the discontinuity is corrected by a holomorphic change of variable in the local coordinates z^β, according to [A07], Proposition 3.9:

$$z^\beta \to z^\beta \cdot h(z^\alpha)^{[\partial\beta]\cdot[\partial\alpha]} \tag{0.8}$$

where here $\alpha \in H_2(Y, L_0, \mathbb{Z})$ represents the class of the Maslov index zero disk with boundary on L_0 a Lagrangian fibre over a point on the wall, and $h(q)$ is a generating function counting such holomorphic disks. Thus we can define a new complex manifold, with the same local coordinates, by composing the *obvious gluing* induced by identifications of fibres of the local system on B_0 with fibres $H_2(Y, L, \mathbb{Z})$ with the automorphism (0.8). These regluings are the instanton corrections, and the modified manifold X should be the mirror. By construction it comes with canonical global holomorphic functions ϑ_i. In particular, the sum $W = \sum_i \vartheta_i$ is a well defined global function, the *Landau–Ginzburg potential*.

0.6.2. *The proofs.* Let us consider first the well-known case that (Y, D) is toric, so that $U := Y \setminus D$ is the structure torus $\mathbb{G}_m^2$. Then the SYZ fibration is smooth, and U does not contain any Maslov index zero disks bounding a fibre. Thus there are no instanton corrections. So in this case the naive description of X as a moduli space of Lagrangians with $U(1)$ connection is correct on the nose. There is a standard description of the mirror of a toric variety, due to Givental [Giv], which is really a simple instance of a fundamental construction of Mumford (with origins in his toric description of the Tate curve). Here the integral affine manifold $B = B_{(Y,D)}$ constructed in §0.3.1 is just $\mathbb{R}^2$ and Σ is the fan defining φ. Suppose we fix the symplectic form ω to represent the first Chern class of an ample line bundle on Y. This specifies, up to a linear function, $\varphi : B \to \mathbb{R}$ a strictly convex Σ-piecewise linear function with integral slopes. Let $\mathcal{Q} \subset \mathbb{P} := B \times \mathbb{R}$ be the set of points on or above the graph of φ. The polyhedron $\mathcal{Q}$ is preserved by translation by $\mathbb{R}_{\geq 0}$ acting on the second factor, and so the polarized toric variety $\mathcal{X}$ associated to $\mathcal{Q}$ (which in this case is affine as $\mathcal{Q}$ is in fact a cone) comes with a map $f : \mathcal{X} \to \mathbb{A}^1$. The special fibre is $\mathbb{V}_n$ and the other fibres are $(\mathbb{G}_m)^2$.

Let $v_i \in B$ be the primitive generator of the ray of Σ corresponding to $D_i \subset D$. Define ϑ_i to be the monomial $z^{(v_i, \varphi(v_i))}$. In the case that Y is Fano, it follows from the work of Cho and Oh [CO06] that the restriction of ϑ_i to the fibre $f^{-1}(1)$ coincides with the function defined in (0.7). If Y is not Fano, then (0.7) will contain some corrections from degenerate Maslov index two disks with components contained in the boundary.

This gives one family for each ample divisor, but it is more convenient to put these all together in one family, as Givental did, giving a single family $\mathcal{X} \to \mathrm{Spec}(\Bbbk[\mathrm{NE}(Y)])$, analogously defined in terms of a canonical (up to linear functions) strictly convex Σ-piecewise linear function

$$\varphi : B \to H_2(Y, \mathbb{R}).$$

Full details are given in §1.2

Now we look for an analogous family in the general case. We have already generalized (§0.3.1) the toric fan corresponding to the pair (Y, D) to the data (B, Σ), with

$B = B_{(Y,D)}$ integral linear with a singularity at the origin. The input for the Mumford construction, the canonical piecewise linear function φ, generalizes in an obvious way, and if we remove the singular point $0 \in \mathbb{V}_n$ there is a straightforward naive generalisation of Mumford's family, a (formal) deformation of $\mathbb{V}_n^o := \mathbb{V}_n \setminus \{0\}$ parameterized by $\operatorname{Spf}(\widehat{\Bbbk[\mathrm{NE}(Y)]})$, the formal completion of the toric variety $\operatorname{Spec} \Bbbk[\mathrm{NE}(Y)]$ at the unique zero-dimensional stratum of the boundary. (As usual in this introduction, we will assume that $\mathrm{NE}(Y)$ is a finitely generated monoid; more generally, we replace $\mathrm{NE}(Y)$ by a finitely generated monoid containing it). This naive deformation can be described via a simple open cover, as follows.

First note that $\mathbb{V}_n^o$ has a natural cover by open sets $U_i := \{x_i \neq 0\} \subset \mathbb{V}_n$ isomorphic to

$$U_i = V(x_{i-1}x_{i+1}) \subset \mathbb{A}^2_{x_{i-1},x_{i+1}} \times (\mathbb{G}_m)_{x_i}$$

which are disjoint except for

$$U_{i,i+1} := U_i \cap U_{i+1} = (\mathbb{G}_m)^2_{x_i,x_{i+1}}.$$

In $\mathbb{V}_n$ they are glued in the obvious way, i.e., via the canonical inclusions

$$U_{i,i+1} = \{x_{i+1} \neq 0\} \subset U_i, \quad U_{i,i+1} = \{x_i \neq 0\} \subset U_{i+1}.$$

The formal deformation is obtained by gluing thickenings of the U_i

$$U_{i,I} := \{x_{i-1}x_{i+1} = z^{[D_i]}x_i^{-D_i^2}\} \subset S_I \times \mathbb{A}^2_{x_{i-1},x_{i+1}} \times (\mathbb{G}_m)_{x_i} \tag{0.9}$$

where $z^{[D_i]} \in \Bbbk[\mathrm{NE}(Y)]$ is the corresponding monomial. Here $I \subset \Bbbk[\mathrm{NE}(Y)]$ is a monomial ideal with $\sqrt{I} = \mathfrak{m}$ the maximal monomial ideal, and $S_I = \operatorname{Spec}(\Bbbk[\mathrm{NE}(Y)]/I)$. The overlaps are relative tori, $U_{i,i+1,I} = S_I \times \mathbb{G}_m^2$, and the gluings are the obvious ones. The details (including a coordinate free description) are given in §2.1. This naive generalization is directly analogous to the naive description of (an open subset of) the mirror as the moduli of smooth special Lagrangian fibres with $U(1)$ connection.

Next, there is a natural way of translating the instanton corrections. The analog of the scattering walls and the attached generating functions for associated holomorphic disks in U is our canonical scattering diagram, see §0.3.2 and §2.2. As in [GPS09], one is able to replace the non-algebro-geometric notion of holomorphic disk with, roughly speaking, finite maps $\mathbb{A}^1 \to U$. Intuitively, each holomorphic disk contributing can be approximated by a global rational curve (meeting D in a single point).

This intuition leads to the scattering diagram $\mathfrak{D}$ of (0.4). The scattering diagram gives the set of walls described in the symplectic setting in §0.6.1. In a sense, our picture can be viewed as an asymptotic version of the chamber structure which would appear in the symplectic setting. For example, B has only a single singularity at the origin, but the more natural picture of having a base of the SYZ fibration with many simple singularities can be recovered by perturbing the single singularity of B. Our scattering diagram $\mathfrak{D}$ can then be viewed as a limit of diagrams, where each diagram is a union of *tropical disks* with leaves at the simple singularities. The diagram $\mathfrak{D}$ can then be viewed as a natural limit of diagrams defined tropically.

We now modify our naive generalization of Mumford's family by exactly the same coordinate transformations (0.8). This procedure is carried out for a general scattering diagram, in §2.2. In this way we obtain a modified family X_I^o (referred to as $X_{I,\mathfrak{D}}^o$ in the main body of the paper, where $\mathfrak{D}$ is the scattering diagram). This is a flat deformation of $\mathbb{V}_n^o$, the punctured n-vertex, over $\operatorname{Spec} \Bbbk[\mathrm{NE}(Y)]/I$. Now we want to extend this modified family to a flat deformation of $\mathbb{V}_n$. By some standard commutative algebra (see Lemma 2.34) it is enough to find lifts to X_I^o of the canonical coordinate functions x_i on $\mathbb{V}_n$. Then these coordinate functions give embeddings $X_I^o \hookrightarrow \mathbb{A}^n \times S_I$, and we obtain X_I by taking the closure in $\mathbb{A}^n \times S_I$.

There is in fact a canonical choice of lift, suggested by the symplectic heuristic. The natural global functions in the symplectic heuristic are the functions ϑ_i of (0.7). This admits a natural translation into tropical geometry. We use the notion of broken line, introduced in [G09] to construct the full Landau-Ginzburg potential for the mirror of $\mathbb{P}^2$.

A rough definition of broken line has already been given in Definition 0.6; see Definition 2.22 for a precise definition. The underlying piecewise straight path represents a *tropical Maslov index two disk.* While the path itself is not a tropical curve (there is no balancing condition at the bends), we can produce one by, for each bending point b of the broken line, gluing on the line segment $\overline{0b}$. Recall any bending point lies on a ray of the scattering diagram — this ray represents the tropicalisation of an $\mathbb{A}^1$-curve. Heuristically we imagine that a subset of this $\mathbb{A}^1$-curve is a Maslov index zero holomorphic disk with boundary on the SYZ fibre L_b. The line segment $\overline{0b} \subset B$ represents the tropicalisation of this disk. Having glued on these line segments, the bending in the definition of broken line yields the usual tropical balancing condition at vertices, and the resulting tree is viewed as the tropicalization of a Maslov index two holomorphic disk with boundary lying in the SYZ fibre over the endpoint of the broken line. This Maslov index two disk is approximated by taking a cylinder over each line segment of the broken line, and gluing on the appropriate disks at each bend. The attached data p_i of Definition 0.6 (or equivalently q_L of Definition 2.22) is analogous to the relative homology class $\beta \in H_2(Y, L)$. The transformation rule for p_i when the path bends is analogous to the change in this class coming from gluing on Maslov index zero disks. The coefficients keep track of the actual count of such disks; this count can be viewed as a generalization of the usual Mikhalkin tropical multiplicity formula [Mk05].

Our theta functions are defined in local charts as sums over monomials attached to broken lines: this is the expression $\mathrm{Lift}_Q(q)$ of Equation (2.13), and is exactly analogous to the canonical global functions ϑ_i of (0.7). Now to complete the construction we have to prove that these local expression *patch*, i.e., transform under change of local variables according to the wall-crossing maps determined by the canonical scattering diagram $\mathfrak{D}$. If this is the case, we say that $\mathfrak{D}$ is *consistent*, see Definition 2.32. Of course this is a completely natural expectation — our scattering diagram is motivated by the scattering walls of the heuristic discussion, and the associated automorphisms are exactly defined so that the local expressions for the ϑ_i patch together. But this is only heuristic reasoning, and our actual proof of consistency proceeds along quite different lines.

We assume now for simplicity that we have a *toric model* $Y \to \bar{Y}$. The broken lines of [G09] were defined in a case where the integral affine manifold B is smooth. In our situation, $B_{(\bar{Y},\bar{D})} = \mathbb{R}^2$ has no singularities (in fact smoothness of B exactly characterizes toric pairs, see Lemma 1.3). The analogous consistency result was proven there for a scattering diagram which had a natural compatibility. This compatibility stated that the composition of scattering automorphisms for a small loop around any point (see Definition 3.23 for the precise notion of path-ordered compositions) is trivial. A cleaner proof of this consistency was given in [CPS], not using the specifics of the $\mathbb{P}^2$ situation used in [G09]. Our strategy is to reduce our consistency question on $B_{(Y,D)}$ to an analogous question on $B_{(\bar{Y},\bar{D})}$, and then apply the method of [CPS].

The existence of the toric model $p : Y \to \bar{Y}$ allows us to define the Gross–Siebert locus: let G be the monomial ideal in $\Bbbk[\mathrm{NE}(Y)]$ generated by classes which are not contracted by p. Then G defines a toric boundary stratum of $\mathrm{Spec}(\Bbbk[\mathrm{NE}(Y)])$; the big torus orbit T of this stratum is what we call the *Gross–Siebert locus.* If J is a monomial ideal with $\sqrt{J} = G$, we obtain a thickening $T_J \subset S_J$ of the Gross–Siebert locus, and then will construct an extension of the family $\mathbb{V}_n \times T$ to a flat family over T_J. We call this the Gross–Siebert family.

This is related to the construction of [GS07] as follows. A divisor $L = p^*A$ for A an ample divisor on $\bar{Y}$ induces a map $\mathbb{A}^1 \to \mathrm{Spec}\, \Bbbk[\mathrm{NE}(Y)]$ with $0 \in \mathbb{A}^1$ mapping into T. Pulling back the family we construct to the completion of $\mathbb{A}^1$ at 0 gives a family constructed using the techniques of [GS07].

We construct the Gross–Siebert family as follows. Restricting [GS07] to the case at hand, [GS07] applies only to affine surfaces with very specific types of singularities, and *not* of the type that $B_{(Y,D)}$ has at the origin. Rather, all singularities must appear in the interiors of rays of the fan on B, and must have monodromy of the form $\begin{pmatrix} 1 & -k \\ 0 & 1 \end{pmatrix}$

for some $k > 0$, with the ray being the invariant direction; in analogy with the Kodaira classification, we call this an I_k singularity. Indeed, one expects a cycle of k two-spheres as fibre over such a point in the SYZ picture.

Here, we can view such a surface as being obtained by *factoring* the complicated singularity $0 \in B$ into I_1 singularities along the edges of Σ, one for each exceptional divisor of $p : Y \to \bar{Y}$, by *moving worms* (in the terminology of [KS06]). We describe the process in Example 1.10. Let B' be the manifold so obtained, choosing arbitrarily the position of each I_1 singularity along its corresponding edge. If we now push the I_1 singularities to infinity (along their invariant directions, which correspond to the rays of the toric fan for $\bar{Y}$) we obtain the toric fan $\Sigma_{\bar{Y}}$ for $\bar{Y}$ in $B_{(\bar{Y},\bar{D})} = \mathbb{R}^2$. The Gross–Siebert deformations are built using a scattering diagram, $\mathfrak{D}'$, constructed by translating one of the key ideas from [KS06]. In particular, there is an algorithm for the construction of $\mathfrak{D}'$, which we now outline.

Let us explain the initial input to the algorithm. If B' had no singularities, then $\mathfrak{D}'$ is empty and one obtains a Mumford deformation in a straightforward way. Locally near a ray ρ of B', one can view this deformation infinitesimally simply by gluing together two standard thickenings of $\mathbb{A}^1 \times \mathbb{G}_m$ to obtain the set $U_{i,J}$ given in (0.9). However, if instead B' is singular, say with k I_1 singularities on a ray of B', then while the thickenings of the irreducible components are canonical (still the irreducible components of $U_{i,J}$), the gluings are not, as the gluing depends on a path chosen connecting the two maximal cells of B' containing the given edge: see [GS08], §2.3 for a detailed discussion of this phenomenon. In particular, the choice of path gives different choices for $U_{i,J}$: if one glues using a path which crosses ρ very far from the origin, so that all the singularities lie between this crossing point and the origin, then the naturally induced gluing yields the choice (0.9) for the glued patch. On the other hand, if one glues using a path which crosses ρ near the origin, so that there are no singularities between this crossing point and the origin, then one obtains a chart

$$U'_{i,J} := \{x_{i-1}x_{i+1} = z^{[D_i]}x_i^{-D_i^2-k}\} \subset S_J \times \mathbb{A}^2_{x_{i-1},x_{i+1}} \times (\mathbb{G}_m)_{x_i}.$$

This ambiguity needs to be dealt with. The suggestion originally made by Kontsevich and Soibelman in [KS06] was to let rays emanate from the singularities. These rays carry gluing information in the form of a function which determines an automorphism. Then when we try to glue two affine pieces together, we need to modify the standard gluing using these automorphisms. In our case, each singularity has two rays emanating from it. Because the automorphisms associated to these can rays differ, it is possible to choose these automorphisms in such a way that the effect of monodromy about the singularity is cancelled out.

In our case, these modified gluings yield a modification of the $U_{i,J}$ of (0.9):

$$U_{i,J} = V(x_{i-1}x_{i+1} - z^{[D_i]}x_i^{-D_i^2}\prod_j(1+z^{[E_{ij}]}x_i^{-1})) \subset S_J \times \mathbb{A}^2_{x_{i-1},x_{i+1}} \times (\mathbb{G}_m)_{x_i}. \tag{0.10}$$

Here E_{ij} are the exceptional divisors ((-1)-curves) for the corresponding toric model $p : Y \to \bar{Y}$ which intersect the boundary divisor $D_i \subset D$. The basic problem in this approach is that in order to ensure compatible gluings, these rays need to extend indefinitely. When rays collide, one obtains new compatibility problems because the associated automorphisms in general fail to commute. This fundamental problem was solved in [KS06], where a canonical procedure for adding new rays to restore compatibility is introduced. This key idea is stated in Theorem 3.24. The Gross–Siebert deformation is obtained using the diagram $\mathfrak{D}'$ which is the output of this procedure applied to $\mathfrak{D}'_0$, the scattering diagram consisting of the rays emanating from the singularities of B'. In fact in our argument, we never work on B', but rather imagine the singularities go off to infinity, yielding the smooth manifold $B_{(\bar{Y},\bar{D})} = \mathbb{R}^2$. The limit of $\mathfrak{D}'_0$ (which we call $\bar{\mathfrak{D}}_0$) on $B_{(\bar{Y},\bar{D})}$ has only the rays emanating from the singularities (now at infinity) heading towards the origin. These rays now come in from infinity.

The diagram $\mathfrak{D}'$ satisfies the necessary compatibility to build an actual deformation, but its role for mirror symmetry was unclear. The meaning of $\mathfrak{D}'$ was clarified by [GPS09]. The main result of that paper is a description of $\mathfrak{D}'$ in terms of relative Gromov-Witten invariants. In particular, $\mathfrak{D}'$ can be constructed by counting rational curves on a toric variety $(\bar{Y},\bar{D})$ meeting the toric boundary at a collection of prescribed points, plus a single extra contact at an unspecified point. More precisely, these invariants involve counting curves on the blow-up of $\bar{Y}$ at the prescribed points, and these curves are precisely the $\mathbb{A}^1$-curves mentioned earlier. Thus the repackaged result of the Kontevich-Soibelman scattering process is our canonical scattering diagram $\mathfrak{D}$. This is the translation into algebraic geometry of the collections of walls and generating functions determined by Maslov index zero disks. Thus starting with the simple input diagram the Kontsevich-Soibelman procedure produces the diagram one expects from the symplectic heuristic.

Given this, we prove consistency of $\mathfrak{D}$, which is a question of equality of functions defined to arbitrary order, after restricting to the Gross–Siebert locus. If we move the singularities to infinity, $\mathfrak{D}'$ induces a diagram $\bar{\mathfrak{D}}$ on $\mathbb{R}^2 = B_{(\bar{Y},\bar{D})}$, which, by construction, satisfies the local consistency condition required for patching. Now we reduce the patching question for the canonical diagram on $B_{(Y,D)}$ to $\bar{\mathfrak{D}}$ on $B_{(\bar{Y},\bar{D})}$. The identity map on the underlying topological spaces gives $\mathbb{Z}$PL identifications $\nu : B \to B' \to \bar{B}$. This allows us to compare the scattering diagrams, $\mathfrak{D}$, $\mathfrak{D}'$ and $\bar{\mathfrak{D}}$ (which lie on different affine manifolds). The scattering diagrams, or more precisely, the functions attached to rays, *are not* identified by ν, see Proposition 3.26. This is clear since the starting diagram $\mathfrak{D}'_0$ changes as we move worms. Nonetheless ν induces a bijection on broken lines, see Lemma 3.30.

This bijection between broken lines allows us to reduce consistency for $\mathfrak{D}$ on $B_{(Y,D)}$ to consistency for $\bar{\mathfrak{D}}$ on $B_{(\bar{Y},\bar{D})}$, and so to the method of [CPS]. This completes the construction of the family in Theorem 0.1.

Remark 0.21. We take this as an indication that it is the collection of broken lines, rather than the scattering diagram, which is the essential object. We will show in a future paper [K3] that in the global K3 context the analog of broken lines are similarly independent of moving worms. Here the invariance is more striking, as the underlying scattering diagrams can be quite wildly different. This is the key fact that will allow us to glue together deformations of the global analog of the n-vertex constructed from different semi-stable models for the mirror degeneration, to produce a canonical deformation, with accompanying theta functions, depending only on the generic fibre of the mirror degeneration.

We conclude this overview by explaining how Looijenga's conjecture follows naturally from the construction of the mirror family. The point is that we can extend our construction (from the formal completion over the maximal ideal) to give a family over the formal completion along *most* of the toric boundary. In particular, let $f : Y \to Y'$ be the contraction of the cusp, and let $I_f \subset \Bbbk[\mathrm{NE}(Y)]$ be the ideal generated by the monomials z^C for curves C not contracted by f. Let T_2 be the corresponding stratum of $\operatorname{Spec}\mathbb{C}[\mathrm{NE}(Y)]$, and assuming there is a toric model $p : Y \to \bar{Y}$, let T_1 be the closure of the Gross–Siebert locus. We extend our family over the formal completion along $(T_1 \cup T_2) \cap S'$ for an open analytic neighbourhood S' of the torus fixed point of $\operatorname{Spec}\mathbb{C}[\mathrm{NE}(Y)]$. Informally, the reader should imagine that we have a holomorphic family defined in a tubular neighborhood of the union of these two strata.

From the explicit charts for smoothings around the Gross–Siebert locus, one can show that this family contains formal smoothings. This can be done because these deformations can be described via local charts that cover all of $\mathbb{V}_n$. Indeed, the complement of the origin is covered by the charts given by (0.10), while a neighbourhood of the origin in fact is described purely torically and is locally isomorphic to a Mumford degeneration. From these explicit charts it is obvious that these are smoothings. Thus by considering the Gross–Siebert locus, we conclude the "generic fibre" of our family is smooth. (Here we use quotes because, as our family is only formal, we do not actually have a generic fibre, but there is a natural formulation, in terms of the scheme-theoretic singular locus of the map, which makes sense). See §4.3.

Next we consider the family over the locus T_2, defined by the ideal I_f. From the naive charts above it is easy to guess the singularities of the fibre of our family over a generic point of the boundary stratum T_2. The local charts (0.9) covering $\mathbb{V}_n^o$ are correct, and we only modify the gluing. Since the monomials z^{D_i} do not lie in the ideal I_f, they are in fact invertible generically along T_2. Thus from (0.9) we see that the double curves of $\mathbb{V}_n^o$ are smoothed, and so the fibre has at worst normal Gorenstein SLC singularities. Furthermore, our scattering diagram is defined in terms of $\mathbb{A}^1$-classes — none of these are contracted by f, and thus modulo I_f, the scattering automorphisms are trivial. One can then show that our family over the open torus orbit of T_2 (intersected with S') specializes to Hirzebruch's original quotient construction for the dual cusp to the cusp of Y'. See Example 1.9. We carry this out in §4. We will see this part of our family — a partial smoothing of $\mathbb{V}_n$ to the dual cusp — is obtained from the purely toric Mumford construction, modulo a natural $\mathbb{Z}$-action. As we have already seen our "generic fibre" is smooth, so in particular the cusp is smoothable. If our family were holomorphic, defined over a tubular neighborhood of the toric boundary, this implication would be obvious. Because we have only formal families the details are a bit more delicate, see §4.3.

Similar constructions to ours appeared in the physics papers [GMN1], [GMN2], and [GMN3], which are in part concerned with a construction of a family of complex manifolds over $\mathbb{P}^1$, the twistor family of some underlying hyperkahler manifold. The construction depends on some additional data which is natural from the point of view of supersymmetric field theory, but is not a priori available in our context; nevertheless it turns out that some of the spaces we discuss also appeared in these papers In these cases: The restriction of the main construction of [GMN1] to any single fiber of the $\mathbb{P}^1$ family (excluding the fibers over 0 and ∞) is equivalent to the gluing we use. Canonical functions identical (we believe) to our theta functions also appeared in [GMN3], as line operators in a certain supersymmetric four-dimensional field theory. In [GMN2], where the hyperkahler manifold in question is Hitchen's integrable system, the line operators are traces of holonomies around loops as in the Fock-Goncharov theory (which [GMN2] in part follows).

0.7. **Acknowledgements.** An initial (and ongoing) motivation for the project was to find a geometric compactification of moduli of polarized K3 surfaces. We received a good deal of initial inspiration in this direction from conversations with V. Alexeev. We learned a great many things from A. Neitzke, especially about the connections of our work with cluster algebras and moduli of local systems. Our thinking about Looijenga pairs was heavily influenced by conversations with R. Friedman and E. Looijenga.

Our Conjecture 0.12 was inspired by talks with M. Abouzaid, D. Auroux and P. Seidel. Many other people have helped us with the project, discussions with D. Allcock, D. Benzvi, V. Fock, D. Freed, A. Goncharov, R. Heitmann, D. Huybrechts, M. Kontsevich, A. Oblomkov, T. Perutz, M. Reid, A. Ritter, B. Siebert, and Y. Soibelman were particularly helpful. We would also like to thank IHÉS for hospitality during the summer of 2009 when part of this research was done. The first author was partially supported by NSF grants DMS-0805328 and DMS-0854987. The second author was partially supported by NSF grant DMS-0968824. The third author was partially supported by NSF grant DMS-0854747.

1. Basics

1.1. **Tropical Looijenga pairs.** We recall the following basic definition. Fix a lattice $M \cong \mathbb{Z}^n$. In what follows, we will always use the notation $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$, $N = \operatorname{Hom}_{\mathbb{Z}}(M, \mathbb{Z})$ and $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$. We denote by $\operatorname{Aff}(M)$ the group of affine linear transformations of the lattice M.

Definition 1.1. An *integral affine manifold* B is a (real) manifold B with an atlas of charts $\{\psi_i : U_i \to M_{\mathbb{R}}\}$ such that $\psi_i \circ \psi_j^{-1} \in \operatorname{Aff}(M)$ for all i, j.

An *integral affine manifold with singularities* B is a (real) manifold B with an open subset $B_0 \subset B$ which carries the structure of an integral affine manifold, and such that $\Delta := B \setminus B_0$, the *singular locus* of B, is a locally finite union of locally closed submanifolds of codimension at least two.

An integral affine manifold with singularities is *(oriented) integral linear* if the transition maps of the atlas lie in $SL_n(\mathbb{Z})$.

If B is an integral affine manifold with singularities, there is a local system Λ_B on B_0 consisting of flat integral vector fields: i.e., if $y_1, \ldots, y_n$ are local integral affine coordinates, then Λ_B is locally given by linear combinations of the vector fields $\partial/\partial y_1, \ldots, \partial/\partial y_n$. If B is clear from context, we drop the subscript B.

Similarly, $\check{\Lambda}_B$ is the dual local system, locally generated by $dy_1, \ldots, dy_n$.

We will be primarily interested in $\dim B = 2$ in this paper, in which case Δ will consist, in all our examples, of a finite number of points. All integral affine manifolds we encounter will in fact be oriented integral linear.

The fundamental example for this paper is the following. Fix a pair (Y, D) where Y is a rational surface and $D = D_1 + \cdots + D_n \in |-K_Y|$ is an anti-canonical cycle of rational curves. We associate to this data a pair (B, Σ), where B is homeomorphic to $\mathbb{R}^2$ with singularity at the origin, and Σ is a decomposition of B into cones. The idea is that we pretend that (Y, D) is toric and we try to build the associated fan. More precisely, the construction is as follows.

We will always take $D_1, \ldots, D_n$ cyclically ordered and indices modulo n, so that $D_i \cap D_j \neq \emptyset$ if and only if the indices i and j are adjacent in the cyclic ordering of $1, \ldots, n$. For each node $p_{i,i+1} := D_i \cap D_{i+1}$ of D we take a rank two lattice $M_{i,i+1}$ with basis v_i, v_{i+1}, and the cone $\sigma_{i,i+1} \subset M_{i,i+1} \otimes_{\mathbb{Z}} \mathbb{R}$ generated by v_i and v_{i+1}. We then glue $\sigma_{i,i+1}$ to $\sigma_{i-1,i}$ along the rays $\rho_i := \mathbb{R}_{\geq 0} v_i$ to obtain a piecewise-linear manifold B homeomorphic to $\mathbb{R}^2$ and a decomposition

$$\Sigma = \{\sigma_{i,i+1} \,|\, 1 \leq i \leq n\} \cup \{\rho_i \,|\, 1 \leq i \leq n\} \cup \{0\}.$$

We define an integral affine structure on $B \setminus \{0\}$. We do this by defining charts $\psi_i : U_i \to M_{\mathbb{R}}$ (where $M = \mathbb{Z}^2$). Here

$$U_i = \mathrm{Int}(\sigma_{i-1,i} \cup \sigma_{i,i+1})$$

and ψ_i is defined on the closure of U_i by

$$\psi_i(v_{i-1}) = (1, 0), \quad \psi_i(v_i) = (0, 1), \quad \psi_i(v_{i+1}) = (-1, -D_i^2),$$

with ψ_i linear on $\sigma_{i-1,i}$ and $\sigma_{i,i+1}$.

A coordinate free description of this structure is given by equation (0.2) in the introduction.

We note this construction makes sense even when $n = 1$, i.e., the anti-canonical divisor D is an irreducible nodal curve. In this case there is one cone $\sigma_{1,1}$, and opposite sides of the cone are identified. (Moreover, the integral affine charts are defined using the integer $D_1^2 - 2$ instead of D_1^2. This is the degree of the normal bundle of the map from the normalization of D_1 to Y.) However, this case will often complicate arguments in this paper, so we will usually replace Y with a surface obtained by blowing up the node of D, and replace D with the reduced inverse image of D under the blowup. This does not change the underlying integral affine manifold with singularities, but refines the decomposition Σ, exactly as in the toric case. See §1.3 for some elementary properties of such *toric blowups*.

Example 1.2. It is easy to see that if Y is a non-singular toric surface and $D = \partial Y$ is the toric boundary of D, then in fact the affine structure on B extends across the origin, identifying (B, Σ) with $(M_{\mathbb{R}}, \Sigma_Y)$, where Σ_Y is the fan for Y. Indeed, if $\rho_j \in \Sigma_Y$ corresponds to the divisor D_j and $\rho_j = \mathbb{R}_{\geq 0} w_j$ with $w_j \in M$ primitive, then it is a standard fact that

$$w_{i-1} + (D_i)^2 w_i + w_{i+1} = 0.$$

Since Y is non-singular, there is always a linear identification $\varphi_i : M \xrightarrow{\sim} \mathbb{Z}^2$ taking w_{i-1} to $(1, 0)$, w_i to $(0, 1)$, and thus w_{i+1} must map to $(-1, -D_i^2)$. Thus on U_i, a chart for the affine structure on B is $\psi_i' = \varphi_i^{-1} \circ \psi_i : U_i \to M_{\mathbb{R}}$. The maps ψ_i' glue to give an integral affine isomorphism $B \to M_{\mathbb{R}}$.

In fact, the converse is also true:

Lemma 1.3. *If the affine structure on* $B_0 = B \setminus \{0\}$ *extends across the origin, then* Y *is toric and* $D = \partial Y$*.*

Proof. We make use of some elementary results which are proved later in the paper. We first note that by Lemma 1.17, we can replace (Y, D) with a non-singular toric blow-up without affecting the affine manifold B. By Proposition 1.19, we can thus assume that there is a smooth toric variety $\bar{Y}$ with toric boundary $\bar{D}$ along with a birational morphism $\pi : (Y, D) \to (\bar{Y}, \bar{D})$ which induces an isomorphism of D with $\bar{D}$. Now if $\bar{D}_i$ is the image of D_i under this map, then $\bar{D}_i^2 \geq D_i^2$.

We first claim that (Y, D) is isomorphic to $(\bar{Y}, \bar{D})$ if and only if equality holds for every i. Indeed, if equality holds for a given i, then π can't contract any curves which intersect D_i. On the other hand, π can't contract any curves contained in $Y \setminus D$ since then D would not be an anti-canonical cycle.

Now assume that (Y, D) is *not* toric, so that π is not an isomorphism. Now in general, $B \setminus \rho_1$ has a coordinate chart $\psi : B \setminus \rho_1 \to M_{\mathbb{R}}$, constructed by gluing together coordinate charts for $\sigma_{1,2}, \ldots, \sigma_{n-1,n}, \sigma_{n,1}$. Note that $\bar{D}_i^2 > D_i^2$ for at least one i, and by choosing ρ_1 appropriately, we can assume that this is the case for some $i \neq 1$. By comparing this chart ψ with the corresponding chart $\bar{\psi}$ constructed using $(\bar{Y}, \bar{D})$, one sees easily that $\psi(B \setminus \rho_1)$ is properly contained in $\bar{\psi}(B \setminus \rho_1) = M_{\mathbb{R}} \setminus \bar{\rho}_1$. Here $\bar{\rho}_1$ is the one-dimensional cone in the fan for $\bar{Y}$ corresponding to $\bar{D}_1$. In fact, $\psi(B \setminus \rho_1)$ is a wedge (possibly non-convex). If the affine structure extended across the origin, then ψ would extend to an isomorphism $\psi : B \to M_{\mathbb{R}}$, which is impossible. $\square$

Remark 1.4. The fact that $B_{(Y,D)}$ is integral linear, as opposed to just affine, and comes divided into cones has a nice parallel in the symplectic heuristic (observed jointly with D. Auroux). Returning to the notation of §0.6.1, with special Lagrangian fibration $f : U \to B$, we can choose the complex structure on (Y, D) so that $\mathrm{Im}(\Omega)|_U$ is exact. Indeed, one can show that if $p : Y \to \bar{Y}$ is a toric blowup, this exactness holds if and only if the blown up points lie in the unit circle of the structure $\mathbb{G}_m$ of the boundary components (up to the action of the big torus of $\bar{Y}$). Write $\mathrm{Im}(\Omega)|_U = d\gamma$. Then for $L = f^{-1}(b)$, $\gamma|_L$ is closed (by the special Lagrangian condition), and thus determines a class $[\gamma|_L] \in H^1(L, \mathbb{R})$, which is independent of the choice of γ.

There is a canonical identification of the tangent space to B at b with $H^1(L, \mathbb{R})$ using the form $\mathrm{Im}(\Omega)$. Given a tangent vector v at b, lift it to a normal vector field v along L. Then the contraction $(\iota(v)\,\mathrm{Im}(\Omega))|_L$ is a well-defined closed one-form on L, hence defining an element of $H^1(L, \mathbb{R})$. Thus the restrictions $\gamma|_L$ determine a canonically defined vector field v on B_0. One can show using the Arnold-Louiville theorem that locally v is flat with respect to a canonical affine flat connection on the tangent bundle to B_0, with v locally of the form $\sum_i (y_i+c_i)\partial_{y_i}$ for some constants c_i and affine coordinates y_i. From this linearity follows: in a neighbourhood of $b \in B_0$, the vector field gives a canonical linear coordinate chart in the tangent space to B_0 at b.

Now we consider the flow of this vector field. The expectation is that the Lagrangian over a generic point will flow to a node of D — in this way each node $p_{i,i+1} = D_i \cap D_{i+1}$ determines a 2-dimensional cone $\sigma_{i,i+1} \subset B$, while each D_i determines an edge $\rho_i \subset B$, those points whose associated Lagrangian approaches the interior $D_i^o := D_i \setminus \{p_{i-1,i}, p_{i,i+1}\}$ of D_i.

We can consider the universal cover $q : \tilde{B} \to B \setminus \{0\}$. This comes with a decomposition $\tilde{\Sigma}$ into cones minus the origin:

$$\tilde{\Sigma} = \{\sigma_{i,i+1} \setminus \{0\} \,|\, i \in \mathbb{Z}\} \cup \{\rho_i \setminus \{0\} \,|\, i \in \mathbb{Z}\}.$$

In addition $\tilde{B}$ has an integral affine structure pulled back from $B \setminus \{0\}$ and a canonical $\mathbb{Z}$-action sending $\sigma_{i,i+1} \to \sigma_{i+n,i+n+1}$. By patching together integral affine coordinate charts, one obtains a canonical (up to integral linear functions) *developing map* $\delta : \tilde{B} \to \mathbb{R}^2$, an integral affine immersion. Negativity of the boundary $D \subset Y$ is translated into convexity properties of the developing map:

Lemma 1.5. *The developing map δ is one-to-one if and only if D is negative semi-definite. In this case the closure of the image is a convex cone, strictly convex if and only if D is negative definite. If the developing map is not one-to-one, then it surjects onto* $\mathbb{R}^2 \setminus \{0\}$.

Proof. We can assume that D contains no (-1)-curves by contracting such curves. This changes Σ but not B or its affine structure, by Lemma 1.17. Then D is negative semi-definite if $D_i^2 \leq -2$ for all i, and negative definite if in addition $D_i^2 < -2$ for at least one i. If $D_i^2 = -2$ for all i, then by the construction of the affine structure, the developing map, up to an integral linear transformation, identifies $\tilde{B}$ with the upper half-plane, with $\sigma_{i,i+1}$ identified with the cone generated by $(i, 1)$ and $(i + 1, 1)$. If $D_i^2 < -2$ for some i, it is then clear that the developing map identifies $\tilde{B}$ with the interior of a strictly convex cone in $\mathbb{R}^2$.

Conversely, if $D_i^2 \geq 0$ for some i, then again by the construction of the affine structure on B, the image of $\sigma_{i-1,i} \cup \sigma_{i,i+1}$ under the developing map contains a half-plane, from which it follows easily that δ cannot be one-to-one and is surjective. □

The following observation will be needed in a future paper, but for now it illustrates the essential difference between these cases. Suppose we have a point $x \in B_0$, and a tangent vector v to B at x. Then there is a ray emanating from x in the direction

v, which is a straight line. If the tangent vector points towards the origin, then this straight line is just the line segment joining x and the origin. Otherwise, one obtains different possible behaviours. For example, in the negative semi-definite cases, by considering the above description of the developing map, it is possible that such a ray wraps around the origin an infinite number of times. In the best case, this does not happen:

Corollary 1.6. *If D is not negative semi-definite, then any such ray in B which does not pass through the origin goes off to infinity in some cone σ of Σ. That is, writing $\gamma\colon [0,\infty) \to B_0$ for an affine parametrization of the ray, we have $\gamma(t) \in \sigma$ for $t \gg 0$.*

Proof. This follows from the fact that δ is surjective in the non-negative semi-definite case, by the previous lemma. $\square$

Choose a ray ρ_i in Σ. Choosing a ray $\tilde{\rho}_i$ in $\tilde{\Sigma}$ mapping to ρ_i determines a canonical section $s : B \setminus \rho_i \to \tilde{B}$ of q such that the closure of the image is bounded by edges $\tilde{\rho}_i$ and $\tilde{\rho}_{i+n}$. We then obtain an integral affine immersion $\delta \circ s : B \setminus \rho_i \to \mathbb{R}^2$.

Lemma 1.7. *The closure $S := \overline{(\delta \circ s)(B \setminus \rho_i)} \subset \mathbb{R}^2$ is convex (resp. strictly convex) if and only if $(\sum_{j \neq i} a_j D_j)^2 \leq 0$ for any collection of integers a_j (resp. < 0 unless all $a_j = 0$).*

Proof. The set S is the support of a fan, Σ_S, given by the closures of the images under δ of cones $\sigma \in \tilde{\Sigma}$ which intersect $s(B \setminus \{\rho_i\})$, along with the faces of these cones. Note that unless Y was already a toric variety, S has two boundary edges, images of $\tilde{\rho}_i$ and $\tilde{\rho}_{i+n}$. Then Σ_S defines a toric variety whose toric divisors, excluding the ones given by the images of $\tilde{\rho}_i$ and $\tilde{\rho}_{i+n}$, have the same intersection numbers as $D_{i+1}, \ldots, D_{i+n-1}$, by the construction of the affine structure. It is then standard for toric varieties that the convexity of S is equivalent to the negative semi-definiteness of the intersection pairing of $D_{i+1}, \ldots, D_{i+n-1}$ and the strict convexity of S is equivalent to the negative definiteness of $D_{i+1}, \ldots, D_{i+n-1}$. $\square$

Example 1.8. Let Y be a del Pezzo surface of degree 5. Thus Y is isomorphic to the blowup of $\mathbb{P}^2$ in 4 points in general position. The surface Y contains exactly 10 (-1)-curves. It is easy to find an anti-canonical cycle D of length 5 among these 10 curves.

In this case, consider $B \setminus \rho_1$. Each chart $\psi_i : U_i \to M_{\mathbb{R}}$ can be composed with an integral linear function on $M_{\mathbb{R}}$ in such a way that the charts ψ_2, ψ_3, ψ_4 and ψ_5 glue to give a chart $\psi : B \setminus \rho_1 \to M_{\mathbb{R}}$. This can be done, for example, with

$$\psi(v_1) = (1,0), \quad \psi(v_2) = (0,1), \quad \psi(v_3) = (-1,1), \quad \psi(v_4) = (-1,0), \quad \psi(v_5) = (0,-1).$$



FIGURE 1.1.

We can then take a chart $\psi' : U_5 \cup U_1 \to M_{\mathbb{R}}$ which agrees with ψ on $\sigma_{5,1}$, and hence takes the values

$$\psi'(v_5) = (0, -1), \quad \psi'(v_1) = (1, -1), \quad \psi'(v_2) = (1, 0),$$

see Figure 1.1.

Thus B, as an affine manifold, can be constructed by cutting $M_{\mathbb{R}}$ along the positive real axis, and then identifying the two copies of the cone $\sigma_{1,2}$ via an integral linear transformation.

Example 1.9 (Cusps)**.** The next example is basic to our proof of Looijenga's conjecture. We begin with background on cusps: By definition, a cusp is a normal surface singularity for which the exceptional locus of the minimal resolution is a cycle of rational curves. The self-intersections of these exceptional curves determine the analytic type of the singularity, see [L73]. Cusps have a quotient construction due to Hirzebruch [Hi73] which we explain here.

Let $M = \mathbb{Z}^2$. Let $T \in \mathrm{SL}(M)$ be a hyperbolic matrix, i.e., with a real eigenvalue $\lambda > 1$. Then T determines a pair of *dual* cusps as follows: Let $w_1, w_2 \in M_{\mathbb{R}}$ be eigenvectors with eigenvalues $\lambda_1 = 1/\lambda$, $\lambda_2 = \lambda$, chosen so that $w_1 \wedge w_2 > 0$ (in the standard counter-clockwise orientation of $\mathbb{R}^2$). Let $\overline{C}, \overline{C}'$ be the strictly convex cones spanned by w_1, w_2 and $w_2, -w_1$, and let C, C' be their interiors, either of which is preserved by T. Let $U_C, U_{C'}$ be the corresponding tube domains, i.e.,

$$U_C := \{z \in M_{\mathbb{C}} | \operatorname{Im}(z) \in C\}/M \subset M_{\mathbb{C}}/M = M \otimes \mathbb{G}_m.$$

T acts freely and properly discontinuously on $U_C, U_{C'}$. The holomorphic hulls of $U_C/\langle T\rangle, U_{C'}/\langle T\rangle$ are normal surface germs of dual cusps. All cusps (and their duals) arise this way. If M is identified with its dual by choosing an isomorphism $\bigwedge^2 M \cong \mathbb{Z}$, the cone C' is identified with the dual cone of C. In this way $C'/\langle T\rangle$ and $C/\langle T\rangle$ are dual integral affine manifolds, which suggests the duality between the corresponding cusps is a form of mirror symmetry.

Now suppose for a Looijenga pair (Y, D) that $D_i^2 \leq -2$ for all i and D is negative definite (which is equivalent to $D_i^2 \leq -3$ for some i). Then we have a contraction $p : Y \to \bar{Y}$ with $\bar{Y}$ having a single cusp singularity. By Lemma 1.5 the developing map produces an infinite fan $\tilde{\Sigma}$ in a lattice $M \simeq \mathbb{Z}^2$ as follows. We take v_0, v_1 to be a positively oriented basis for $M \simeq \mathbb{Z}^2$, and define v_i for $i \in \mathbb{Z}$ by the relation

$$v_{i-1} + (D^2_{i \bmod n})v_i + v_{i+1} = 0.$$

We take the two-dimensional cones of $\tilde{\Sigma}$ to be the cones generated by v_i and v_{i+1}, $i \in \mathbb{Z}$. If we define $T \in \mathrm{SL}(M)$ by $T(v_0) = v_n$ and $T(v_1) = v_{n+1}$, then $T(v_i) = v_{i+n}$ for each i. Necessarily T is hyperbolic and $|\tilde{\Sigma}|$ is the cone C defined above. Then $U_C/\langle T\rangle$ is a punctured analytic neighborhood of a cusp singularity isomorphic to $z := p(D) \in \bar{Y}$. See [L81], §III.2.

Let $P \in X$ be the dual cusp, thus the holomorphic hull of $U_{C'}/\langle T\rangle$. Here we provide an alternative formulation of Looijenga's condition on smoothability of $P \in X$ (i.e., that the cycle of rational curves for the dual cusp occur as the boundary of a Looijenga pair) which we can phrase just in terms of the hyperbolic matrix T. We claim that Looijenga's condition (that there exists a rational surface Y with anti-canonical boundary divisor D which can be contracted to the dual cusp) is equivalent to the following: There exists distinct primitive vectors $w_1, \ldots, w_r$, spanning rays in counter-clockwise cyclic order in M, and positive integers $n_1, \ldots, n_r$, such that, defining $T_i \in \mathrm{SL}(M)$ by

$$T_i x = x - n_i(w_i \wedge x)w_i,$$

we have

$$T^{-1} = T_r \cdots T_1.$$

That is, there is a factorization of the monodromy matrix T^{-1} of $B_0 := C/\langle T\rangle$ into matrices T_i conjugate to $\begin{pmatrix} 1 & -n_i \\ 0 & 1 \end{pmatrix}$, with the invariant vectors w_i in anticlockwise cyclic order.

To see this, first suppose given such a factorization. Let Σ be a complete fan in $\mathbb{R}^2$ including the rays generated by the w_i together with additional rays so that each cone is convex and generated by a basis of $\mathbb{Z}^2$. Let $\bar{Y}$ be the associated smooth toric surface.

Blowup n_i interior points of the boundary divisor corresponding to w_i for each i. Let (Y', D') be the pair so obtained. By our assumption the integral affine manifold B_0 associated to Y' has monodromy matrix T^{-1}, so is the integral affine manifold $C/\langle T\rangle$ described above. It follows that D' can be contracted to the dual cusp.

Conversely, given (Y, D), after a toric blowup $Y' \to Y$ there exists a toric model $Y' \to \bar{Y}$ by Proposition 1.19. So we obtain a factorization of the monodromy matrix by the reverse of the above process.

The factorization determines a *topological* smoothing of $P \in X$: The projection

$$M_{\mathbb{C}} \to M_{\mathbb{R}}, \quad (z_1, z_2) \mapsto (\operatorname{Im} z_1, \operatorname{Im} z_2)$$

induces an $S^1 \times S^1$-fibration $f^0 \colon X \setminus \{P\} \to C/\langle T\rangle$. We have $B = (C \cup \{0\})/\langle T\rangle = (C/\langle T\rangle) \cup \{0\}$. f^0 extends to a continuous map $f \colon X \to B$, $P \mapsto 0$. Topologically, the existence of the above factorisation allows us to deform the singular fibration f to a $S^1 \times S^1$-fibration with ordinary (Lefschetz) singularities, that is, each singular fibre is a pinched torus and the local monodromy is the Dehn twist in the vanishing cycle δ, acting on $H_1(S^1 \times S^1, \mathbb{Z})$ by $\gamma \mapsto \gamma - (\delta \cap \gamma)\delta$. The monodromy of f at infinity (corresponding to the link L of the singularity $P \in X$) is the matrix T^{-1}, and the above factorization is obtained by expressing this in terms of the local monodromies around the singular fibres of the deformed fibration.

The smoothing we construct for $P \in X$ in our proof of Corollary 0.5 will induce exactly this *topological* smoothing of f. The factoring of the singularity into simple singularities is an instance of the following:

Example 1.10 (Moving worms)**.** Let $p : (Y, D) \to (\bar{Y}, \bar{D})$ be the blowup at $x \in \bar{D}_i \subset \bar{Y}$ which is a smooth point of $\bar{D}$. Let (B, Σ) and $(\bar{B}, \bar{\Sigma})$ be the associated integral affine manifolds with fans. The map

$$p : B \to \bar{B}$$

which identifies each cone in Σ with the corresponding cone in $\bar{\Sigma}$ is a $\mathbb{Z}$PL-isomorphism, and an isomorphism of integral affine manifolds outside of ρ_i, but it is not affine along ρ_i. There is a natural one parameter family of integral affine manifolds interpolating between the two structures by a process Kontsevich and Soibelman [KS06] call *moving worms*. Precisely, choose a point $y \in \rho_i$. Put a new singular affine structure on B by defining a Σ-piecewise linear function to be linear if its restriction to a neighbourhood of $(y, +\infty) \subset \rho_i$ is B-linear, and is $\bar{B}$-linear on $B \setminus [y, +\infty)$. Call the resulting manifold B_y. The map $p : B_y \to \bar{B}$ is a linear isomorphism near 0 (indeed off of $[y, +\infty)$). This new manifold B_y has an I_1, or *focus-focus*, singularity at y, with invariant direction ρ_i, and is non-singular elsewhere. Now under $p : B \to \bar{B}$ we view an I_1 singularity as having moved off of $0 \in B$ and migrated to infinity along the ray ρ_i.

One can iterate this process. If p is now the blowup at m (for simplicity, distinct) points $x_{ij} \in D_i \subset \bar{Y}$, all non-singular points of D, we can similarly define an m-parameter family of integral affine structures, with each p_{ij} determining an I_1 singularity along ρ_i. For example if p is a toric model, i.e., $(\bar{Y}, \bar{D})$ is a smooth toric surface, then we obtain an m-parameter family of structures with m I_1 singularities located along rays of Σ, and non-singular elsewhere. In particular the toric model determines a factorisation of the singularity B into m simple I_1 singularities, and then we *smooth* the manifold by letting these singularities migrate along their invariant direction to infinity.

Moving worms are particularly natural in terms of birational geometry of maximally unipotent $K3$ degenerations. Though it is not needed for the proofs of any of our results, we explain this connection, as the underlying philosophy is relevant to our proof of Theorem 0.1:

Remark 1.11 (Tropical K3 surfaces). Let $p : \mathcal{Y} \to \Delta$ be a type III semi-stable degeneration of $K3$ surfaces. Recall that this means $\mathcal{Y}$ is a smooth 3-fold, Δ is an analytic disk, p is proper, the generic fibre Y^{gen} is a smooth $K3$ surface, and the special fibre Y is a normal crossing divisor with dual complex B_Y a triangulation of S^2. Each vertex v of B_Y corresponds to an irreducible component $Y_v \subset Y$ which is a rational surface with anti-canonical cycle $D_v := \mathrm{Sing}(Y) \cap Y_v$. Each edge of the triangulation of B corresponds to the intersection of two irreducible components, and each 2-simplex corresponds to a triple point of Y.

Let $v_0v_1v_2$, $v_0v_1v_3$ be two triangles of the triangulation of B sharing the edge v_0v_1. We define the affine structure on the union of these two triangles by declaring that a piecewise affine function φ is in fact affine if and only if

$$((\varphi(v_1) - \varphi(v_0))D_{v_1v_0} + (\varphi(v_2) - \varphi(v_0))D_{v_2v_0} + (\varphi(v_3) - \varphi(v_0))D_{v_3v_0}) \cdot D_{v_1v_0} = 0$$

where D_{vw} for adjacent vertices $v, w \in B$ means the corresponding rational curve $Y_v \cap Y_w \subset Y_w$. We use the convention that $D_{vw} \cdot D_{v'w}$ is computed in Y_w. That this is independent of the choice of one of the two vertices on the edge v_0v_1 is equivalent to the condition

$$D^2_{v_0v_1} + D^2_{v_1v_0} = -2, \tag{1.1}$$

which follows from the existence of the semistable smoothing $\mathcal{Y}/\Delta$, see, e.g., [F83a]. The open star of the vertex v in B_Y (the complement to the union of triangles not containing v) can then be identified with an open neighbourhood of the origin in $B_{(Y_v,D_v)}$, the integral affine manifold with singularities associated to the pair (Y_v, D_v).

There can be many semi-stable models $\mathcal{Y} \to \Delta$ with the same generic fibre. Any two are related by simple flops. Suppose for example one component $Y_v \subset Y$ contains a (-1)-curve, $E \subset Y_v$ not contained in $D_v \subset Y_v$. Then $E \cap D_v$ is necessarily a single smooth point. Let $D_\omega \subset D_v$ be the component containing this point; this is a stratum of Y corresponding to an edge ω. Let the second endpoint of ω be w, so that $Y_w \subset Y$ is the other component containing D_ω. Let $p : Y_v \to \overline{Y}_v$ be the blowdown. We can extend this to $p : Y \to \bar{Y}$. Then $\bar{Y}$ is again normal crossing and K-trivial, and its dual complex is the same triangulation of B_Y. We can then define as above an affine structure on the quadrilateral containing ω as a diagonal, using $v_0 = v$ or $v_0 = w$. This gives two affine structures, and now, as (1.1) fails, these affine structures are not compatible. If we choose (arbitrarily) a point $x \in \omega$, there is a canonical structure on this quadrilateral away from x, using the affine structure defined using $v_0 = v$ in a neighbourhood of v and the affine structure defined using $v_0 = w$ in a neighbourhood of w. Since these two affine structures are compatible on the interior of the two-cells of B, this gives an integral affine structure $B^x_{\bar{Y}}$. It is then the case that $Y \to \bar{Y}$ extends to a small contraction $\mathcal{Y} \to \overline{\mathcal{Y}}$. Let $\mathcal{Y}' \to \overline{\mathcal{Y}}$ be the flop, which is again a type III semi-stable degeneration. Note $B_Y, B_{Y'}, B_{\bar{Y}}$ and their triangulations are canonically identified as $\mathbb{Z}$PL-manifolds. The change in integral affine structure is exactly the moving worm procedure: under the flop an I_1 singularity *departs* from $v \in B_Y$, migrates along ω and into $w \in B_{Y'}$. Thus the smooth $K3$ surface Y^{gen} determines an integral affine structure on S^2 canonically up to moving worms, and thus structures independent of moving worms are of particular interest. We will see that our broken lines are one example.

Now suppose $p : \mathcal{Y} \to \overline{\mathcal{Y}}$ is a *toric model*, i.e., the restriction to each irreducible component $p : Y_v \to \bar{Y}_v$ of Y is a toric model. Then $\bar{Y}$ (and a choice of position for each singular point on $B_{\bar{Y}}$) determines an integral affine structure on $B_{\bar{Y}}$ which is smooth at each vertex. The family $\overline{\mathcal{Y}} \to \Delta$ is what Gross and Siebert [GS07] call a *toric degeneration*, and $B_{\bar{Y}}$ is the associated integral affine manifold with singularities of the type that they consider.

1.2. **The Mumford degeneration.**

The toric case of Theorem 0.1 is a special case of a construction due to Mumford which we now recall in a form convenient for our purposes.

Fix a toric monoid P and a lattice $M = \mathbb{Z}^n$. The monoid P is defined by a convex cone $\sigma_P \subset P^{\text{gp}}_{\mathbb{R}}$ with $P = \sigma_P \cap P^{\text{gp}}$. Fix a fan Σ in $M_{\mathbb{R}}$, whose support, $|\Sigma|$, is convex. In what follows, we view $B = |\Sigma|$ as an affine manifold with boundary. We denote by $\Sigma_{\max}$ the set of maximal cones in Σ.

Definition 1.12. A Σ-*piecewise linear function* $\varphi : |\Sigma| \to P^{\mathrm{gp}}_{\mathbb{R}}$ is a continuous function such that for each $\sigma \in \Sigma_{\max}$, $\varphi|_{\sigma}$ is given by an element $\varphi_{\sigma} \in \mathrm{Hom}_{\mathbb{Z}}(M, P^{\mathrm{gp}}) = N \otimes_{\mathbb{Z}} P^{\mathrm{gp}}$.

For each codimension one cone $\rho \in \Sigma$ contained in two maximal cones $\sigma_+, \sigma_- \in \Sigma_{\max}$, we can write

$$\varphi_{\sigma_+} - \varphi_{\sigma_-} = n_{\rho} \otimes p_{\rho,\varphi}$$

where $n_{\rho} \in N$ is the unique primitive element annihilating ρ and positive on σ_+, and $p_{\rho,\varphi} \in P^{\mathrm{gp}}$. We call $p_{\rho,\varphi}$ the *bending parameter*. Note (as the notation suggests) it depends only on the codimension one cone ρ (not on the ordering of σ_+, σ_-).

We then say a Σ-piecewise linear function $\varphi : |\Sigma| \to P^{\mathrm{gp}}$ is P-*convex* if for every codimension one cone $\rho \in \Sigma$, $p_{\rho,\varphi} \in P$. We say φ is *strictly* P-*convex* if for every codimension one cone $\rho \in \Sigma$, $p_{\rho,\varphi} \in P \setminus P^{\times}$, where $P^{\times}$ is the group of invertible elements of P.

Example 1.13. Take a complete fan Σ in $M_{\mathbb{R}}$. This defines a toric variety $Y = Y_{\Sigma}$, which we assume is non-singular. We let $P \subset P^{\mathrm{gp}}$ be given by the cone of effective curves,

$$\mathrm{NE}(Y) \subset A_1(Y) = H_2(Y, \mathbb{Z}).$$

Each codimension one cone $\rho \in \Sigma$ corresponds to a one-dimensional toric stratum $D_{\rho} \subset \partial Y$, hence a class $[D_{\rho}] \in \mathrm{NE}(Y) = P$.

Lemma 1.14. *There is a* Σ*-piecewise linear strictly* P*-convex function* $\varphi : M \to P^{\mathrm{gp}}$ *with*

$$p_{\rho,\varphi} = [D_{\rho}] \tag{1.2}$$

for each codimension one cone $\rho \in \Sigma$*. Up to a linear function,* φ *is unique.*

Proof. We will only need this result in two dimensions, so we only prove it in this case. This is trivial unless Σ is complete, so let's assume completeness. Let $\rho_1, \ldots, \rho_s$ be the rays in the fan Σ in cyclic order. It is sufficient to show that

$$\sum_{i=1}^{s} n_{\rho_i} \otimes [D_{\rho_i}] = 0$$

in $N \otimes_{\mathbb{Z}} P^{\mathrm{gp}}$, where n_{ρ_i} is chosen, say, to be positive on ρ_{i+1}. However, using a suitable choice of identification of $\bigwedge^2 M \cong \mathbb{Z}$, we note that n_{ρ_i} can be identified with the functional on M given by $m \mapsto m_i \wedge m$, where m_i is a primitive generator of ρ_i. Thus we need to show that

$$\sum_{i=1}^{s} m_i \otimes [D_{\rho_i}] = 0.$$

This we can show by evaluating the left-hand side on an element $n \in N$, getting

$$\sum_{i=1}^{s} \langle n, m_i \rangle [D_{\rho_i}].$$

However, this is precisely the divisor of zeroes and poles of z^n, and hence is zero in $P^{\mathrm{gp}} = H_2(Y, \mathbb{Z})$. $\square$

Given a Σ-piecewise linear and P-convex function $\varphi : |\Sigma| \to P^{\mathrm{gp}}$ we can define a monoid $P_\varphi \subset M \times P^{\mathrm{gp}}$ by

$$P_\varphi := \{(m, \varphi(m) + p) \,|\, m \in |\Sigma|, p \in P\}. \tag{1.3}$$

The convexity condition implies that P_φ is closed under addition. Furthermore, we have a natural inclusion $P \hookrightarrow P_\varphi$ given by $p \mapsto (0, p)$, which gives us a flat morphism

$$f : \operatorname{Spec} \Bbbk[P_\varphi] \to \operatorname{Spec} \Bbbk[P].$$

It is easy to see that a general fibre of f is isomorphic to the algebraic torus $\Bbbk[M]$: in fact, if we consider the big torus orbit $U = \operatorname{Spec} \Bbbk[P^{\mathrm{gp}}] \subset \operatorname{Spec} \Bbbk[P]$, $f^{-1}(U) = U \times \operatorname{Spec} \Bbbk[M]$.

We now describe the fibres over other toric strata of $\operatorname{Spec} \Bbbk[P]$. Let $x \in \operatorname{Spec} \Bbbk[P]$ be a point in the torus orbit corresponding to a face $Q \subset P$. Then by replacing P with the localized monoid $P - Q$ obtained by inverting all elements of Q, we may assume that x is contained in the smallest toric stratum of $\operatorname{Spec} \Bbbk[P]$. Consider the composed map

$$\bar{\varphi} : |\Sigma| \xrightarrow{\varphi} P^{\mathrm{gp}} \to P^{\mathrm{gp}}/P^{\times}.$$

Note $\bar{\varphi}$ is also piecewise linear. Let $\bar{\Sigma}$ be the fan (of not necessarily convex cones) whose maximal cones are the maximal domains of linearity of $\bar{\varphi}$. Then $f^{-1}(x)$ can be written as

$$f^{-1}(x) = \operatorname{Spec} \Bbbk[\bar{\Sigma}].$$

Here,

$$\Bbbk[\bar{\Sigma}] = \bigoplus_{m \in M \cap |\Sigma|} \Bbbk z^m$$

with multiplication given by

$$z^m \cdot z^{m'} = \begin{cases} z^{m+m'} & \text{if } m, m' \text{ lie in a common cone of } \bar{\Sigma}, \\ 0 & \text{otherwise.} \end{cases} \tag{1.4}$$

In particular, the irreducible components of $f^{-1}(x)$ are the toric varieties $\operatorname{Spec} \Bbbk[\sigma \cap M]$ for $\sigma \in \bar{\Sigma}_{\max}$.

In the particular case that $\operatorname{rank} M = 2$ and Σ defines a non-singular complete surface with n toric divisors, suppose φ is strictly convex. Then if x is a point of the smallest toric stratum of $\operatorname{Spec}\Bbbk[P]$, then $f^{-1}(x)$ is just $\mathbb{V}_n \subset \mathbb{A}^n$, the reduced cyclic union of coordinate $\mathbb{A}^2$'s:

$$\mathbb{V}_n = \mathbb{A}^2_{x_1,x_2} \cup \mathbb{A}^2_{x_2,x_3} \cup \cdots \cup \mathbb{A}^2_{x_n,x_1} \subset \mathbb{A}^n_{x_1,\ldots,x_n}.$$

We call $\mathbb{V}_n$ the *vertex*, or more specifically, the *n-vertex.*

We will need in the sequel the degenerate case of the n-vertex for $n = 2$. This is a union of two affine planes and can be described as the double cover

$$\mathbb{V}_2 = \operatorname{Spec}\Bbbk[x_1, x_2, y]/(y^2 - x_1^2x_2^2) = \mathbb{A}^2_{x_1,x_2} \cup \mathbb{A}^2_{x_2,x_1}. \tag{1.5}$$

Of course, this does not appear as a central fibre of a Mumford degeneration. Analogously, one can define

$$\mathbb{V}_1 = \operatorname{Spec}\Bbbk[x, y, z]/(xyz - x^2 - z^3), \tag{1.6}$$

the affine cone over a nodal cubic.

Example 1.15. In Example 1.13, with the choice of φ given by Lemma 1.14, the family

$$\operatorname{Spec}\Bbbk[P_\varphi] \to \operatorname{Spec}\Bbbk[\mathrm{NE}(Y)]$$

in fact gives the family of mirror manifolds to the toric variety Y, as constructed by Givental [Giv].

In fact, the mirror of a toric variety also includes the data of a *Landau-Ginzburg potential*, which is a regular function. In this case the potential is

$$W = \sum_\rho z^{(m_\rho,\varphi(m_\rho))}$$

where we sum over all rays $\rho \in \Sigma$, and $m_\rho \in M$ denotes the primitive generator of ρ. If Y is not Fano, then this is actually not the correct Landau-Ginzburg potential; rather, the potential receives corrections which can be viewed as coming from degenerate holomorphic disks on Y with irreducible components mapping into D. See Remark 0.18.

1.3. **Some toric constructions.**

We collect here a few simple observations which we use repeatedly throughout the paper.

Definition 1.16 (Toric Blowup)**.** Let Y be a rational surface and D an anti-canonical cycle of rational curves on Y. A *toric blow-up* of the pair (Y, D) is a birational morphism $\pi : \tilde{Y} \to Y$ such that if $\tilde{D}$ is the reduced scheme structure on $\pi^{-1}(D)$, then $\tilde{D}$ is an anti-canonical cycle of rational curves on $\tilde{Y}$.

Given (B, Σ), a *refinement* is a pair $(B, \tilde{\Sigma})$, where $\tilde{\Sigma}$ is a decomposition of B into rational polyhedral cones refining Σ.

Lemma 1.17. *There is a one-to-one correspondence between toric blow-ups of* (Y, D) *and refinements of* (B, Σ). *Furthermore, if* $(\tilde{Y}, \tilde{D})$ *is a non-singular toric blow-up of* (Y, D), *and* $(\tilde{B}, \tilde{\Sigma})$ *is the affine manifold with singularities constructed from* $(\tilde{Y}, \tilde{D})$, *then* $\tilde{B}$ *and* B *are isomorphic as integral affine manifolds with singularities in such a way that* $\tilde{\Sigma}$ *is the corresponding refinement of* Σ.

Proof. Let $\pi : \tilde{Y} \to Y$ be a toric blow-up. It follows easily from the condition that $\pi^{-1}(D)_{\mathrm{red}}$ is an anti-canonical divisor that $\pi : \tilde{Y} \setminus \pi^{-1}(\mathrm{Sing}(D)) \to Y \setminus \mathrm{Sing}(D)$ is an isomorphism. Thus π is a blow-up along a subscheme supported on $\mathrm{Sing}(D)$. Let $x \in \mathrm{Sing}(D)$ be a double point of D, corresponding to a cone $\sigma \in \Sigma$. Note σ can be viewed as a rational polyhedral cone defining a non-singular toric variety $X_\sigma \cong \mathbb{A}^2$. Then étale locally near x, the pair (Y, D) is isomorphic to the pair $(X_\sigma, \partial X_\sigma)$. One can then check that in this local model, the only possible blow-ups satisfying the definition of toric blow-ups come from subdivisions of the cone σ, i.e., toric blow-ups of X_σ. Indeed, the exceptional divisors of toric blowups are the only divisors with log discrepancy -1. This gives the desired correspondence. The second statement is then easily checked. □

Definition 1.18. A *toric model* of (Y, D) is a birational morphism $(Y, D) \to (\bar{Y}, \bar{D})$ to a smooth toric surface with its toric boundary such that $D \to \bar{D}$ is an isomorphism.

Proposition 1.19. *Given* (Y, D) *there exists a toric blowup* $(\tilde{Y}, \tilde{D})$ *which has a toric model* $(\tilde{Y}, \tilde{D}) \to (\bar{Y}, \bar{D})$.

Proof. First observe:

1. Let $p : Y \to Y'$ be the blowdown of a (-1)-curve not contained in D, and $D' := p_*(D) \subset Y'$. If the proposition holds for (Y', D') then it holds for (Y, D).
2. Let $Y'' \to Y$ be the blowup at a node of D, and $D'' \subset Y''$ the reduced inverse image of D. The proposition holds for (Y'', D'') if and only if it holds for (Y, D).

By using (1) and (2) repeatedly we may assume Y is minimal, and thus is either a ruled surface or is $\mathbb{P}^2$. In the latter case, by blowing up a node of D we reduce to the ruled case. So we have $q : Y \to \mathbb{P}^1$ a ruling. We next consider the number of components of D contained in fibres of q. There cannot be more than two such components, for otherwise D cannot be a cycle. If there are precisely two such components, then D necessarily has precisely four components, and it is then easy to check that D is the toric boundary of Y, for a suitable choice of toric structure on Y. In this case the proposition obviously holds. Otherwise let $D' \subset D$ be the union of components not contained in fibres. If D' has a node, then we can blowup the node, blowdown the strict transform of the fibre (through the node), increasing the number of components of D contained in fibres.

After carrying out this procedure for each node of D', we are then in one of two cases.

Case I. D has two components contained in fibres, and then we are done.

Case II. D consists of a fibre f and a non-singular irreducible two-section D' of q. Note that since $D' + f \sim -K_Y$ and Y is isomorphic to the Hirzebruch surface $\mathbb{F}_e$ for some e, we can write $\operatorname{Pic} Y = \mathbb{Z}C_0 \oplus \mathbb{Z}f$, with $C_0^2 = -e$ and $-K_Y = 2C_0 + (e+2)f$. Thus $D' \sim 2C_0 + (e+1)f$ and $C_0 \cdot D' = -e+1$. Since C_0 is not contained in D', $e = 0$ or 1.

If $e = 0$, then there is a second ruling $q' : Y \to \mathbb{P}^1$, with D' and f sections of this ruling. In this case, we follow the same procedure as above of blowing up nodes for this new ruling, arriving in Case I.

If $e = 1$, then C_0 is disjoint from D'. Blowing down C_0, we obtain $\mathbb{P}^2$, and can then blowup one of the nodes of the image of $D' \cup f$. Using this new ruled surface, we can again blowup a node and find ourselves back in Case I. $\square$

2. Modified Mumford deformations

As throughout the paper, (Y, D) is a Looijenga pair, where D is a cycle of length n, and $(B = B_{(Y,D)}, \Sigma)$ is the tropicalisation defined in §1.1. Next we explain how to generalize Mumford's family, to give a canonical formal deformation of $\mathbb{V}_n^o$ associated to (Y, D). In local coordinates this is described in §0.6.2; here we give an abstract coordinate free description. Locally on B_0 the picture is toric and we have Mumford's deformations described in §1.2. As Mumford's construction is functorial, the deformations built locally patch canonically together. Here are the details.

2.1. Generalities on integral linear manifolds and multi-valued functions.

Let us consider a somewhat more general situation, in which B_0 is an integral linear surface. For any locally constant sheaf $\mathcal{F}$ on B_0, and any simply connected subset $\tau \subset B_0$ we write $\mathcal{F}_\tau$ for the stalk of this local system at any point of τ (as any two such stalks are canonically identified by parallel transport). In particular, we apply this for the sheaf Λ of integral constant vector fields, and $\Lambda_\mathbb{R} = \Lambda \otimes_\mathbb{Z} \mathbb{R}$.

One feature of linear manifolds is that for any simply connected open set $U \subset B_0$, there is a canonical linear immersion $U \to \Lambda_{\mathbb{R},\tau}$, compatible with parallel transport inside U. The construction is functorial in the obvious way for linear maps between linear manifolds, and this allows us to generalize many convex geometric notions from vector spaces to linear manifolds.

A rational polyhedral fan Σ on B_0 is a polyhedral decomposition of B_0 whose restriction to each linear chart $B_0 \supset U \to \mathbb{R}^2$ is the restriction of a fan in the usual sense; the reader should think of the example of §1.1.

For each cone $\tau \in \Sigma$, and each cone $\sigma \in \Sigma$ containing τ, there is a canonical inclusion $\sigma \subset \Lambda_{\tau,\mathbb{R}}$. We define the *localized fan* $\tau^{-1}\Sigma$ in $\Lambda_{\tau,\mathbb{R}}$ as the collection of cones

$$\tau^{-1}\Sigma = \{\sigma + \mathbb{R}\tau \subset \Lambda_{\tau,\mathbb{R}} \,|\, \sigma \in \Sigma, \tau \subset \sigma\}.$$

When τ is positive dimensional, this is a fan of not strictly convex cones: if τ is two-dimensional, then $\tau^{-1}\Sigma$ consists just of the entire space $\Lambda_{\tau,\mathbb{R}}$, while if τ is one-dimensional, it consists of the tangent line to τ in $\Lambda_{\tau,\mathbb{R}}$ along with the two closed half-planes with boundary this tangent line.

Remark 2.1. Note that if q is a point of B_0 with $q \in \sigma \in \Sigma$, and $\tau \subset \sigma$, then the canonical embedding of a neighbourhood of τ in $\Lambda_{\tau,\mathbb{R}}$ identifies q with a point of $\Lambda_{\tau,\mathbb{R}}$. We shall use this identification freely in what follows.

Let $P \subseteq P^{\mathrm{gp}}$ be a toric monoid as in §1.2. Let $\pi : \mathbb{P}_0 \to B_0$ be a linear locally trivial principal $P^{\mathrm{gp}}_{\mathbb{R}} := P^{\mathrm{gp}} \otimes_{\mathbb{Z}} \mathbb{R}$ bundle. In other words, $\mathbb{P}_0$ is an integral linear manifold, π is linear, we have local trivializations $\pi^{-1}(U) = U \times P^{\mathrm{gp}}_{\mathbb{R}}$, and the transition functions respect the translation action on the second factor. We write $\mathcal{P} := \pi_*\Lambda_{\mathbb{P}_0}$ (the local system whose sections over a simply connected open set $U \subset B_0$ are the constant integral tangent vectors on $\pi^{-1}(U)$). Thus we can write $\mathcal{P}_\tau$ for $\tau \in \Sigma$, and this coincides with the stalk of $\Lambda_{\mathbb{P}_0}$ at any point $x \in \pi^{-1}(\tau)$.

We use this setup to generalize the kind of piecewise linear function which appeared in §1.2. In particular, one can consider Σ-piecewise linear sections $\varphi : B_0 \to \mathbb{P}$. These are continuous sections of π which are linear on cones of Σ. Furthermore, one can define convexity in the same sense as Definition 1.12. Indeed, using a trivialization $\pi^{-1}(U) = U \times P^{\mathrm{gp}}_{\mathbb{R}}$ in a neighbourhood of a point of τ, we can describe φ by giving a piecewise linear function $\varphi_U : U \to P^{\mathrm{gp}}_{\mathbb{R}}$, and so, as in Definition 1.12, there is a bending parameter p_{ρ,φ_U}. We say φ is convex (strictly convex) if $p_{\rho,\varphi_U} \in P$ ($P \setminus P^\times$) for all ρ, U. Note p_{ρ,φ_U} is defined independently of the trivialization, since changing the choice of trivialization only changes φ_U by a linear function.

To see how such bundles and sections arise, we now specialize to the case $B = B_{(Y,D)}$, and $B_0 \subset B$ the smooth locus (which is $B \setminus \{0\}$ unless (Y,D) is toric, in which case $B_0 = B = \mathbb{R}^2$). Recall from §1.1 that B_0 is covered by open sets $U_1, \ldots, U_n$ on which charts are defined, with $U_i = \mathrm{Int}(\sigma_{i-1,i} \cup \sigma_{i,i+1})$ and $\rho_i = \sigma_{i-1,i} \cap \sigma_{i,i+1}$.

Definition 2.2. A *($P^{\mathrm{gp}}_{\mathbb{R}}$-valued) Σ-piecewise linear multivalued function* on B is a collection $\varphi = \{\varphi_i\}$ with φ_i a Σ-piecewise linear function on U_i with values in $P^{\mathrm{gp}}_{\mathbb{R}}$.

Note this is equivalent to giving a $\rho_i^{-1}\Sigma$-piecewise linear function $\varphi_i : \Lambda_{\mathbb{R},\rho_i} \to P^{\mathrm{gp}}_{\mathbb{R}}$ for each ray $\rho_i \in \Sigma$. Two such functions φ, φ' are said to be *equivalent* if $\varphi_i - \varphi'_i$ is linear for each i. Note the equivalence class of φ is determined by the collection of bending parameters $p_{\rho,\varphi} \in P$.

We drop the modifiers Σ and P when they are clear from context.

The collection $\{\varphi_i\}$ determines a $P^{\mathrm{gp}}_{\mathbb{R}}$-principal bundle $\pi : \mathbb{P}_0 \to B_0$ with a piecewise linear convex section $\varphi : B_0 \to \mathbb{P}_0$ in an obvious way: we glue $U_i \times P^{\mathrm{gp}}_{\mathbb{R}}$ to $U_{i+1} \times P^{\mathrm{gp}}_{\mathbb{R}}$ along $(U_i \cap U_{i+1}) \times P^{\mathrm{gp}}_{\mathbb{R}}$ by

$$(x, p) \to (x, p + \varphi_{i+1}(x) - \varphi_i(x)).$$

By construction we have local sections $x \mapsto (x, \varphi_i(x))$ which patch to give the piecewise linear section φ. One checks immediately the isomorphism class (of the $P^{\mathrm{gp}}_{\mathbb{R}}$-principal bundles together with the section) depends only on the equivalence class of $\{\varphi_i\}$.

This construction gives rise to a sheaf $\mathcal{P} := \pi_* \Lambda_{\mathbb{P}_0}$ on B_0, and taking the differential of the projection π gives a canonical exact sequence

$$0 \to \underline{P}^{\mathrm{gp}} \to \mathcal{P} \xrightarrow{r} \Lambda_{B_0} \to 0 \tag{2.1}$$

of local systems on B_0, where $\underline{P}^{\mathrm{gp}}$ denotes the constant local system. We shall write Λ for Λ_{B_0}, as we shall only consider sheaves on B_0.

Example 2.3. Our standard example, fundamental to this paper, will be as follows. Suppose P is a monoid which comes with a homomorphism $\eta : \mathrm{NE}(Y) \to P$ of monoids.

Choose φ by specifying φ_i on U_i by the formula

$$p_{\rho_i, \varphi_i} = \eta([D_i]).$$

Such a φ is well-defined up to linear functions, and always exists. This is always convex, and is strictly convex provided $\eta([D_i])$ is not invertible for any i.

In terms of the symplectic heuristic, in the case that η induces an isomorphism between $H_2(Y, \mathbb{Z})$ and P^{gp}, we view the sequence (2.1) as analogous to the relative homology sequence

$$0 \to H_2(Y, \mathbb{Z}) \to H_2(Y, L, \mathbb{Z}) \to H_1(L, \mathbb{Z}) \to 0 \tag{2.2}$$

where L is a Lagrangian fibre of the SYZ fibration. Of course varying L this becomes a sequence of local systems.

We continue to assume we have a Σ-piecewise linear P-convex section $\varphi : B_0 \to \mathbb{P}_0$ as above, with $B = B_{(Y,D)}$. We explain how Mumford's construction determines a canonical formal deformation of $\mathbb{V}^o_n$.

For each τ, φ determines a $\tau^{-1}\Sigma$-piecewise linear P-convex function $\varphi_\tau : \Lambda_{\mathbb{R},\tau} \to \mathcal{P}_{\mathbb{R},\tau}$ (note the target is a $P^{\mathrm{gp}}_{\mathbb{R}}$ torsor, so P-convex makes sense).

Now define the toric monoid $P_{\varphi_\tau} \subset \mathcal{P}_\tau$ by

$$P_{\varphi_\tau} := \{q \in \mathcal{P}_\tau | q = p + \varphi_\tau(m) \text{ for some } p \in P,\ m \in \Lambda_\tau\}. \tag{2.3}$$

By the definition of convex, we have canonical inclusions

$$P_{\varphi_\rho} \subset P_{\varphi_\sigma} \subset \mathcal{P}_\rho \tag{2.4}$$

whenever $\rho \subset \sigma \in \Sigma$. If $\rho \in \Sigma$ is a ray with $\rho \subset \sigma_\pm \in \Sigma_{\max}$ we have the equality

$$P_{\varphi_{\sigma_+}} \cap P_{\varphi_{\sigma_-}} = P_{\varphi_\rho}. \tag{2.5}$$

Definition 2.4 (Monomial ideals)**.** A *(monoid) ideal* of a monoid P is a subset $I \subset P$ such that $p \in I, q \in P$ implies $p + q \in I$. An ideal determines a *monomial ideal* in the monoid ring $\Bbbk[P]$, generated by monomials z^p for $p \in I$. We also denote this ideal by I, hopefully with no confusion. As a consequence, we shall sometimes write certain ideal operations either additively or multiplicatively, i.e., for $J \subset P$,

$$kJ = \{p_1 + \cdots + p_k \,|\, p_i \in J, 1 \leq i \leq k\},$$

and the corresponding monomial ideal is J^k.

Let $\mathfrak{m} = P \setminus P^\times$. This is the unique maximal ideal of P, defining a monomial ideal $\mathfrak{m} \subset \Bbbk[P]$. Note $\Bbbk[P]/\mathfrak{m} \cong \Bbbk[P^\times]$.

We say an ideal $I \subset P$ is $\mathfrak{m}$-*primary* if $\sqrt{I} = \mathfrak{m}$, in which case the same holds for the associated monomial ideal $I \subset \Bbbk[P]$.

In this paper we only consider *toric monoids* P, that is, $P = \sigma \cap L$ where L is a free abelian group of finite rank and $\sigma \subset L_\mathbb{R}$ is a rational polyhedral cone (thus $\Bbbk[P]$ is the coordinate ring of an affine toric variety over $\Bbbk$). In particular, $\Bbbk[P]$ is Noetherian. If σ is strictly convex then $\mathfrak{m}$ is the maximal ideal corresponding to the unique torus fixed point of $\operatorname{Spec} \Bbbk[P]$.

Fix an ideal $I \subset P$. Define $I_{\tau,\sigma} \subset P_{\varphi_\tau}$ for $\tau, \sigma \in \Sigma$ with $\tau \subset \sigma \in \Sigma_{\max}$, $\tau \neq \{0\}$, by

$$I_{\tau,\sigma} := \{q \in P_{\varphi_\tau} | q - \varphi_\sigma(r(q)) \in I\}. \tag{2.6}$$

Finally, for $\tau_1 \subset \tau_2$,

$$I_{\tau_1,\tau_2} := \bigcup_{\sigma \supseteq \tau_2} I_{\tau_1,\sigma}, \tag{2.7}$$

Note the union is just the single ideal $I_{\tau_1,\sigma=\tau_2}$ defined in (2.6) except in the case $\tau_1 = \tau_2 = \rho$ is a ray of the fan.

Now we define

$$R_{\tau_1,\tau_2,I} := \Bbbk[P_{\varphi_{\tau_1}}]/I_{\tau_1,\tau_2}.$$

We note that in the special case $\tau_1 = \tau_2 = \sigma \in \Sigma_{\max}$, then $I_{\sigma,\sigma} = I \cdot \Bbbk[P_{\varphi_\sigma}]$ and

$$\begin{aligned} R_{\sigma,\sigma,I} &= \Bbbk[P_{\varphi_\sigma}]/I\Bbbk[P_{\varphi_\sigma}] \\ &\cong \Bbbk[\Lambda_\sigma] \otimes_\Bbbk \Bbbk[P]/I. \end{aligned}$$

For ρ a ray and σ a two-dimensional cell with $\rho \subset \sigma$, there is a natural surjection

$$R_{\rho,\sigma,I} \twoheadrightarrow R_{\rho,\rho,I}.$$

By (2.4), there is also a natural injective map

$$R_{\rho,\sigma,I} \to R_{\sigma,\sigma,I}$$

which is the localization at $z^{\varphi_\rho(m)}$ for any $m \in \rho^{-1}\sigma$ that generates $\Lambda_\rho/(\mathbb{R}\rho \cap \Lambda_\rho)$.

From this system of rings we can build a scheme X_I^o over $\operatorname{Spec} \Bbbk[P]/I$ as follows. First, if $\rho \subset \sigma_\pm$, let

$$R_{\rho,I} := R_{\rho,\sigma_+,I} \times_{R_{\rho,\rho,I}} R_{\rho,\sigma_-,I}.$$

Note $R_{\rho,I}$ is a $\Bbbk[P]/I$-algebra.

Lemma 2.5. *Let $I \subset P$ be a monomial ideal. Then $R_{\rho,I} = \Bbbk[P_{\varphi_\rho}] \otimes_{\Bbbk[P]} \Bbbk[P]/I$. So if $\rho = \rho_i$, then $\operatorname{Spec} R_{\rho,I} \to \operatorname{Spec} \Bbbk[P]/I$ is the base-change of the Mumford degeneration induced by the PL function φ_i on the localized fan $\rho^{-1}\Sigma$.*

Proof. It is easy to check that the map

$$\Bbbk[P_{\varphi_\rho}]/I\Bbbk[P_{\varphi_\rho}] \to R_{\rho,I}$$

given by $f \mapsto (f \bmod I_{\rho,\sigma_+}, f \bmod I_{\rho,\sigma_-})$ is an isomorphism, using the fact that

$$I\Bbbk[P_{\varphi_\rho}] = I_{\rho,\sigma_+} \cap I_{\rho,\sigma_-}.$$

□

Remark 2.6. Since $\operatorname{Spec} R_{\rho,I} \to \operatorname{Spec} \Bbbk[P]/I$ is a base-change of the Mumford degeneration, we can in fact say what a fibre of this morphism is over a point x in the smallest toric stratum of $\operatorname{Spec} \Bbbk[P]$, i.e., a point in $\operatorname{Spec} \Bbbk[P]/\mathfrak{m}$. This depends on whether $p_{\rho,\varphi} \in P$ is invertible or not. If it is not invertible, then the fibre is $\operatorname{Spec} \Bbbk[\rho^{-1}\Sigma] \cong \operatorname{Spec} \Bbbk[x,y,z^{\pm 1}]/(xy)$. If $p_{\rho,\varphi}$ is invertible, then the fibre is $\operatorname{Spec} \Bbbk[\mathbb{Z}^2]$. In this latter case, in fact the surjective maps $R_{\rho,\sigma_\pm,I} \to R_{\rho,\rho,I}$ are isomorphisms, and the fibred product $R_{\rho,I}$ is isomorphic to any of these three rings.

We now have natural maps

$$\psi_{\rho,\pm} : R_{\rho,I} \to R_{\sigma_\pm,\sigma_\pm,I}$$

given by $\psi_{\rho,\pm}(z^p) = z^p$, using the inclusions $P_{\varphi_\rho} \subset P_{\varphi_{\sigma_\pm}}$ of (2.4). One sees easily that the $\psi_{\rho,\pm}$ are in fact localization maps, given by localization at $z^{\varphi_\rho(m)}$ for any $m \in \rho^{-1}\sigma_\pm$ that generates $\Lambda_\rho/(\mathbb{R}\rho \cap \Lambda_\rho)$. Thus if $\rho \subset \sigma$, we obtain open subschemes

$$U_{\rho,\sigma,I} := \operatorname{Spec} R_{\sigma,\sigma,I} \subset U_{\rho,I} := \operatorname{Spec} R_{\rho,I}.$$

Of course, if $\rho_\pm$ are the two edges of σ, $U_{\rho_-,\sigma,I}$ and $U_{\rho_+,\sigma,I}$ are canonically isomorphic. Note also that

$$U_{\rho,\sigma_+,I} \cap U_{\rho,\sigma_-,I} \cong \operatorname{Spec} \Bbbk[\Lambda_\rho] \times \operatorname{Spec}(\Bbbk[P]/I)_{z^{p_{\rho,\varphi}}} \cong (\mathbb{G}_m)^2 \times \operatorname{Spec}(\Bbbk[P]/I)_{z^{p_{\rho,\varphi}}} \tag{2.8}$$

Note that if $p_{\rho,\varphi} \in \sqrt{I}$ then the localization $(\Bbbk[P]/I)_{z^{p_{\rho,\varphi}}}$ is zero, and the intersection is empty. This motivates the following definition.

Definition 2.7. Given (B, Σ), φ a PL multivalued convex function on B with values in $P^{\mathrm{gp}}_{\mathbb{R}}$, and $I \subseteq P$ an ideal, we define Σ_I to be the decomposition of B into cones obtained by removing any ray $\rho \in \Sigma$ such that $p_{\rho,\varphi} \notin \sqrt{I}$, so that Σ is a refinement of Σ_I. Then Σ_I is a fan of not necessarily convex cones.

We can now define our analogue of the Mumford degeneration.

Definition-Lemma 2.8. Suppose that Σ_I contains at least two rays. Then the canonical identifications of the open sets $U_{\rho_i,\sigma_{i,i+1},I}$ with $U_{\rho_{i+1},\sigma_{i,i+1},I}$ generate an equivalence relation on $\coprod_i U_{\rho_i,I}$, and the quotient by this equivalence relation defines a scheme X_I^o over $\operatorname{Spec} \Bbbk[P]/I$.

Proof. Let ρ_i, ρ_j be two rays of Σ contained in the same maximal cone $\sigma \in \Sigma_I$. Assume without loss of generality that $i < j$ and $\rho_{i+1}, \ldots, \rho_{j-1}$ also lie in σ. Let $U_{ij} \subseteq U_{\rho_i,\sigma_{i,i+1}} \subseteq U_{\rho_i,I}$ be the open set defined by localizing $R_{\sigma_{i,i+1},\sigma_{i,i+1},I}$ at $\prod_{k=i+1}^{j-1} z^{p_{\rho_k,\varphi}}$, and define $U_{ji} \subseteq U_{\rho_j,\sigma_{j-1,j}} \subseteq U_{\rho_j,I}$ by localizing $R_{\sigma_{j-1,j},\sigma_{j-1,j},I}$ at this same element. One sees from (2.8) that the gluings of $U_{\rho_k,I}$ with $U_{\rho_{k+1},I}$, $i \leq k < j$, induce an isomorphism $\psi_{ij} : U_{ij} \to U_{ji}$. One checks easily that these isomorphisms satisfy the requirements for gluing data for schemes along open subsets, see e.g., [H77], Ex. II 2.12. $\square$

Remarks 2.9. (1) X_I^o only depends on the equivalence class of φ, since the monoids P_{φ_τ}, P_{φ_σ} are canonically defined, independently of the choice of representative for φ.

(2) One could work in the case that only one ray ρ satisfies $p_{\rho,\varphi} \in \sqrt{I}$ (recall in Definition 2.8 we require at least two such rays). In this case we are identifying two disjoint open subsets of $U_{\rho,I}$ via an étale equivalence relation, so a priori this only gives an algebraic space. To avoid such technicalities we do not consider this case, though after straightforward adjustments our method does apply. For $\sqrt{I} = I_q$ the ideal associated to a contraction (a morphism with connected fibres) $q : Y \to Z$, this means we require $q|_{D_i}$ is finite for at least two components D_i (though again, this is purely to avoid the need for algebraic spaces).

(3) If $p_{\rho,\varphi} \in P^\times$ for all rays ρ in Σ one obtains a single open set, isomorphic to an algebraic torus over $\operatorname{Spec} \Bbbk[P]/I$, and dividing out by the identifications would divide this torus out by the action of $T \in \mathrm{SL}_2(\mathbb{Z})$ representing the monodromy of the

singularity of B. This quotient is badly behaved in general — it does not have the structure of a scheme or complex analytic space (it is not Hausdorff). However, when D is negative definite and $\Bbbk = \mathbb{C}$, let $f : Y \to Y'$ be the contraction of D. Suppose P is a toric monoid containing $\mathrm{NE}(Y)$ such that the classes of the D_i's generate a face of P. If we take $J := I_f$ to be the complement of this face, and I to be an ideal with $\sqrt{I} = J$, then after restricting to an appropriate analytic open set, the action is properly discontinuous and so the quotient has the structure of a complex analytic space, see §4.1. For example, if we take $I = J$ the prime ideal, the resulting family is a degeneration of the dual cusp to $\mathbb{V}_n$; the generic fibre is exactly the quotient construction of the dual cusp given in Example 1.9. This is why Looijenga's conjecture follows naturally from Theorem 0.1. We give the details in §4.

We first analyze this construction in the purely toric case:

Lemma 2.10. *For (Y, D) toric and $\mathbb{P} = B \times P^{\mathrm{gp}}_{\mathbb{R}}$, X^o_I is an open subscheme of the Mumford degeneration* $\operatorname{Spec} \Bbbk[P_\varphi]/I\Bbbk[P_\varphi]$, *and*

$$H^0(X^o_I, \mathcal{O}_{X^o_I}) = \Bbbk[P_\varphi]/I\Bbbk[P_\varphi].$$

Proof. Note that for $\tau \in \Sigma$, the monoid P_{φ_τ} is isomorphic to the localization of P_φ along the face $\{(m, \varphi(m)) \,|\, m \in \tau \cap M\}$. Thus $\operatorname{Spec} \Bbbk[P_{\varphi_\tau}]$ is an open subset of $\operatorname{Spec} \Bbbk[P_\varphi]$ and $\operatorname{Spec} \Bbbk[P_{\varphi_\tau}] \otimes_{\Bbbk[P]} \Bbbk[P]/I$ is an open subset of $\operatorname{Spec} \Bbbk[P_\varphi]/I\Bbbk[P_\varphi]$. Furthermore, the gluing procedure constructing X^o_I is clearly compatible with these inclusions, so X^o_I is an open subscheme of $\operatorname{Spec} \Bbbk[P_\varphi]/I\Bbbk[P_\varphi]$. Next, looking at the fibre over a closed point, one sees easily that the underlying topological space of these fibres is obtained just by removing the zero-dimensional torus orbit from the corresponding fibre of the Mumford degeneration. The closed fibres of the Mumford degeneration are S_2 by [A02], 2.3.19. Thus by Lemma 2.11, the result follows. ☐

Lemma 2.11. *Let $\pi : \mathcal{X} \to S$ be a flat family of surfaces such that the fibre $\mathcal{X}_s$ satisfies Serre's condition S_2 for each $s \in S$. Let $i \colon \mathcal{X}^o \subset \mathcal{X}$ be the inclusion of an open subset such that the complement has finite fibres. Then $i_*\mathcal{O}_{\mathcal{X}^o} = \mathcal{O}_{\mathcal{X}}$. Similarly, if $\mathcal{F}$ is a coherent sheaf on S then $i_*(\mathcal{O}_{\mathcal{X}^0} \otimes \pi^*\mathcal{F}) = \mathcal{O}_{\mathcal{X}} \otimes \pi^*\mathcal{F}$.*

Proof. For the first statement see, e.g., [H04], Lemma A.3, (the assumption that the fibres are semi log canonical is not used). The second statement follows from the first by dévissage. ☐

Definition 2.12. Let $B_0(\mathbb{Z})$ denote the set of points of B_0 with integral coordinates in an integral affine chart. We also write $B(\mathbb{Z}) = B_0(\mathbb{Z}) \cup \{0\}$.

Remark 2.13. Recall that by our definition an integral affine manifold has a distinguished atlas of charts with transition functions lying in $\mathrm{GL}(n,\mathbb{Z}) \ltimes \mathbb{Z}^n$. Thus the integral points of an integral affine manifold are well defined. Often a weaker definition is used where one only requires that the transition functions lie in $\mathrm{GL}(n,\mathbb{Z}) \ltimes \mathbb{R}^n$ (for example in the theory of integrable systems, see e.g., [GS06], Example 1.17). In this case one cannot define integral points. For an integrable system $(Y,\omega) \to B$, the induced affine structure on the base B is integral in the strong sense if the class of the symplectic form $[\omega] \in H^2_{\mathrm{dR}}(Y,\mathbb{R})$ is integral, see [GS06], Example 1.17 or [KS06], 3.1.1.

Given the description of Remark 2.6, the following lemma is obvious.

Lemma 2.14. *Suppose we are given φ and a monomial ideal $I \subset P$ such that Σ_I contains at least two rays, so that X_I^o is defined. Assume further that all cones in Σ_I are strictly convex.*

1. *If $x \in \operatorname{Spec} \Bbbk[P]/I$ is a closed point, let Σ_x denote the refinement of Σ_I obtained by adding any ray ρ of Σ for which $p_{\rho,\varphi} \notin \sqrt{I}$ but $z^{p_{\rho,\varphi}}$ lies in the maximal ideal of $\Bbbk[P]/I$ corresponding to x. Suppose Σ_x contains at least three rays. Then the fibre of $X_I^o \to \operatorname{Spec} \Bbbk[P]/I$ over x is $(\operatorname{Spec} \Bbbk[\Sigma_x]) \setminus \{0\}$. Here, $\Bbbk[\Sigma_x]$ denotes the $\Bbbk$-algebra with a $\Bbbk$-basis $\{z^m \,|\, m \in B(\mathbb{Z})\}$ with multiplication given exactly as in (1.4), and 0 is the closed point whose ideal is generated by $\{z^m \,|\, m \neq 0\}$.*
2. *Suppose that every cone in Σ_x is integral affine isomorphic to the first quadrant and Σ_x contains $p \geq 3$ rays. If $x \in \operatorname{Spec} \Bbbk[P]/I$ is a closed point, then the fibre of $X_I^o \to \operatorname{Spec} \Bbbk[P]/I$ over x is $\mathbb{V}_p^o$, the p-vertex minus the origin.*

Remark 2.15. If the hypothesis of Lemma 2.14, (2) is not satisfied then each closed fibre of $X_I^o \to \operatorname{Spec} \Bbbk[P]$ is a degenerate cusp [KSB88], 4.20. A *degenerate cusp* is a union of cyclic quotient singularities glued pairwise along toric boundary components in cyclic order.

Remark 2.16. We show that the convexity condition in Lemma 2.14 is satisfied in the cases we shall consider in this paper. We assume φ is defined using a pair (Y,D) as in Example 2.3, and $p_{\rho,\varphi} = \eta([D_\rho]) \in \sqrt{I}$ for at least two rays $\rho \in \Sigma$. Furthermore, we shall assume that $\sqrt{I}$ is a prime ideal.

Suppose $D_1,\ldots,D_r$ are consecutive components of the boundary such that $\eta([D_i]) \notin \sqrt{I}$, for some $r \leq n-2$. We consider two cases. First suppose the union of the cones $\sigma_{0,1},\ldots,\sigma_{r,r+1}$ is not convex. Then there exist $a_1,\ldots,a_r \in \mathbb{Z}_{\geq 0}$, such that $(\sum a_i D_i)^2 > 0$. (This follows from the toric case.) So $\sum a_i D_i$ lies in the interior of the cone of curves $\mathrm{NE}(Y)$. Since we are assuming $\sqrt{I}$ is prime, it follows that $\eta([C]) \notin \sqrt{I}$ for all $[C] \in \mathrm{NE}(Y)$, so in particular, $\eta([D_j]) \notin \sqrt{I}$ for all $j = 1,..,n$, contradicting our assumption.

Second suppose the union of the cones $\sigma_{0,1},\ldots,\sigma_{r,r+1}$ is a half-space. Then there exist $a_1,\ldots,a_r\in\mathbb{Z}_{>0}$ such that $(\sum a_iD_i)^2=0$, and $\sum a_iD_i$ defines a ruling $f\colon Y\to\mathbb{P}^1$. (Indeed, as in the toric case, there is a composition $g\colon Y\to Y'$ of contractions of (-1)-curves, with exceptional locus contained in $D_1+\cdots+D_r$, such that $D_1+\cdots+D_r$ contracts to a smooth rational curve F with $F^2=0$. Then the curve F defines a ruling of Y', so $g^*F=\sum a_iD_i$ defines a ruling of Y.) It follows that $\eta([C])\notin\sqrt{I}$ for all curves $C\subset Y$ contracted by f, again by our assumption of primality. Note that D_j is contracted by f for $j\neq r+1,n$, and D_{r+1},D_n are sections of f.

If (Y,D) is toric, the deformation $X_I^o\to\operatorname{Spec}\Bbbk[P]/I$ is a Mumford degeneration (see Lemma 2.10), and the closed fibres are isomorphic to the normal crossing surface $V(x_1x_3)\subset\mathbb{A}^2_{x_1,x_3}\times\mathbb{G}_{m,x_2}$.

If (Y,D) is not toric, then after a toric blowup $\pi\colon(\tilde{Y},\tilde{D})\to(Y,D)$, there exists a (-1)-curve $C\subset\tilde{Y}$, which is not contained in the boundary $\tilde{D}$ and is contracted by $f\circ\pi$. (Indeed, since $\rho(Y)>n-2$ we see that either there is a (-1)-curve C not contained in D and contracted by f, or $r=n-2$ and the fiber F through $p:=D_{n-1}\cap D_n$ is irreducible. In the latter case let $\pi\colon\tilde{Y}\to Y$ be the blowup of p and C be the strict transform of F.) We will later assume that either $\eta([D_j])\in\sqrt{I}$ for $j=1,\ldots,n$, or $\eta(\beta)\in\sqrt{I}$ for each $\mathbb{A}^1$-*class* β (see Definition 3.3), in particular, $\eta(\pi_*[C])\in\sqrt{I}$. So this case is not considered.

2.2. **Scattering diagrams on B.** Next we translate into algebraic geometry the instanton corrections. To construct our mirror family we will use the canonical scattering diagram $\mathfrak{D}^{\text{can}}$ (which is the translation of the instanton corrections associated to Maslov index zero disks), but as the regluing process works for any scattering diagram (and we will make use of this greater generality in the sequel [K3]), we carry it out for an arbitrary scattering diagram.

We continue with the notation of the previous sections, with (Y,D), $(B=B_{(Y,D)},\Sigma)$, P, and φ given. We also fix a monomial ideal $J\subset P$ such that $J=\sqrt{J}$. For any $\tau\in\Sigma$, $\tau\neq 0$, we have the monoid ring $\Bbbk[P_{\varphi_\tau}]$ and the ideal $J_{\tau,\tau}\subset\Bbbk[P_{\varphi_\tau}]$. We denote by

$$\widehat{\Bbbk[P_{\varphi_\tau}]}$$

the completion of the ring $\Bbbk[P_{\varphi_\tau}]$ with respect to the ideal $J_{\tau,\tau}$.

We will now define a *scattering diagram*, which encodes ways of modifying the construction of X_I^o.

Definition 2.17. A *scattering diagram* for the data (B,Σ), P,φ, and J is a set

$$\mathfrak{D}=\{(\mathfrak{d},f_{\mathfrak{d}})\}$$

where

1. $\mathfrak{d} \subset B$ is a ray in B with endpoint the origin with rational slope. $\mathfrak{d}$ may coincide with a ray of Σ, or lie in the interior of a two-dimensional cone of Σ.
2. Let $\tau_{\mathfrak{d}} \in \Sigma$ be the smallest cone containing $\mathfrak{d}$. Then $f_{\mathfrak{d}}$ is a formal sum
$$f_{\mathfrak{d}} = 1 + \sum_p c_p z^p \in \widehat{\Bbbk[P_{\varphi_{\tau_{\mathfrak{d}}}}]}$$
for $c_p \in \Bbbk$ and p running over elements of $P_{\varphi_{\tau_{\mathfrak{d}}}}$ such that $r(p) \neq 0$ and $r(p)$ is tangent to $\mathfrak{d}$. We further require that $\mathfrak{d}$ be an *outgoing ray*, in which case $r(p)$, viewed as a tangent vector at an interior point of $\mathfrak{d}$, always points towards the origin, or an *incoming ray*, in which case $r(p)$ always points away from the origin.
3. If $\dim \tau_{\mathfrak{d}} = 2$, or if $\dim \tau_{\mathfrak{d}} = 1$ and $p_{\tau_{\mathfrak{d}},\varphi} \notin J$, then $f_{\mathfrak{d}} \equiv 1 \mod J$.
4. For any ideal $I \subset P$ with $\sqrt{I} = J$, there are only a finite number of $(\mathfrak{d}, f_{\mathfrak{d}}) \in \mathfrak{D}$ such that $f_{\mathfrak{d}} \not\equiv 1 \mod I_{\tau_{\mathfrak{d}},\tau_{\mathfrak{d}}}$.

Construction 2.18. We now explain how a scattering diagram $\mathfrak{D}$ is used to modify the construction of X_I^o. Suppose we are given a scattering diagram $\mathfrak{D}$ for the data (B, Σ), P, φ and J, and an ideal I with $\sqrt{I} = J$. We assume that Σ_I contains at least two rays.

We will use the scattering diagram $\mathfrak{D}$ to both modify the definition of the rings $R_{\rho,I}$ as well as the gluings of the schemes defined by these rings. First, we modify the definition of $R_{\rho,I}$, setting for a ray $\rho \in \Sigma$ contained in two maximal cells $\sigma_\pm \in \Sigma$

$$R_{\rho,I} := R_{\rho,\sigma_-,I} \times_{(R_{\rho,\rho,I})_{f_\rho}} R_{\rho,\sigma_+,I}. \tag{2.9}$$

Here, $f_\rho \in R_{\rho,\rho,I}$ is defined by

$$f_\rho = \prod_{\substack{(\mathfrak{d}, f_{\mathfrak{d}}) \in \mathfrak{D} \\ \mathfrak{d} = \rho}} f_{\mathfrak{d}} \mod I_{\rho,\rho}.$$

This makes sense by condition (4) of the definition of scattering diagram. Furthermore, the maps in the fibred product are given by

$$R_{\rho,\sigma_-,I} \to (R_{\rho,\rho,I})_{f_\rho}$$

being the composition of the canonical surjection with the localization, while the map

$$R_{\rho,\sigma_+,I} \to (R_{\rho,\rho,I})_{f_\rho}$$

is given by the composition of the canonical surjection with the localization and then the map

$$\begin{aligned} (R_{\rho,\rho,I})_{f_\rho} &\to (R_{\rho,\rho,I})_{f_\rho} \\ z^p &\mapsto z^p f_\rho^{\langle n_\rho, r(p)\rangle}, \end{aligned}$$

where $n_\rho \in \Lambda_\rho^*$ annihilates the tangent space to ρ and is positive on σ_+. One notes easily that $R_{\rho,I}$ does not depend on which maximal cone containing ρ is called σ_+.

Set

$$U_{\rho,I} := \operatorname{Spec} R_{\rho,I}.$$

Lemma 2.19. *Let ρ be a ray in Σ corresponding to a component $D_\rho \subset D$ and $m_\rho \in \rho$ its primitive integral generator. Let $\rho \subset \sigma_\pm$, and let ρ_+ and ρ_- be the other boundary rays of σ_+ and σ_- respectively. Define*

$$R'_{\rho,I} := \frac{(\Bbbk[P]/I)[X_+, X_-, X^{\pm 1}]}{(X_+X_- - z^{p_{\rho,\varphi}} X^{-D_\rho^2} f_\rho)} \tag{2.10}$$

Here we view f_ρ as an element of $(\Bbbk[P]/I)[X^{\pm 1}]$ by identifying X with $z^{\varphi_\rho(m_\rho)} \in \Bbbk[P_{\varphi_\rho}]$. The map

$$(\Bbbk[P]/I)[X_+, X_-, X^{\pm 1}] \to R_{\rho,\sigma_-,I} \times R_{\rho,\sigma_+,I}$$

given by

$$\begin{aligned} X &\mapsto (z^{\varphi_\rho(m_\rho)}, z^{\varphi_\rho(m_\rho)}) \\ X_- &\mapsto (z^{\varphi_\rho(m_{\rho_-})}, f_\rho z^{\varphi_\rho(m_{\rho_-})}) \\ X_+ &\mapsto (f_\rho z^{\varphi_\rho(m_{\rho_+})}, z^{\varphi_\rho(m_{\rho_+})}) \end{aligned}$$

induces an isomorphism $h : R'_{\rho,I} \to R_{\rho,I}$. Furthermore, the natural map $U_{\rho,I} \to$ $\operatorname{Spec}(\Bbbk[P]/I)$ *is flat.*

Proof. First, it is easy to check that the image of the given map takes values in the fibre product $R_{\rho,I}$. Furthermore, $X_-X_+ - z^{p_{\rho,\varphi}} X^{-D_\rho^2} f_\rho$ is mapped to

$$(f_\rho z^{\varphi_\rho(m_{\rho_-})+\varphi_\rho(m_{\rho_+})} - f_\rho z^{p_{\rho,\varphi}} z^{-D_\rho^2\varphi_\rho(m_\rho)}, f_\rho z^{\varphi_\rho(m_{\rho_-})+\varphi_\rho(m_{\rho_+})} - f_\rho z^{p_{\rho,\varphi}} z^{-D_\rho^2\varphi_\rho(m_\rho)}).$$

However, $m_{\rho_-} + D_\rho^2 m_\rho + m_{\rho_+} = 0$ as elements of Λ_ρ, so one sees in fact that $\varphi_\rho(m_{\rho_-}) + \varphi_\rho(m_{\rho_+}) = p_{\rho,\varphi} - D_\rho^2\varphi_\rho(m_\rho)$. Thus the above element of the product is zero, and so our map induces a map $h : R'_{\rho,I} \to R_{\rho,I}$.

To show h is an isomorphism, we proceed as in [GS07], Lemma 2.34. Note that the rings $R_- = R_{\rho,\sigma_-,I}$ and $R_+ = R_{\rho,\sigma_+,I}$ are generated as $S = (\Bbbk[P]/I)[X^{\pm 1}]$-modules by $1, x^j, y^k$ for $j, k > 0$, where $x = z^{\varphi_\rho(m_{\rho_-})}$ and $y = z^{\varphi_\rho(m_{\rho_+})}$. Also the S-submodules of R_- (R_+) generated by x^j, $j \geq 0$ (y^k, $k \geq 0$) are free direct summands. So for $g_\pm \in R_\pm$, we can write uniquely

$$\begin{aligned} g_- &= \sum_{j\geq 0} a_j x^j + h_-(y) \\ g_+ &= \sum_{k\geq 0} b_k y^k + h_+(x), \end{aligned}$$

with $a_i, b_j \in S$ and $h_- \in S[y]$, $h_+ \in S[x]$, with $h_\pm(0) = 0$. So (g_-, g_+) lies in the fibred product if and only if

$$a_0 = b_0, \quad h_-(y) = \sum_{k>0} b_k f_\rho^k y^k, \quad h_+(x) = \sum_{j>0} a_j f_\rho^j x^j$$

as elements of $(R_{\rho,\rho,I})_{f_\rho}$. But if this is the case then (g_-, g_+) is the image of

$$\sum_{j\geq 0} a_j X_-^j + \sum_{k>0} b_k X_+^k.$$

This shows that the map h is surjective. Injectivity follows easily after noting that $R'_{\rho,I}$ is a free S-module with basis X_-^j, X_+^k, $j \geq 0$, $k > 0$.

The claimed flatness now is clear from the form of $R'_{\rho,I}$. $\square$

Remark 2.20. In the case of the canonical scattering diagram $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$, and $\sqrt{I} = I_p$ the monomial ideal for the Gross–Siebert locus, i.e., the toric stratum associated to a toric model $p : Y \to \bar{Y}$, the Lemma shows that $U_{\rho,I}$ is given by Equation (0.10), see Corollary 3.16.

Note that there are canonical maps

$$\psi_{\rho,\pm} : R_{\rho,I} \to R_{\sigma_\pm,\sigma_\pm,I}.$$

These are induced by the natural projections $R_{\rho,I} \to R_{\rho,\sigma_\pm,I}$ followed by the natural inclusions $R_{\rho,\sigma_\pm,I} \hookrightarrow R_{\sigma_\pm,\sigma_\pm,I}$. In terms of the description of $R_{\rho,I}$ as the ring $R'_{\rho,I}$, the map $\psi_{\rho,\pm}$ can be described as the localization of $R'_{\rho,I}$ at $X_\pm$. Thus $\operatorname{Spec} R_{\sigma_\pm,\sigma_\pm,I}$ is identified with an open subset $U_{\rho,\sigma_\pm,I}$ of $U_{\rho,I}$.

Note that if $p_{\rho,\varphi} \notin J$, then (3) of Definition 2.17 tells us that f_ρ is already invertible in $R_{\rho,\rho,I}$, so the localization isn't necessary. However, $\psi_{\rho,+} \circ \psi_{\rho,-}^{-1}$, where defined, is not the trivial identification used in Definition-Lemma 2.8, but is twisted by the automorphism $z^p \mapsto z^p f_\rho^{-\langle n_\rho, r(p)\rangle}$.

Next, consider $(\mathfrak{d}, f_\mathfrak{d}) \in \mathfrak{D}$ with $\mathfrak{d} \subset \sigma \in \Sigma_{\max}$ and $\mathfrak{d}$ not contained in a ray of Σ. Let γ be a path in B_0 which crosses $\mathfrak{d}$ transversally at time t_0. Then define

$$\theta_{\gamma,\mathfrak{d}} : R_{\sigma,\sigma,I} \to R_{\sigma,\sigma,I}$$

by

$$\theta_{\gamma,\mathfrak{d}}(z^p) = z^p f_\mathfrak{d}^{\langle n_\mathfrak{d}, r(p)\rangle}$$

where $n_\mathfrak{d} \in \Lambda_\sigma^*$ is primitive and satisfies, with m a non-zero tangent vector of $\mathfrak{d}$,

$$\langle n_\mathfrak{d}, m\rangle = 0, \quad \langle n_\mathfrak{d}, \gamma'(t_0)\rangle < 0.$$

Note that $f_\mathfrak{d}$ is invertible in $R_{\sigma,\sigma,I}$ since $f_\mathfrak{d} \equiv 1 \mod J_{\sigma,\sigma}$, so $f_\mathfrak{d} - 1$ is nilpotent in $R_{\sigma,\sigma,I}$.



FIGURE 2.1. The path γ. The solid lines indicate the fan, the dotted lines are additional rays in $\mathfrak{D}$. The solid lines may also support rays in $\mathfrak{D}$

Let $\mathfrak{D}_I \subset \mathfrak{D}$ be the finite set of rays $(\mathfrak{d}, f_{\mathfrak{d}})$ with $f_{\mathfrak{d}} \not\equiv 1 \mod I_{\tau_{\mathfrak{d}},\tau_{\mathfrak{d}}}$. For a path γ wholly contained in the interior of $\sigma \in \Sigma_{\max}$ and crossing elements of $\mathfrak{D}_I$ transversally, we define

$$\theta_{\gamma,\mathfrak{D}} := \theta_{\gamma,\mathfrak{d}_n} \circ \cdots \circ \theta_{\gamma,\mathfrak{d}_1},$$

where γ crosses precisely the elements $(\mathfrak{d}_1, f_{\mathfrak{d}_1}), \ldots, (\mathfrak{d}_n, f_{\mathfrak{d}_n})$ of $\mathfrak{D}_I$, in the given order. Note that if two rays $\mathfrak{d}_i, \mathfrak{d}_{i+1}$ in fact coincide as subsets of B, then $\theta_{\gamma,\mathfrak{d}_i}$ and $\theta_{\gamma,\mathfrak{d}_{i+1}}$ commute, so the ordering is not important for overlapping rays.

To construct $X^o_{I,\mathfrak{D}}$, we modify the gluings of the sets $U_{\rho,I}$ along the open subsets $U_{\rho,\sigma,I}$. For each $\sigma \in \Sigma_{\max}$ with edges ρ, ρ', we have canonical identifications

$$U_{\rho,\sigma,I} = \operatorname{Spec} R_{\sigma,\sigma,I} = U_{\rho',\sigma,I}.$$

Hence we can modify this gluing via any automorphism. We do this by choosing a path $\gamma : [0,1] \to B$ whose image is contained in the interior of σ, with $\gamma(0)$ a point in σ close to ρ and $\gamma(1) \in \sigma$ close to ρ', chosen so that γ crosses every ray $(\mathfrak{d}, f_{\mathfrak{d}})$ of $\mathfrak{D}_I$ with $\tau_{\mathfrak{d}} = \sigma$ exactly once, see Figure 2.1.

We then obtain an automorphism

$$\theta_{\gamma,\mathfrak{D}} : R_{\sigma,\sigma,I} \to R_{\sigma,\sigma,I},$$

hence an isomorphism

$$\theta_{\gamma,\mathfrak{D}} : U_{\rho',\sigma,I} \to U_{\rho,\sigma,I}.$$

Gluing together the open sets $U_{\rho',\sigma,I} \subset U_{\rho',I}$ and $U_{\rho,\sigma,I} \subset U_{\rho,I}$ via $\theta_{\gamma,\mathfrak{D}}$, we obtain a scheme $X^o_{I,\mathfrak{D}}$. $\square$

2.3. **Broken lines.** We continue to fix a rational surface with anti-canonical cycle (Y, D) as usual, giving (B, Σ), as well as a monoid P, a P-convex function φ on B, $J \subset P$ an ideal with $\sqrt{J} = J$, and a scattering diagram $\mathfrak{D}$ for this data. As explained in the previous section, $\mathfrak{D}$ determines an *instanton-corrected* deformation $X^o_{I,\mathfrak{D}}$ of $\mathbb{V}^o_n$. As explained in §0.6.2, this extends to a deformation of $\mathbb{V}_n$ if and only if the coordinate functions x_i on $\mathbb{V}^o_n$ lift to $X^o_{I,\mathfrak{D}}$. The symplectic heuristic suggests a natural way of producing such lifts. Translated into algebraic geometry this leads to the notion of *broken line*.

Definition 2.21. Let B be an integral affine manifold. An integral affine map $\gamma : (t_1, t_2) \to B$ from an open interval (t_1, t_2) is a continuous map such that for any integral affine coordinate chart $\psi : U \to \mathbb{R}^n$ of B, $\psi \circ \gamma : \gamma^{-1}(U) \to \mathbb{R}^n$ is integral affine, i.e., is given by $t \mapsto tv + b$ for some $v \in \mathbb{Z}^n$ and $b \in \mathbb{R}^n$.

Note that for an integral affine map, $\gamma'(t) \in \Lambda_{B,\gamma(t)}$.

Definition 2.22. A *broken line* γ in B for $q \in B_0(\mathbb{Z})$ with endpoint $Q \in B_0$ is a proper continuous piecewise integral affine map $\gamma : (-\infty, 0] \to B_0$, together with, for each $L \subset (-\infty, 0]$ a maximal connected domain of linearity of γ, a choice of monomial $m_L = c_L z^{q_L}$ where $c_L \in \Bbbk$ and $q_L \in \Bbbk[\Gamma(L, \gamma^{-1}(\mathcal{P})|_L)]$, satisfying the following properties.

1. For the unique unbounded domain of linearity L, $\gamma|_L$ goes off to infinity in a cone $\sigma \in \Sigma_{\max}$ as $t \to -\infty$, and $q \in \sigma$. Furthermore, using the identification of the stalk $\mathcal{P}_x$ for $x \in \sigma$ with $\mathcal{P}_\sigma$, $m_L = z^{\varphi_\sigma(q)}$ (see Remark 2.1).
2. For each L and $t \in L$, $-r(q_L) = \gamma'(t)$. Also $\gamma(0) = Q \in B_0$.
3. Let $t \in (-\infty, 0)$ be a point at which γ is not linear, passing from domain of linearity L to L'. If $\gamma(t) \in \tau \in \Sigma$, then $\mathcal{P}_{\gamma(t)} = \mathcal{P}_\tau$, so that we can view $q_L \in \mathcal{P}_\tau$ and $r(q_L) \in \Lambda_\tau$. Let $\mathfrak{d}_1, \ldots, \mathfrak{d}_p \in \mathfrak{D}$ be the rays of $\mathfrak{D}$ that contain $\gamma(t)$, with attached functions $f_{\mathfrak{d}_j}$. Let $n = n_{\mathfrak{d}_j}$ be the element of Λ^*_τ used to define $\theta_{\gamma,\mathfrak{d}_j}$. Expand
$$\prod_{j=1}^{p} f_{\mathfrak{d}_j}^{\langle n, r(q_L)\rangle} \tag{2.11}$$
as a formal power series in $\widehat{\Bbbk[P_{\varphi_\tau}]}$. Then there is a term cz^q in this sum with
$$m_{L'} = m_L \cdot (cz^q).$$

Remark 2.23. Using the notation of item (3) above, by item (2) of the definition,
$$\langle n, r(q_L)\rangle > 0. \tag{2.12}$$
This is vital to interpret (2.11). Indeed, if τ is a ray, $f_{\mathfrak{d}_i}$ need not be invertible in $\widehat{\Bbbk[P_{\varphi_\tau}]}$, so (2.12) tells us that (2.11) makes sense in this ring.

The next lemma and corollary are vital for interpreting the monomials m_L:

Lemma 2.24. *Let* $\sigma_-, \sigma_+ \in \Sigma_{\max}$ *be the two maximal cones containing the ray* $\rho \in \Sigma$. *If* $q \in P_{\varphi_{\sigma_-}}$ *with* $-r(q) \in \operatorname{Int}(\rho^{-1}\sigma_+) \subset \Lambda_\rho \otimes_{\mathbb{Z}} \mathbb{R}$, *then*

$$q \in P_{\varphi_\rho} = P_{\varphi_{\sigma_-}} \cap P_{\varphi_{\sigma_+}}.$$

Proof. By the definitions there exist $p, p_{\rho,\varphi} \in P$ and $n_\rho \in \Lambda_\rho^*$ annihilating the tangent space to ρ and positive on σ_+ such that

$$q = \varphi_{\sigma_-}(r(q)) + p$$
$$\varphi_{\sigma_+}(-r(q)) = \varphi_{\sigma_-}(-r(q)) + \langle n_\rho, -r(q)\rangle p_{\rho,\varphi}.$$

Since $\langle n_\rho, -r(q)\rangle > 0$,

$$q = \varphi_{\sigma_+}(r(q)) + p + \langle n_\rho, -r(q)\rangle p_{\rho,\varphi} \in P_{\varphi_{\sigma_+}}.$$

□

An immediate consequence of this lemma is

Corollary 2.25. (1) *Let* $\gamma : [t_1, t_2] \to B_0$ *be integral affine. Suppose that* $\gamma(t_1) \in \tau_1$, $\gamma(t_2) \in \tau_2$. *Suppose also we are given a section* $q \in \Gamma(\gamma^{-1}\mathcal{P})$ *such that* $-r(q) = \gamma'(t)$ *for each* t. *If*

$$q(t_1) \in P_{\varphi_{\tau_1}} \subset \mathcal{P}_{\tau_1} = \mathcal{P}_{\gamma(t_1)},$$

then

$$q(t_2) \in P_{\varphi_{\tau_2}} \subset \mathcal{P}_{\tau_2} = \mathcal{P}_{\gamma(t_2)}.$$

(2) *If* γ *is a broken line,* $t \in L$ *a maximal domain of linearity with* $\gamma(t) \in \tau$, *then*

$$q_L \in P_{\varphi_\tau} \subset \mathcal{P}_\tau = \mathcal{P}_{\gamma(t)}.$$

Proof. The first item follows immediately from the lemma. The second item follows from the fact that if $t \ll 0$ lies in the unbounded domain of linearity with $\gamma(t) \in \sigma$, then $m_L = z^{\varphi_\tau(q)} \in P_{\varphi_\sigma}$ by construction. Then this holds for all t by item (1) and Definition 2.22, (3). □

The convexity of φ puts further restrictions on the monomial decorations of a broken line.

Definition 2.26. Let $J \subset P$ be a proper monoid ideal. For $p \in J$ there exists a maximal $k \geq 1$ such that $p = p_1 + \cdots + p_k$ with $p_i \in J$. We define $\operatorname{ord}_J(p)$ to be this maximum, and $\operatorname{ord}_J(p) = 0$ if $p \in P \setminus J$.

For $x \in \tau$, $q \in P_{\varphi_\tau}$, define $\mathrm{ord}_{J,x}(q) := \mathrm{ord}_J(q - \varphi_\tau(r(q)))$. This measures how high q is above the graph of φ_τ. If γ is a broken line and $t \in L$ a maximal domain of linearity, define

$$\mathrm{ord}_{J,\gamma}(t) = \mathrm{ord}_{J,\gamma(t)}(q_L),$$

using $\gamma(t) \in \tau$ and $q_L \in P_{\varphi_\tau} \subset \mathcal{P}_{\gamma(t)}$.

Lemma 2.27. *Let γ be a broken line. Then if $t < t'$,*

$$\mathrm{ord}_{J,\gamma}(t) \leq \mathrm{ord}_{J,\gamma}(t'),$$

with strict inequality if either t and t' lie in different domains of linearity or for some t'' with $t < t'' < t'$, $\gamma(t'')$ lies in a ray $\rho \in \Sigma$ with bending parameter $p_{\rho,\varphi} \in J$.

Proof. This is immediate from the definitions and the proof of Lemma 2.24. $\square$

Definition 2.28. For I an ideal in P with $\sqrt{I} = J$, let

$$\mathrm{Supp}_I(\mathfrak{D}) := \bigcup_{\mathfrak{d}} \mathfrak{d}$$

where the union is over all $(\mathfrak{d}, f_{\mathfrak{d}}) \in \mathfrak{D}$ such that $f_{\mathfrak{d}} \not\equiv 1 \mod I_{\tau_{\mathfrak{d}},\tau_{\mathfrak{d}}}$. By Definition 2.17, (4), this is a finite union.

Definition 2.29. Let I be an ideal of P with $\sqrt{I} = J$, and let $Q \in B \setminus \mathrm{Supp}_I(\mathfrak{D})$, $Q \in \tau \in \Sigma$. For $q \in B_0(\mathbb{Z})$, define

$$\mathrm{Lift}_Q(q) := \sum_{\gamma} \mathrm{Mono}(\gamma) \in \Bbbk[P_{\varphi_\tau}]/I \cdot \Bbbk[P_{\varphi_\tau}], \tag{2.13}$$

where the sum is over all broken lines γ for q with endpoint Q, and $\mathrm{Mono}(\gamma)$ denotes the monomial attached to the last domain of linearity of γ. The fact that $\mathrm{Lift}_Q(q)$ lies in the stated ring follows from:

Lemma 2.30. *Let $Q \in \sigma \in \Sigma_{\max}$, $q \in B_0(\mathbb{Z})$. Let I be an ideal with $\sqrt{I} = J$. Assume that $p_{\rho,\varphi} \in J$ for at least one ray $\rho \in \Sigma$. Then the following hold:*

(1) *The collection of γ in Definition 2.29 with*

$$\mathrm{Mono}(\gamma) \notin I \cdot \Bbbk[P_{\varphi_\sigma}]$$

is finite.

(2) *If one boundary ray of the connected component of $B \setminus \mathrm{Supp}_I(\mathfrak{D})$ containing Q is a ray $\rho \in \Sigma$, then $\mathrm{Mono}(\gamma) \in \Bbbk[P_{\varphi_\rho}]$, and the collection of γ with*

$$\mathrm{Mono}(\gamma) \notin I \cdot \Bbbk[P_{\varphi_\rho}]$$

is finite.

Proof. Note there is some k such that $J^k \subset I$ because $\Bbbk[P]$ is Noetherian. If γ is a broken line with $\mathrm{Mono}(\gamma) \notin I \cdot \Bbbk[P_{\varphi_\sigma}]$, then it is easy to see that γ crosses the rays of Σ in a cyclic order, and so in any set of at least n consecutive rays of Σ that it crosses, there is at least one ray ρ with $p_{\rho,\varphi} \in J$. By Lemma 2.27, $\mathrm{ord}_{J,\gamma}$ increases every time γ crosses such a ray, and also every time γ bends. Once $\mathrm{ord}_{J,\gamma} \geq k$, $\mathrm{Mono}(\gamma) \in I \cdot \Bbbk[P_{\varphi_\sigma}]$. Hence there is an absolute bound on the number of rays of Σ that γ can cross, and the number of times γ can bend. Once the initial direction of γ is fixed, it's easy to see that the set of all possible broken lines satisfying the condition of (1) is finite, yielding the finiteness of (1).

The argument for the finiteness statement in (2), once the first part of (2) is established, is the same. For the first part of (2), consider a broken line γ contributing to $\mathrm{Lift}_Q(q)$. We take $Q \in \sigma_+$, in the notation of Lemma 2.24. Write $\mathrm{Mono}(\gamma) = c_L z^{q_L}$. If $r(q_L) \in \rho^{-1}\sigma_+$ then the statement follows from Lemma 2.24. Otherwise (by the definition of broken line) γ crosses ρ, which is the last ray of Σ and the last ray of $\mathrm{Supp}_I(\mathfrak{D})$ it crosses before reaching Q. Now the result follows from Lemma 2.24 and the definition of broken line. □

Remark 2.31 (Heuristic meaning of $\mathrm{Lift}_Q(q)$). Assume $\mathfrak{D}$ is the canonical scattering diagram $\mathfrak{D}^{\mathrm{can}}$. Let $q \in B_0(\mathbb{Z})$ and $q = mv$, for $v \in B_0(\mathbb{Z})$ the primitive integral point on the ray $\rho_q := \mathbb{R}_{\geq 0} q \subset B$. Then v determines a boundary divisor D_v on a toric blowup $(\tilde{Y}, \tilde{D})$ of (Y, D). Let γ be a broken line contributing to $\mathrm{Lift}_Q(q)$, with $\mathrm{Mono}(\gamma) = cz^w \in \Bbbk[\mathcal{P}_Q]$. The point Q heuristically corresponds to a Lagrangian $L \subset U := Y \setminus D$ and γ to a holomorphic disk with boundary on L and meeting $\tilde{D}$ in a single point, lying in the interior D_v^o of D_v, with multiplicity m. Then $w \in \mathcal{P}_Q = H_2(Y, L)$ is the relative homology class of the disk, and the monomial function z^w (on the SYZ dual X of $U \to B$ viewed now as the instanton corrected moduli space of special Lagrangians with $U(1)$ connection) is as defined in (0.6). The monomials z^q are only locally defined, but we should expect that the sum $\mathrm{Lift}_Q(q)$ makes global sense. For example, in the special case $q = v_i$ it corresponds to the canonical global function ϑ_i obtained from counting Maslov index two disks meeting D_i. In particular, we expect the $\mathrm{Lift}_Q(q)$ to transform the same way as the z^q, i.e., by change of coordinate according to the symplectomorphisms coming from the canonical scattering diagram $\mathfrak{D}^{\mathrm{can}}$. In this section we are considering an arbitrary scattering diagram, so we introduce the necessary patching conditions as an axiom. In subsequent sections we'll prove that $\mathfrak{D}^{\mathrm{can}}$ has this property.

Definition 2.32. Assume that $p_{\rho,\varphi} \in J$ for at least one ray $\rho \in \Sigma$. We say a scattering diagram $\mathfrak{D}$ is *consistent* if for all ideals $I \subset P$ with $\sqrt{I} = J$ and for all $q \in B_0(\mathbb{Z})$, the following holds. Let $Q \in B_0$ be chosen so that the line joining the origin and Q has irrational slope, and $Q' \in B_0$ similarly. Then:

(1) If $Q, Q' \in \sigma \in \Sigma_{\max}$, then we can view $\mathrm{Lift}_Q(q)$ and $\mathrm{Lift}_{Q'}(q)$ as elements of $R_{\sigma,\sigma,I}$, and as such, we have

$$\mathrm{Lift}_{Q'}(q) = \theta_{\gamma,\mathfrak{D}}(\mathrm{Lift}_Q(q))$$

for γ a path contained in the interior of σ connecting Q to Q'.

(2) If $Q \in \sigma_-$ and $Q' \in \sigma_+$ with $\sigma_\pm \in \Sigma_{\max}$ and $\rho = \sigma_+ \cap \sigma_-$ a ray, and furthermore Q and Q' are contained in connected components of $B \setminus \mathrm{Supp}_I(\mathfrak{D})$ whose closures contain ρ, then viewing $\mathrm{Lift}_Q(q) \in R_{\rho,\sigma_-,I}$, $\mathrm{Lift}_{Q'}(q) \in R_{\rho,\sigma_+,I}$, the pair

$$(\mathrm{Lift}_Q(q), \mathrm{Lift}_{Q'}(q))$$

lies in the ring $R_{\rho,I}$.

Of course the definition is introduced so that the following holds:

Theorem 2.33. *Let φ be a multivalued piecewise linear function satisfying the hypotheses of Lemma 2.14. Let $\mathfrak{D}$ be a consistent scattering diagram and $I \subset P$ an ideal with $\sqrt{I} = J$. Set*

$$X_I := \operatorname{Spec} \Gamma(X^o_{I,\mathfrak{D}}, \mathcal{O}_{X^o_{I,\mathfrak{D}}}).$$

Since $X^o_{I,\mathfrak{D}}$ has the structure of a scheme over $\operatorname{Spec} \Bbbk[P]/I$, so does X_I, which we write as

$$f_I : X_I \to \operatorname{Spec} \Bbbk[P]/I.$$

Suppose that for every closed point $x \in \operatorname{Spec} \Bbbk[P]/I$, Σ_x satisfies the hypotheses of Lemma 2.14, (1). Then

(1) *X_I contains $X^o_{I,\mathfrak{D}}$ as an open subset, f_I is flat with fibre over a closed point x of $\operatorname{Spec} \Bbbk[P]/I$ a surface, isomorphic to $\operatorname{Spec} \Bbbk[\Sigma_x]$ in general and to the p-vertex $\mathbb{V}_p$ for some p if the hypothesis of Lemma 2.14, (2), holds.*

(2) *For each $q \in B(\mathbb{Z})$, there is a section $\vartheta_q \in \Gamma(X_I, \mathcal{O}_{X_I})$, and the set $\{\vartheta_q \,|\, q \in B(\mathbb{Z})\}$ is a free $(\Bbbk[P]/I)$-module basis for $\Gamma(X_I, \mathcal{O}_{X_I})$.*

Proof. We define $\vartheta_0 = 1$. We now construct ϑ_q for $q \in B_0(\mathbb{Z})$: this is the main point of Definition 2.32. For each ray $\rho \in \Sigma$ contained in $\sigma_\pm \in \Sigma_{\max}$, choose two points $Q^\pm_\rho \in B$, one each in the two connected components of $B \setminus (\mathrm{Supp}_I(\mathfrak{D}) \cup \rho)$ which are adjacent to ρ, with $Q^+_\rho \in \sigma_+$ and $Q^-_\rho \in \sigma_-$. Then $\mathrm{Lift}_{Q^\pm_\rho}(q)$ is a well-defined element of $R_{\rho,\sigma_\pm,I}$, independent of the particular choice of $Q^\pm_\rho$: given a choice say of $Q = Q^+_\rho$ and another choice Q', we take a path γ connecting Q and Q' contained wholly in the connected component of $B \setminus (\mathrm{Supp}_I(\mathfrak{D}) \cup \rho)$ containing Q and Q'. By Definition 2.32, (1), it then follows that $\mathrm{Lift}_Q(q) = \mathrm{Lift}_{Q'}(q)$. Furthermore, by Definition 2.32, (2),

$(\mathrm{Lift}_{Q_\rho^-}(q), \mathrm{Lift}_{Q_\rho^+}(q))$ defines an element $\mathrm{Lift}_\rho(q) \in R_{\rho,I}$. It then follows via another application of Definition 2.32, (1), applied to the path of Figure 2.1, that if ρ, ρ' are adjacent rays in Σ, then $\mathrm{Lift}_\rho(q)$ and $\mathrm{Lift}_{\rho'}(q)$ glue under the identification of open subsets of $U_{\rho,I}$ and $U_{\rho',I}$ given by $\theta_{\gamma,\mathfrak{D}}$. Thus all these elements of the rings $R_{\rho,I}$ glue to give a regular function on $X^o_{I,\mathfrak{D}}$, by construction of this latter space. This regular function is what we call ϑ_q. By the definition of X_I, $\vartheta_q \in \Gamma(X_I, \mathcal{O}_{X_I})$.

Now note that $X^o_{J,\mathfrak{D}} = X^o_J$ as defined in §2.1. Indeed, for any $(\mathfrak{d}, f_\mathfrak{d}) \in \mathfrak{D}$ with $\dim \tau_\mathfrak{d} = 2$ we have $f_\mathfrak{d} \equiv 1 \mod J$, so the open sets $U_{\rho_i,J}$, $U_{\rho_{i+1},J}$ are glued trivially. Similarly, if $\dim \tau_\mathfrak{d} = 1$ and $p_{\tau_\mathfrak{d},\varphi} \notin J$, then $f_\mathfrak{d} \equiv 1 \mod J$, so the sets $U_{\tau_\mathfrak{d},J}$ coincide in both constructions. If $p_{\tau_\mathfrak{d},\varphi} \in J$, then by Lemma 2.19, again these open sets coincide in both constructions.

We verify explicitly that the map

$$\bigoplus_{q\in B(\mathbb{Z})} (\Bbbk[P]/J)\cdot\vartheta_q \to \Gamma(X^o_J, \mathcal{O}_{X^o_J}) \tag{2.14}$$

is an isomorphism. We assume that Σ_I has at least three rays, leaving the straightforward modifications in the case of two rays to the reader. We first prove surjectivity. Write $U_i := U_{\rho_i,J}$ and $U_{i,i+1} := U_i \cap U_{i+1}$. Let $h \in \Gamma(X^o_J, \mathcal{O}_{X^o_J})$ be a global function on X^o_J. Observe that X^o_J is reduced and $\bigcup U_{i,i+1}$ is a dense open set. So it suffices to show that the restriction of h to this open set coincides with the restriction of a $(\Bbbk[P]/J)$-linear combination of the ϑ_q. Write $h_{i,i+1} = h|_{U_{i,i+1}}$ and $h_{i,i+1} = \sum_{q\in\Lambda_{\sigma_{i,i+1}}} h_{i,i+1,q}$, where $h_{i,i+1,q}$ is a sum of monomials in $r^{-1}(q) \subset \mathcal{P}_{\sigma_{i,i+1}}$. Then the $h_{i,i+1,q}$ satisfy the following compatibility. Let $\gamma\colon [0,1] \to B_0$ be a path given by a straight line segment in the direction q with endpoints $\gamma(0)$, $\gamma(1)$ lying in the interiors of $\sigma_{i,i+1}$ and $\sigma_{j,j+1}$. Then $h_{i,i+1,q}$ is obtained from $h_{j,j+1,q}$ by parallel transport in $\mathcal{P}$ along γ^{-1} followed by reduction modulo J. (In the case that γ crosses a single ray of Σ, this follows from the definition of U_j and the inclusions $U_{j-1,j} \subset U_j$, $U_{j,j+1} \subset U_j$. The general case follows by induction.) In particular, if γ crosses a ray of Σ_I then $h_{i,i+1,q} = 0$ (cf. Lemma 2.27). Let σ be the maximal cone of Σ_I containing $\sigma_{i,i+1}$. Then the above compatibility yields the following. If the ray $\mathbb{R}_{\geq 0}\cdot q$ is not contained in σ then $h_{i,i+1,q} = 0$. Otherwise, let $\sigma_{j,j+1} \subset \sigma$ be a maximal cone of Σ containing q. Then $h_{i,i+1,q}$ is obtained from $h_{j,j+1,q}$ by parallel transport in $\mathcal{P}$ along a path in σ from $\sigma_{j,j+1}$ to $\sigma_{i,i+1}$. Now for $q \in B(\mathbb{Z})$ let $\sigma_{j,j+1}$ be a maximal cone of Σ containing q and write $h_{j,j+1,q} = a_q z^{\varphi(q)}$ where $a_q \in \Bbbk[P]/J$. We deduce that $h = \sum_{q\in B(\mathbb{Z})} a_q\vartheta_q$. Thus (2.14) is surjective. Finally, injectivity of (2.14) is clear by restriction to the open sets $U_{i,i+1}$. Hence (2.14) is an isomorphism as required.

It follows that $X_J := \operatorname{Spec}\Gamma(X^o_J, \mathcal{O}_{X^o_J})$ is flat over $\operatorname{Spec}(\Bbbk[P]/J)$ and the fiber over a closed point x is given by $\operatorname{Spec}\Bbbk[\Sigma_x]$.

Now let I be a J-primary ideal. Let $i\colon X_J^o \subset X_J$ be the inclusion. Define a ringed space X_I' with underlying topological space X_J by $\mathcal{O}_{X_I'} := i_*\mathcal{O}_{X^o_{I,\mathfrak{D}}}$. Then the natural map $\mathcal{O}_{X_I'} \to \mathcal{O}_{X_J}$ is surjective by the existence of the lifts ϑ_q. Thus $X_I'/\operatorname{Spec}(\Bbbk[P]/I)$ is a flat deformation of $X_J/\operatorname{Spec}(\Bbbk[P]/J)$ by Lemma 2.34 below. Now since X_J is affine it follows that X_I' is also affine, so $X_I' = X_I := \operatorname{Spec}\Gamma(X^o_{I,\mathfrak{D}}, \mathcal{O}_{X^o_{I,\mathfrak{D}}})$.

We showed above that the ϑ_q form a $(\Bbbk[P]/J)$-module basis of $\Gamma(X_J, \mathcal{O}_{X_J})$. Now since $X_I/\operatorname{Spec}(\Bbbk[P]/I)$ is a flat infinitesimal deformation of $X_J/\operatorname{Spec}(\Bbbk[P]/J)$ it follows that the ϑ_q form a $\Bbbk[P]/I$-module basis of $\Gamma(X_I, \mathcal{O}_{X_I})$, see Lemma 2.35 below. □

Lemma 2.34. *Let X_0/S_0 be a flat family of surfaces such that the fibres satisfy Serre's condition S_2. Let $i\colon X_0^o \subset X_0$ be the inclusion of an open subset such that the complement has finite fibres. Note that $i_*\mathcal{O}_{X_0^o} = \mathcal{O}_{X_0}$ by Lemma 2.11.*

Let $S_0 \subset S$ be an infinitesimal thickening of S_0 and let $X^o \to S$ be a flat deformation of X_0^o/S_0 over S. Define a family of ringed spaces $X \to S$ by $\mathcal{O}_X := i_*\mathcal{O}_{X^o}$.

Then X/S is a flat deformation of X_0/S_0 (that is, X/S is flat and $X_0 = X \times_S S_0$) if and only if the map

$$\mathcal{O}_X := i_*\mathcal{O}_{X^o} \to i_*\mathcal{O}_{X_0^o} = \mathcal{O}_{X_0} \tag{2.15}$$

is surjective.

Proof. The condition is clearly necessary.

Conversely, suppose (2.15) is surjective. Let $\mathcal{I} \subset \mathcal{O}_S$ be the nilpotent ideal defining $S_0 \subset S$. Let X_n^o/S_n denote the nth order infinitesimal thickening of X_0/S_0 determined by X^o/S, that is, $\mathcal{O}_{X_n^o} = \mathcal{O}_{X^o}/\mathcal{I}^{n+1}\cdot\mathcal{O}_{X^o}$ and $\mathcal{O}_{S_n} = \mathcal{O}_S/\mathcal{I}^{n+1}$. Define X_n/S_n by $\mathcal{O}_{X_n} := i_*\mathcal{O}_{X_n^o}$. Note that $\mathcal{O}_{X_n} \to \mathcal{O}_{X_0}$ is surjective because $\mathcal{O}_X \to \mathcal{O}_{X_0}$ is surjective by assumption. We show by induction on n that X_n/S_n is a flat deformation of X_0/S_0. For $n = 0$ there is nothing to prove. Suppose the induction hypothesis is true for n. Since X^o_{n+1}/S_{n+1} is flat (being the restriction of the flat family X^o/S to S_{n+1}) we have a short exact sequence

$$0 \to \mathcal{I}^{n+1}/\mathcal{I}^{n+2} \otimes \mathcal{O}_{X_0^o} \to \mathcal{O}_{X^o_{n+1}} \to \mathcal{O}_{X^o_n} \to 0.$$

Applying i_* we obtain an exact sequence

$$0 \to i_*(\mathcal{I}^{n+1}/\mathcal{I}^{n+2} \otimes \mathcal{O}_{X_0^o}) \to \mathcal{O}_{X_{n+1}} \to \mathcal{O}_{X_n}$$

By Lemma 2.11 the first term is equal to $\mathcal{I}^{n+1}/\mathcal{I}^{n+2} \otimes \mathcal{O}_{X_0}$. Moreover, the last arrow is surjective because $\mathcal{O}_{X_{n+1}} \to \mathcal{O}_{X_0}$ is surjective, $\mathcal{O}_{X_n}/\mathcal{I}\cdot\mathcal{O}_{X_n} = \mathcal{O}_{X_0}$ by the induction hypothesis, and $\mathcal{I}$ is nilpotent. So we have an exact sequence

$$0 \to \mathcal{I}^{n+1}/\mathcal{I}^{n+2} \otimes \mathcal{O}_{X_0} \to \mathcal{O}_{X_{n+1}} \to \mathcal{O}_{X_n} \to 0. \tag{2.16}$$

It follows that $\mathcal{O}_{X_{n+1}}/\mathcal{I}^{n+1}\cdot\mathcal{O}_{X_{n+1}} = \mathcal{O}_{X_n}$ (using again that $\mathcal{O}_{X_{n+1}} \to \mathcal{O}_{X_0}$ is surjective). Now by [Ma89], Theorem 22.3, p. 174, the exact sequence (2.16) shows that X_{n+1}/S_{n+1} is a flat deformation of X_0/S_0. $\square$

Lemma 2.35. *Let $A \to B$ be a flat homomorphism of Noetherian rings and $I \subset A$ a nilpotent ideal. Suppose given a set S of elements of B such that the reductions of the elements of S form an A/I-module basis of B/IB. Then S is an A-module basis of B.*

Proof. Since I is nilpotent and S generates B/IB it is clear that S spans B. So we have an exact sequence

$$0 \to K \to A^S \to B \to 0.$$

Tensoring with A/I we obtain an exact sequence

$$0 \to K/IK \to (A/I)^S \to B/IB \to 0$$

using flatness of B over A. We deduce that $K/IK = 0$ by our assumption, hence $K = 0$ because I is nilpotent. $\square$

Proposition 2.36. *Let $X_I/S_I := \operatorname{Spec}(\Bbbk[P]/I)$ be the family of Theorem 2.33. Then the relative dualizing sheaf ω_{X_I/S_I} is trivial. It is generated by the global section Ω given on local patches $U_\rho = \operatorname{Spec} R_{\rho,I}$ by $\operatorname{dlog} X^- \wedge \operatorname{dlog} X = \operatorname{dlog} X \wedge \operatorname{dlog} X^+$. Here we have used the notation of Lemma 2.19 and have labelled the rays ρ_-, ρ, ρ_+ in anticlockwise order.*

Proof. By the adjunction formula for the closed embedding

$$U_\rho \subset \mathbb{A}^2_{X_-,X_+} \times \mathbb{G}_{m,X} \times S_I,$$

the dualizing sheaf ω_{X_I/S_I} is freely generated over U_ρ by the local section in the statement. These sections patch to give a generator Ω of $\omega_{X^o_{I,\mathfrak{D}}/S_I}$ because the scattering automorphisms preserve the torus invariant differentials. Both ω_{X_I/S_I} and $\mathcal{O}_{X_I}$ satisfy the relative S_2 property $i_*i^*\mathcal{F} = \mathcal{F}$ where $i\colon X^o_I \subset X_I$ is the inclusion ([H04], Appendix), hence ω_{X_I/S_I} is freely generated by Ω. $\square$

2.4. **The algebra structure.**

In the previous section, we saw that the $\Bbbk[P]/I$-algebra

$$R_I = \Gamma(X^o_{I,\mathfrak{D}}, \mathcal{O}_{X^o_{I,\mathfrak{D}}})$$

defining the flat deformation X_I has a $\Bbbk[P]/I$-module basis of theta functions $\{\vartheta_m \mid m \in B(\mathbb{Z})\}$. Here we derive a description of the multiplication rule on R_I using the geometry of the integral affine manifold B. We will use this (in the case of the canonical scattering diagram $\mathfrak{D}^{\mathrm{can}}$) in §5 to prove our deformation is algebraic when D is not negative semi-definite.

Definition 2.37. For a broken line γ with endpoint $Q \in \tau \in \Sigma$, define $s(\gamma) \in \Lambda_\tau$, $c(\gamma) \in \Bbbk[P]$ by demanding that

$$\mathrm{Mono}(\gamma) = c(\gamma) \cdot z^{\varphi_\tau(s(\gamma))}.$$

Write $\mathrm{Limits}(\gamma) = (q, Q)$ if γ is a broken line for q and has endpoint $Q \in B$.

Theorem 2.38. *Let $q_1, q_2 \in B(\mathbb{Z})$. In the canonical expansion*

$$\vartheta_{q_1} \cdot \vartheta_{q_2} = \sum_{q \in B(\mathbb{Z})} \alpha_q \vartheta_q,$$

where $\alpha_q \in \Bbbk[P]/I$ for each q, we have

$$\alpha_q = \sum_{\substack{(\gamma_1,\gamma_2)\\ \mathrm{Limits}(\gamma_i)=(q_i,z)\\ s(\gamma_1)+s(\gamma_2)=q}} c(\gamma_1)c(\gamma_2)$$

Here $z \in B_0$ is a point very close to q contained in a cell τ, and we identify q with a point of Λ_τ using Remark 2.1.

Proof. To see what the coefficient of ϑ_q is, choose a point $z \in B$ very close to q, and describe the product using the lifts of z^{q_1}, z^{q_2} at z:

$$(\mathrm{Lift}_z(q_1))(\mathrm{Lift}_z(q_2)) = \sum_{q'} \alpha_{q'} \,\mathrm{Lift}_z(q').$$

Now observe first that there is only one broken line γ with endpoint z and $s(\gamma) = q \in \Lambda_\tau$: this is the broken line whose image is $z + \mathbb{R}_{\geq 0} q$. Indeed, the final segment of such a γ is on this ray, and this ray meets no scattering rays, so the broken line cannot bend. Thus the coefficient of $\mathrm{Lift}_z(q)$ on the right-hand side of the above equation can be read off by looking at the coefficient (in $\Bbbk[P]/I$) of $z^{\varphi_\tau(q)}$. This gives the desired description. □

3. The canonical scattering diagram

Here we give the precise definition of $\mathfrak{D}^{\mathrm{can}}$, the scattering diagram suggested by the *regluing* in the symplectic heuristic, §0.6.1. As explained in §0.3.2, it is defined in terms of $\mathbb{A}^1$-classes. We begin by recalling necessary facts about relative Gromov-Witten invariants used to count them.

Definition 3.1. Let $(\tilde{Y}, \tilde{D})$ be a rational surface with $\tilde{D}$ an anti-canonical cycle of rational curves, and let C be an irreducible component of $\tilde{D}$. Consider a class $\beta \in H_2(\tilde{Y}, \mathbb{Z})$ such that

$$\beta \cdot \tilde{D}_i = \begin{cases} k_\beta & \tilde{D}_i = C \\ 0 & \tilde{D}_i \neq C \end{cases} \tag{3.1}$$

for some $k_\beta > 0$. Let F be the closure of $\tilde{D} \setminus C$, and let

$$\tilde{Y}^o := \tilde{Y} \setminus F, \quad C^o := C \setminus F.$$

Let $\overline{\mathfrak{M}}(\tilde{Y}^o/C^o, \beta)$ be the moduli space of stable relative maps of genus zero curves representing the class β with tangency of order k_β at an unspecified point of C^o. (See [Li00], [Li02] for the algebraic definition for these relative Gromov-Witten invariants, and [LR01], [IP03] for the original symplectic definitions.) We refer to β informally as an $\mathbb{A}^1$-*class*. The virtual dimension of this moduli space is

$$-K_{\tilde{Y}} \cdot \beta + (\dim \tilde{Y} - 3) - (k_\beta - 1) = 0.$$

Here the first two terms give the standard dimension formula for the moduli space of stable rational curves in $\tilde{Y}$ representing the class β, and the term $k_\beta - 1$ is the change in dimension given by imposing the k_β-fold tangency condition. The moduli space carries a virtual fundamental class. Furthermore, we have

Lemma 3.2. *$\overline{\mathfrak{M}}(\tilde{Y}^o/C^o, \beta)$ is proper over $\Bbbk$.*

Proof. This follows as in the proof of [GPS09], Theorem 4.2. In brief, let R be a valuation ring with residue field K, with $S = \operatorname{Spec} R$, $T = \operatorname{Spec} K$. We would like to extend a morphism $T \to \overline{\mathfrak{M}}(\tilde{Y}^o/C^o, \beta)$ to S. We know that the moduli space $\overline{\mathfrak{M}}(\tilde{Y}/C, \beta)$ is proper, so we obtain a family of relative stable maps $\mathcal{C} \to S$ to $\tilde{Y}$. We just need to show that in fact the image of the closed fibre $\mathcal{C}_0$ lies in $\tilde{Y}^o$. However, the argument in the proof of [GPS09], Theorem 4.2 shows that if the image of $\mathcal{C}_0$ intersects F, then $\mathcal{C}_0$ must be of genus at least 1, which is not the case. $\square$

Given this, we define

$$N_\beta := \int_{[\overline{\mathfrak{M}}(\tilde{Y}^o/C^o,\beta)]^{vir}} 1.$$

Morally, one should view N_β as counting maps from affine lines to $\tilde{Y} \setminus \tilde{D}$ whose closures represent the class β.

In what follows, we fix as usual the pair (Y, D), $B = B_{(Y,D)}$, and φ the function given by Example 2.3 for some choice of $\eta : \mathrm{NE}(Y) \to P$.

Definition 3.3. Assume P is equipped with a homomorphism

$$\eta : \mathrm{NE}(Y) \to P.$$

Let φ be the multi-valued piecewise linear function on (B, Σ) given by Example 2.3. Fix a ray $\mathfrak{d} \subset B$ with endpoint the origin, with rational slope. If $\mathfrak{d}$ coincides with a ray of Σ, set $\Sigma' := \Sigma$; otherwise, let Σ' be the refinement of Σ obtained by adding the ray $\mathfrak{d}$. This gives a toric blow-up $\pi : \tilde{Y} \to Y$ (the identity in the first case) by Lemma 1.17. Let $C \subset \pi^{-1}(D)$ be the irreducible component corresponding to $\mathfrak{d}$.

Let $\tau_{\mathfrak{d}} \in \Sigma$ be the smallest cone containing $\mathfrak{d}$. Let $m_{\mathfrak{d}} \in \Lambda_{\tau_{\mathfrak{d}}}$ be a primitive generator of the tangent space to $\mathfrak{d}$, pointing away from the origin. Define

$$f_{\mathfrak{d}} := \exp\left[\sum_{\beta} k_\beta N_\beta z^{\eta(\pi_*(\beta)) - \varphi_{\tau_{\mathfrak{d}}}(k_\beta m_{\mathfrak{d}})}\right].$$

Here the sum is over all classes $\beta \in H_2(\tilde{Y}, \mathbb{Z})$ satisfying (3.1). Note that if $N_\beta \neq 0$, then necessarily $\overline{\mathfrak{M}}(\tilde{Y}^o/C^o, \beta)$ is non-empty, and thus $\beta \in \mathrm{NE}(\tilde{Y})$, so $\pi_*(\beta) \in \mathrm{NE}(Y)$. We note that Σ' can be replaced with a further refinement, as $\tilde{Y}^o/C^o$, and so the numbers N_β, does not depend on the particular choice of refinement.

We define

$$\mathfrak{D}^{\mathrm{can}} := \{(\mathfrak{d}, f_{\mathfrak{d}}) \,|\, \mathfrak{d} \subset B \text{ a ray of rational slope}\}.$$

We call a class $\beta \in H_2(\tilde{Y}, \mathbb{Z})$ an $\mathbb{A}^1$-*class* if $N_\beta \neq 0$.

Remark 3.4. In theory, one should be able to use logarithmic Gromov-Witten invariants [GS11] to define N_β without the technical trick of blowing up and working on an open variety. This would be done by working relative to D, and counting rational curves of class β with one point mapping to the boundary with specified orders of tangency exactly as done in §0.4. However, we do not yet know that this way of defining N_β coincides with the method given above, and the definition given above was used in the arguments of [GPS09], on which we rely.

Lemma 3.5. *Let $J \subset P$ be an ideal with $\sqrt{J} = J$. Suppose the map $\eta\colon \mathrm{NE}(Y) \to P$ satisfies the following conditions:*

1. *For any ray $\mathfrak{d} \subset B$ of rational slope, let $\pi : \tilde{Y} \to Y$ be the corresponding blow-up. We require that if $\dim \tau_{\mathfrak{d}} = 2$ or $\dim \tau_{\mathfrak{d}} = 1$ and $p_{\tau_{\mathfrak{d}},\varphi} \notin J$ then for any $\mathbb{A}^1$-class β contributing to $f_{\mathfrak{d}}$, we have $\eta(\pi_*(\beta)) \in J$.*
2. *For any ideal I with $\sqrt{I} = J$, there are only a finite number of $\mathfrak{d}$ and $\mathbb{A}^1$-classes β such that $\eta(\pi_*(\beta)) \notin I$.*

Then $\mathfrak{D}^{\mathrm{can}}$ is a scattering diagram for the data $(B, \Sigma), P, \varphi$, and J.

Proof. Note that

$$z^{\eta(\pi_*(\beta)) - \varphi_{\tau_{\mathfrak{d}}}(k_\beta m_{\mathfrak{d}})} \in I_{\tau_{\mathfrak{d}}, \tau_{\mathfrak{d}}}$$

if and only if $\eta(\pi_*(\beta)) \in I$. So the hypotheses of the lemma imply conditions (2)-(4) of Definition 2.17. □

Example 3.6. Let $\sigma \subset A_1(Y) \otimes_{\mathbb{Z}} \mathbb{R}$ be a strictly convex rational cone containing $\mathrm{NE}(Y)$. (This can be obtained as the dual of a strictly convex rational cone in $\mathrm{Pic}(Y) \otimes_{\mathbb{Z}}$

$\mathbb{R}$ which spans this latter space and is contained in the nef cone.) Let $P = \sigma \cap A_1(Y)$. Since σ is strictly convex, $P^\times = 0$. For any $\mathfrak{m}$-primary ideal I, $P \setminus I$ is a finite set. Let $\eta : \mathrm{NE}(Y) \to P$ be the inclusion. Then the finiteness hypotheses of the above Lemma hold for $J = \mathfrak{m}$ (note that the conditions (3.1) determine $\beta \in A_1(\tilde{Y})$ given $\pi_*(\beta)$).

Example 3.7. We return to the example (Y, D) of a del Pezzo surface together with a cycle of 5 (-1)-curves studied in Example 1.8. Let $P = \mathrm{NE}(Y)$ and η be the identity. Then $\mathfrak{D}^{\mathrm{can}}$ consists of five rays:

$$\mathfrak{D}^{\mathrm{can}} = \{(\rho_i, 1 + z^{[E_i] - \varphi_{\rho_i}(v_i)}) \mid 1 \leq i \leq 5\}.$$

Here E_i is the unique (-1)-curve in Y which is not contained in D and meets D_i transversally, and v_i is the primitive generator of the ray ρ_i corresponding to D_i. To derive this formula from the above definition of the canonical scattering diagram one needs to show that the only possible stable relative maps contributing to $\mathfrak{D}^{\mathrm{can}}$ are multiple covers of the E_i's, and that a k-fold multiple cover contributes a Gromov-Witten invariant of $(-1)^{k-1}/k^2$. It is easier to compute this using the main result of [GPS09], which is done by way of Proposition 3.26. See Example 3.28.

If we accept this description of $\mathfrak{D}^{\mathrm{can}}$, then we can describe all broken lines and the multiplication law given by this diagram.

We first note that no broken line can wrap around $0 \in B$, i.e., if a broken line leaves a cone $\sigma \in \Sigma_{\max}$, it will never return to that cone. It is enough to check this for a straight line (as the bending in any broken line is always *away* from the origin), and this is easily verified, using e.g., Figure 1.1.

Next, since the only scattering rays are the rays $\rho \in \Sigma$, if $q, Q \in \sigma \in \Sigma_{\max}$, then the obvious straight line is the unique broken line for q with endpoint Q. Thus if we describe ϑ_q in the open subset of $X^o_{I,\mathfrak{D}^{\mathrm{can}}}$ corresponding to σ, ϑ_q is just the toric monomial $z^{\varphi_\sigma(q)}$. It follows that

$$\vartheta_{v_i}^a \vartheta_{v_{i+1}}^b = \vartheta_{av_i + bv_{i+1}}$$

for $a, b \geq 0$. In particular, the ϑ_{v_i}'s generate the $\Bbbk[P]/I$-algebra $\Gamma(X_I, \mathcal{O}_{X_I})$, and the algebra structure is determined once we compute $\vartheta_{v_i} \cdot \vartheta_{v_{i+2}}$.

We consider a broken line for v_i. One checks the following, using Figure 1.1 and the above description of $\mathfrak{D}^{\mathrm{can}}$: The broken line can cross at most two rays of Σ, and it bends at most once, at the last ray of Σ that it crosses. See Figure 3.1. From this one deduces using Theorem 2.38:

$$\vartheta_{v_{i-1}} \vartheta_{v_{i+1}} = z^{[D_i]}(\vartheta_{v_i} + z^{[E_i]}). \tag{3.2}$$

The term $z^{[D_i]} \cdot \vartheta_{v_i}$ corresponds to two straight broken lines for v_{i-1}, v_{i+1}, with endpoint the point v_i of ρ_i. The term $z^{[D_i]} \cdot z^{[E_i]}$ is the coefficient of $1 = \vartheta_0$. To compute this we



FIGURE 3.1. The different types of broken lines in Example 3.7.

use the invariance of broken lines, and so choose a generic point Q near 0 and compute the coefficient α_0 of ϑ_0 using pairs γ_i as in Theorem 2.38 whose final directions are opposite, i.e., $s(\gamma_1) + s(\gamma_2) = 0$. If we take $Q \in \sigma_{i,i+1}$, then there is exactly one term contributing to α_0: γ_1 will bend once where it crosses ρ_i, and γ_2 is straight.

One can check that the five equations (3.2) define X_I. These equations are algebraic, and in fact define a flat family over $\operatorname{Spec} \Bbbk[\mathrm{NE}(Y)]$. (This is always the case in the non-negative semi-definite case, see Corollary 5.10). The fibre of this family over the unit in the open torus orbit of the base is obtained by specializing to $z^{[D_i]} = z^{[E_i]} = 1$, we obtain the A_2-cluster algebra, see [G07].

Next we prove that $\mathfrak{D}^{\mathrm{can}}$ is consistent — thus completing our construction of our mirror family X_I for any I with $\sqrt{I} = J$. Consistency is an entirely reasonable expectation — in the symplectic heuristic the regluings from $\mathfrak{D}^{\mathrm{can}}$ are *derived* from the consistency condition. But of course that was a purely heuristic discussion, and our actual proof is based on rather different ideas. As we explain in §0.6.2, the main point is that $\mathfrak{D}^{\mathrm{can}}$ is a limit (under moving worms) of scattering diagrams built from Lemma 3.24 — diagrams about which we know essentially nothing *except* that they satisfy a natural local consistency requirement (the composition of scattering automorphisms associated to a small loop around a smooth point is the identity), and thus are consistent by an argument from [CPS].

Our goal now is to prove the following (the final step in the construction of our mirror family):

Theorem 3.8. *Suppose that we are given a map* $\eta : \mathrm{NE}(Y) \to P$ *such that* φ *is defined as in Example 2.3 by* $p_{\rho,\varphi} = \eta([D_\rho])$. *Suppose furthermore the following conditions hold:*

(I) *For any* $\mathbb{A}^1$*-class* β, $\eta(\pi_*(\beta)) \in J$;

(II) *For any ideal* I *with* $\sqrt{I} = J$, *there are only a finite number of* $\mathbb{A}^1$*-classes* β *such that* $\eta(\pi_*(\beta)) \notin I$.

(III) $\eta([D_\rho]) \in J$ *for at least two boundary components* $D_\rho \subset D$.

Then $\mathfrak{D}^{\mathrm{can}}$ *is a consistent scattering diagram.*

We include here an observation we will need later showing that the canonical scattering diagram only depends on the deformation class of (Y, D).

Lemma 3.9. *Let* $(\mathcal{Y}, \mathcal{D}) \to S$ *be a flat family of pairs over a connected base* S, *with each fibre* $(\mathcal{Y}_s, \mathcal{D}_s)$ *being a non-singular rational surface with anti-canonical cycle. Suppose further that there is a trivialization* $\mathcal{D} \cong D \times S$. *Then for any* $s, s' \in S$, $(\mathcal{Y}_s, \mathcal{D}_s)$ *and* $(\mathcal{Y}_{s'}, \mathcal{D}_{s'})$ *induce the same canonical scattering diagram, after making an identification of* $H_2(\mathcal{Y}_s, \mathbb{Z})$ *with* $H_2(\mathcal{Y}_{s'}, \mathbb{Z})$ *via parallel transport with respect to the Gauss-Manin connection along any path from* s *to* s'.

Proof. It is enough to show that the numbers N_β are deformation invariants in the above sense, i.e., if we are given a family $\pi : (\tilde{\mathcal{Y}}, \tilde{\mathcal{D}}) \to S$ with each fibre as in Definition 3.1, with an irreducible component $\mathcal{C} \subset \tilde{\mathcal{D}}$, then the number

$$N_{\beta,s} := \int_{[\overline{\mathfrak{M}}(\tilde{\mathcal{Y}}^o_s/\mathcal{C}^o_s,\beta)]^{vir}} 1$$

is independent of s. Indeed, once this is shown, then if $N_{\beta,s} \neq 0$, necessarily β defines a class in $\mathrm{NE}(\mathcal{Y}_s)$, as well as in $\mathrm{NE}(\mathcal{Y}_{s'})$, under the chosen identification. This invariance follows from the standard argument that (relative) Gromov-Witten invariants are deformation invariants, with a little care because our target spaces are open. For this, one considers the moduli space $\overline{\mathfrak{M}}(\tilde{\mathcal{Y}}^o/\mathcal{C}^o, \beta)$ of stable maps to $\tilde{\mathcal{Y}}^o$ relative to $\mathcal{C}^o$ and whose composition with π is constant. Then one has a map $\psi : \overline{\mathfrak{M}}(\tilde{\mathcal{Y}}^o/\mathcal{C}^o, \beta) \to S$ whose fibre over s is $\overline{\mathfrak{M}}(\tilde{\mathcal{Y}}^o_s/\mathcal{C}^o_s, \beta)$. Letting ξ be the inclusion of this fibre in $\overline{\mathfrak{M}}(\tilde{\mathcal{Y}}^o/\mathcal{C}^o, \beta)$, deformation invariance will follow if we know that

$$\xi^![\overline{\mathfrak{M}}(\tilde{\mathcal{Y}}^o/\mathcal{C}^o, \beta)]^{vir} = [\overline{\mathfrak{M}}(\tilde{\mathcal{Y}}^o_s/\mathcal{C}^o_s, \beta)]^{vir}$$

and ψ is proper. The first statement is standard in Gromov-Witten theory. The second point, the properness of ψ, follows exactly as in the proof of Lemma 3.2. □

3.1. **Reduction to the Gross–Siebert locus.** We now begin the proof of Theorem 3.8, following the outline given in §0.6.2. We will, however, prove a number of lemmas in a slightly more general context, as we will need some more general consistency results in [K3]. We assume we are given (Y, D), $\eta : \mathrm{NE}(Y) \to P$ and φ defined as in Example 2.3, and a radical ideal $J \subseteq P$. Suppose we are given a scattering diagram $\mathfrak{D}$ for this data; the application in this paper will be $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$. In particular, the hypotheses of Theorem 3.8 imply $\mathfrak{D}^{\mathrm{can}}$ is a scattering diagram for this data.

We first show that we are free to perform toric blowups on (Y, D).

Proposition 3.10. *Let $p : (\tilde{Y}, \tilde{D}) \to (Y, D)$ be a toric blowup. Then if we take $\tilde{\eta} := \eta \circ p_* : \mathrm{NE}(\tilde{Y}) \to P$, then $\mathfrak{D}$ can also be viewed as a scattering diagram for $B_{(\tilde{Y},\tilde{D})}, P$. Furthermore, if $\mathfrak{D}$ is consistent for this latter data, it is consistent for the data $B_{(Y,D)}, P$.*

Proof. Decorate notation, writing for example $\tilde{B}, \tilde{\Sigma}$ for the singular affine manifold with subdivision into cones associated to $(\tilde{Y}, \tilde{D})$. By Lemma 1.17, we have a canonical identification of the underlying singular affine manifolds $B = \tilde{B}$, and $\tilde{\Sigma}$ is the refinement of Σ obtained by adding one ray for each p-exceptional divisor. We have multi-valued piecewise linear functions φ on B and $\tilde{\varphi}$ on $\tilde{B}$. We can in fact choose representatives so that $\tilde{\varphi} = \varphi$. Indeed, $p_{\rho,\tilde{\varphi}} = \eta(p_*([\tilde{D}_\rho]))$ where $\tilde{D}_\rho$ is the irreducible component of $\tilde{D}$ corresponding to ρ. But $p_*([\tilde{D}_\rho]) = 0$ if $\rho \notin \Sigma$, and $p_*([\tilde{D}_\rho]) = [D_\rho]$ if $\rho \in \Sigma$. Thus $\tilde{\varphi}$ in fact has the same domains of linearity as φ, and the same bending parameters, so we can choose representatives which agree.

As a consequence, we note that the sheaves $\mathcal{P}$ and $\tilde{\mathcal{P}}$ on B_0 defined using φ and $\tilde{\varphi}$ coincide. Furthermore, if $\tilde{\rho} \subset \tilde{\sigma}$ are cones in $\tilde{\Sigma}$, with $\rho \in \Sigma$ the smallest cone containing $\tilde{\rho}$ and $\sigma \in \Sigma$ the smallest cone containing $\tilde{\sigma}$, there is a canonical identification of P_{φ_ρ} with $P_{\tilde{\varphi}_{\tilde{\rho}}}$ and a canonical isomorphism

$$\tilde{R}_{\tilde{\rho},\tilde{\sigma},I} \cong R_{\rho,\sigma,I}; \tag{3.3}$$

note the slightly non-trivial case when $\dim \tilde{\rho} = 1$ but $\dim \rho = 2$, in which case we use the fact that $p_{\tilde{\rho},\tilde{\varphi}} = 0$.

Using these identifications, we can view $\mathfrak{D}$ as living on $\tilde{B}$, and as such, one sees from the definition that $\mathfrak{D}$ is a scattering diagram for the data $\tilde{B}, P, \tilde{\varphi}$.

Now suppose $\sqrt{I} = J$. One observes that the set of broken lines contributing to $\mathrm{Lift}_Q(q)$ are the same whether we are working in B or $\tilde{B}$. Thus if $Q \in \tilde{\sigma} \in \tilde{\Sigma}_{\max}$, $\mathrm{Lift}_Q(q) \in \tilde{R}_{\tilde{\sigma},\tilde{\sigma},I}$, defined using $\tilde{B}$, coincides under the isomorphism (3.3) with $\mathrm{Lift}_Q(q) \in R_{\sigma,\sigma,I}$. From this one sees easily that if $\mathfrak{D}$ is consistent for $\tilde{Y}$, it is consistent for Y. ☐

Corollary 3.11. *Given Y, P, η, J satisfying the hypotheses of Theorem 3.8, then Theorem 3.8 holds for this data if it holds for the data $\tilde{Y}, P, \tilde{\eta}, J$.*

Proof. By the proposition, one just needs to check that the canonical scattering diagrams defined using Y or $\tilde{Y}$ are identical. Indeed, given a ray $\mathfrak{d} \subset B$, we can choose a refinement Σ' of Σ which is also a refinement of $\tilde{\Sigma}$, giving maps $\tilde{\pi} : Y' \to \tilde{Y}$ and $\pi : Y' \to Y$. Then for an $\mathbb{A}^1$-class $\beta \in H_2(Y', \mathbb{Z})$, $\eta(\pi_*(\beta)) = \tilde{\eta}(\tilde{\pi}_*(\beta))$, and so $f_{\mathfrak{d}}$ is the same for $\tilde{Y}$ and Y. $\square$

Proposition 3.12. *Suppose we are given monoid homomorphisms*

$$\mathrm{NE}(Y) \xrightarrow{\bar{\eta}} \bar{P} \xrightarrow{\psi} P$$

with $\eta = \psi \circ \bar{\eta}$. *Let* $\overline{\varphi}$ *be the multi-valued piecewise linear function defined using* $\bar{P}, \bar{\eta}$, *so that* $\varphi = \psi \circ \overline{\varphi}$ *is the function defined using* P, η. *Then* ψ *induces maps* $\psi : \bar{P}_{\overline{\varphi}_\tau} \to P_{\varphi_\tau}$. *For any* $f \in \widehat{\mathbb{k}[\bar{P}_{\overline{\varphi}_\tau}]}$, *we can then view* $\psi(f)$ *as a formal sum of monomials in* $\mathbb{k}[P_{\varphi_\tau}]$, *though in general* $\psi(f)$ *need not lie in* $\widehat{\mathbb{k}[P_{\varphi_\tau}]}$. *Suppose* $\bar{\mathfrak{D}}$ *is a scattering diagram for the data* $B, \bar{P}, \bar{\mathfrak{m}} = \bar{P} \setminus \bar{P}^\times$, *such that* $\mathfrak{D} = \psi(\bar{\mathfrak{D}})$ *is a scattering diagram for the data* B, P, J, *where*

$$\psi(\mathfrak{D}) = \{(\mathfrak{d}, \psi(f_{\mathfrak{d}})) \,|\, (\mathfrak{d}, f_{\mathfrak{d}}) \in \mathfrak{D}\}.$$

Thus in particular $\psi(f_{\mathfrak{d}})$ *lies in* $\widehat{\mathbb{k}[P_{\varphi_\tau}]}$ *for each* $\mathfrak{d}$. *If* $\bar{\mathfrak{D}}$ *is consistent for* $\bar{P}, \bar{\eta}, \bar{\mathfrak{m}}$, *then* $\mathfrak{D}$ *is consistent for* P, η, J.

Proof. The monoids P and $\bar{P}$ yield sheaves $\mathcal{P}$ and $\bar{\mathcal{P}}$ over B_0. The map $\psi : \bar{P} \to P$ induces a map of sheaves $\psi : \bar{\mathcal{P}} \to \mathcal{P}$ using $\varphi = \psi \circ \bar{\varphi}$, and hence it also induces monoid homomorphisms $\psi : \bar{P}_{\bar{\varphi}_\tau} \to P_{\varphi_\tau}$.

Let $q \in B_0(\mathbb{Z})$. Then if $\bar{\gamma}$ is a broken line for q with endpoint Q with respect to the barred data, i.e., $\bar{P}$, $\bar{\mathcal{P}}$ etc., we can construct what we shall call $\psi(\bar{\gamma})$. This will be the data required for defining a broken line for the unbarred data, consisting of the map $\psi(\bar{\gamma}) : (-\infty, 0] \to B$ coinciding with $\bar{\gamma}$, and we simply apply ψ to the monomial $m_L(\bar{\gamma})$ attached to a domain of linearity L of $\bar{\gamma}$ to get the attached monomial for $\psi(\bar{\gamma})$. This is not a broken line for the unbarred data, as condition (3) of Definition 2.22 need not hold, precisely because $\psi : \bar{\mathcal{P}} \to \mathcal{P}$ need not be injective.

To rectify this, fix an ideal $I \subset P$ with $\sqrt{I} = J$, $Q \in \sigma \in \Sigma$, and let $\bar{\mathfrak{B}}$ be the set of broken lines $\bar{\gamma}$ for the barred data with endpoint Q such that $\psi(\mathrm{Mono}(\bar{\gamma})) \notin I \cdot \mathbb{k}[P_{\varphi_\sigma}]$. The same finiteness argument of Lemma 2.30 shows that $\bar{\mathfrak{B}}$ is a finite set. We then define an equivalence relation on $\bar{\mathfrak{B}}$ by saying $\bar{\gamma}_1 \sim \bar{\gamma}_2$ provided $\psi(\bar{\gamma}_1)$ and $\psi(\bar{\gamma}_2)$ coincide except possibly for the $\mathbb{k}$-valued coefficients of the monomials attached to the domains of linearity. Given an equivalence class $\xi \subset \bar{\mathfrak{B}}$ with respect to this equivalence relation, we will show there is exactly one broken line γ_ξ for the unbarred data such that

$$\sum_{\bar{\gamma} \in \xi} \psi(\mathrm{Mono}(\bar{\gamma})) = \mathrm{Mono}(\gamma_\xi). \tag{3.4}$$

Furthermore, every broken line γ for the unbarred data with $\mathrm{Mono}(\gamma) \notin I \cdot \Bbbk[P_{\varphi_\sigma}]$ arises in this way.

Define γ_ξ to be the broken line with underlying map given by any element of ξ with the following attached monomials. For any domain of linearity $L = [s, t]$ for γ_ξ, choose a maximal subset $\xi_L \subset \xi$ of broken lines such that $\bar{\gamma}_1|_{(-\infty,t]} \neq \bar{\gamma}_2|_{(-\infty,t]}$ for any $\bar{\gamma}_1, \bar{\gamma}_2 \in \xi_L$. Then define

$$m_L(\gamma_\xi) = \sum_{\bar{\gamma} \in \xi_L} m_L(\psi(\bar{\gamma})).$$

One then checks easily that γ_ξ is a broken line, now satisfying (3) of Definition 2.22, and (3.4) is satisfied since for L the last domain of linearity of γ_ξ, one takes $\xi_L = \xi$. Furthermore, it is easy to see that any broken line for the unbarred data with the same underlying map and attached monomials at most differing by their coefficients from γ_ξ must in fact coincide with γ_ξ. This shows the claim.

Since the broken lines in $\bar{\mathfrak{B}}$ only involve bending at a finite number of places and crossing rays of Σ only a finite number of times, there is some $k > 0$ such that for any $\bar{\gamma} \in \bar{\mathfrak{B}}$, $\mathrm{Mono}(\bar{\gamma}) \in \Bbbk[\bar{P}_{\bar{\varphi}_\tau}]$ does not lie in $\bar{\mathfrak{m}}^k \cdot \Bbbk[\bar{P}_{\bar{\varphi}_\tau}]$. If we then take $\bar{I} = \bar{\mathfrak{m}}^k + \psi^{-1}(I)$, then it is clear from (3.4) that

$$\psi(\overline{\mathrm{Lift}}_Q(q)) = \mathrm{Lift}_Q(q),$$

where $\overline{\mathrm{Lift}}_Q(q)$ is the lift defined with respect to the ideal $\bar{I}$ and the other barred data, and $\mathrm{Lift}_Q(q)$ is defined with respect to the unbarred data and the ideal I. Thus if consistency holds for the barred data, it holds for the unbarred data. $\square$

As a consequence of Proposition 1.19 and Corollary 3.11, in order to prove Theorem 3.8 (i.e., with $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$), we may assume we have a toric model $p : (Y, D) \to (\bar{Y}, \bar{D})$ with $\bar{D} = \bar{D}_1 + \cdots + \bar{D}_n$. Furthermore, by replacing (Y, D) with a deformation equivalent pair and using Lemma 3.9, we can assume that p is the blowup at distinct points x_{ij}, $1 \leq j \leq \ell_i$, along $\bar{D}_i$, with exceptional divisors E_{ij}. Let $\rho_i \in \Sigma$ be the ray corresponding to the proper transform of $\bar{D}_i$.

By Proposition 3.12, we can replace P with a better suited choice of monoid. We shall do this as follows in the case that $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$. As in Example 3.6, the nef cone $\overline{\mathcal{K}}(Y) \subset A^1(Y, \mathbb{R})$ contains a strictly convex rational polyhedral cone σ, so $\sigma^\vee \subset A_1(Y, \mathbb{R})$ is a strictly convex rational polyhedral cone containing $\mathrm{NE}(Y)$. The map $\eta : \mathrm{NE}(Y) \to P$ induces a map $\eta : A_1(Y, \mathbb{R}) \to P^{\mathrm{gp}}_{\mathbb{R}}$. There is some rational polyhedral cone $\sigma_P \subset P^{\mathrm{gp}}_{\mathbb{R}}$ such that $P = \sigma_P \cap P^{\mathrm{gp}}$. In addition, let H be an ample divisor on $\bar{Y}$, so that $\mathrm{NE}(Y) \cap (p^*H)^\perp$ is a face of $\mathrm{NE}(Y)$, generated by the classes $[E_{ij}]$. Now take

$$\sigma_{\bar{P}} = \eta^{-1}(\sigma_P) \cap \sigma^\vee \cap \{q \in A_1(Y, \mathbb{R}) \,|\, p^*H \cdot q \geq 0\},$$

and take

$$\bar{P} = \sigma_{\bar{P}} \cap A_1(Y, \mathbb{Z}).$$

As $\sigma_{\bar{P}}$ is strictly convex, $(\bar{P})^\times = \{0\}$, $\bar{\mathfrak{m}} = \bar{P} \setminus \{0\}$, and if $\bar{I}$ is an $\bar{\mathfrak{m}}$-primary ideal, $\bar{P} \setminus \bar{I}$ is finite. Thus the hypotheses of Theorem 3.8 trivially hold for $\bar{\eta} : \mathrm{NE}(Y) \to \bar{P}$. By Proposition 3.12, we can replace P with $\bar{P}$ to prove Theorem 3.8.

To summarize, we will now make four assumptions in order to prove a given scattering diagram $\mathfrak{D}$ is consistent; the above discussion shows these assumptions can be made in the case of $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$.

Assumptions 3.13.

- *There is a toric model*
$$p : (Y, D) \to (\bar{Y}, \bar{D})$$
which blows up distinct points x_{ij} on D_i, with exceptional divisors E_{ij}.
- $\eta : \mathrm{NE}(Y) \to P$ *is an inclusion, and* $P^\times = \{0\}$.
- *There is a face of P whose intersection with* $\mathrm{NE}(Y)$ *is* $\mathrm{NE}(Y) \cap (p^*H)^\perp$. *Let G be the prime monomial ideal given by the complement of this face.*
- $J = \mathfrak{m} = P \setminus \{0\}$.

We assume we are given a scattering diagram $\mathfrak{D}$ for this data.

Note that $G \neq \mathfrak{m}$ unless p is an isomorphism.

Definition 3.14. The *Gross–Siebert locus* is the open torus orbit T of $\operatorname{Spec} \Bbbk[P]/G$.

The next step will be to extend our family to the formal completion of $\operatorname{Spec} \Bbbk[P]$ along the toric boundary stratum associated to G, and then check the explicit equalities in Theorem 3.8 after restricting to the formal thickening of the Gross–Siebert locus. This reduces us to the two-dimensional situation of [GS07]. In terms of the integral affine manifold, we will pass from $B_{(Y,D)}$ to the smooth manifold $B_{(\bar{Y},\bar{D})} = \mathbb{R}^2$ by first factoring the singularity into a collection of I_1-singularities, one for each point blown-up under p (this yields the manifold B' of §0.6.2) and then pushing these singularities to infinity along their invariant directions. We will be left with a scattering diagram in ordinary affine space, the context of the broken line construction in [G09].

We first analyze an important property of $\mathfrak{D}^{\mathrm{can}}$ in this situation; we will then continue our analysis assuming that $\mathfrak{D}$ shares this property.

For each ray ρ_i in Σ, we have a unique ray $(\rho_i, f_{\rho_i}) \in \mathfrak{D}^{\mathrm{can}}$ with support ρ_i. The following describes f_{ρ_i} mod G.

Lemma 3.15.

$$f_{\rho_i} = g_{\rho_i} \prod_{j=1}^{\ell_i} (1 + b_{ij} X_i^{-1})$$

where $X_i = z^{\varphi_{\rho_i}(v_i)}$ *with* v_i *a primitive generator of* ρ_i*,* $b_{ij} = z^{\eta([E_{ij}])}$ *and* $g_{\rho_i} \equiv 1$ mod G*. The* j^{th} *term of the product is the contribution from* $\mathbb{A}^1$*-classes coming from multiple covers of the* p*-exceptional divisor* E_{ij}*, and* g_{ρ_i} *is the product of contributions from all other* $\mathbb{A}^1$*-classes.*

Proof. Note that in defining f_{ρ_i} using the definition of the canonical scattering diagram, we take $\tilde{Y} = Y$. Now the only terms that contribute to f_{ρ_i} mod G will involve classes $\beta \in \mathrm{NE}(Y) \subset A_1(Y)$ with $\eta(\beta) \notin G$, so in particular, such a β must be a linear combination $\sum_{j=1}^{\ell_i} c_j[E_{ij}]$, with $k_\beta = \sum c_j$. Furthermore, if $f : C \to Y$ contributes to N_β, $f(C)$ must be contained in $\bigcup_{i,j} E_{ij}$. Indeed, if $f(C)$ has an irreducible component D not contained in this set, then $\eta([D]) \in G$, so $\eta(f_*([C])) \in G$, as G is an ideal. But $\eta(f_*([C])) = \eta(\beta)$, which we have assumed is not an element of G.

Since $f(C)$ is connected and intersects D_i, we now see that the image of f is E_{ij} for some j, and in particular, f is a degree k_β cover of E_{ij}. Then Theorem 6.1 of [GPS09] tells us that the contribution from k_β-fold multiple covers of E_{ij} is $(-1)^{k_\beta - 1}/k_\beta^2$. From this we conclude that

$$\begin{aligned} f_{\rho_i} &= \exp\left(h + \sum_{k=1}^{\infty}\sum_{j=1}^{\ell_i} k_\beta \left(\frac{(-1)^{k_\beta-1}}{k_\beta^2}\right)(b_{ij}X_i^{-1})^{k_\beta}\right) \\ &= \exp(h)\prod_{j=1}^{\ell_i}(1 + b_{ij}X_i^{-1}) \end{aligned}$$

where $h \in G$. We take $g_{\rho_i} = \exp(h)$. □

Corollary 3.16. $U_{\rho_i,G}$ *is isomorphic to the hypersurface of Equation* (0.10)*.*

Proof. Since the $g_{\rho_i} \equiv 1$ mod G, they are already units in $R_{\rho_i,\sigma_\pm,G}$ and so do not contribute to the isomorphism type of the fibre product, equation (2.9). Now the result follows from Lemma 2.19. □

Corollary 3.17. $\mathfrak{D}^{\mathrm{can}}$ *is a scattering diagram for the data* (B, Σ)*,* P*,* φ *and* G*.*

Proof. Fixing an $I \subset P$ with $\sqrt{I} = G$, there exists a bound n such that $q \in P \setminus I$ implies $q \cdot p^*H < n$, where H is a fixed ample divisor on $\bar{Y}$. Thus if β is an $\mathbb{A}^1$-class with $\eta(\pi_*(\beta)) \in P \setminus I$, there are only a finite number of choices for $p_*\pi_*\beta$, and for each of these choices, there are only a finite number of choices of $\pi_*\beta$. This shows condition (4) in Definition 2.17 of scattering diagrams, from which also follows condition (2). Note that $p_{\rho_i,\varphi} = [D_i] \in G$ for each i so condition (3) is vacuous for $\dim \tau_{\mathfrak{d}} = 1$. If $\dim \tau_{\mathfrak{d}} = 2$, any contributing $\mathbb{A}^1$-class β satisfies $\pi_*\beta \in G$, so (3) holds. □

Theorem 3.18. *We follow the above notation.*

(1) $\mathfrak{D}^{\mathrm{can}}$ *is consistent as a scattering diagram for* (B,Σ)*,* P*,* φ*, and* G*.*
(2) *Theorem 3.8 follows from (1).*

Proof. We shall for now only show item (2). This just follows from the series of reductions of Theorem 3.8 already made and the observation that if I' is an $\mathfrak{m}$-primary ideal, then since $G\subset\mathfrak{m}$ one can find some k such that $kG\subset I'$. To show consistency holds for the ideal I', we use (1) to observe consistency holds for the ideal $I=kG$, and this gives the desired result. □

Remark 3.19. Given a consistent scattering diagram $\mathfrak{D}$ for (B,Σ), P, φ, and G, Theorem 2.33 shows that with $\sqrt{I}=G$,

$$X_I := \operatorname{Spec}\Gamma(X^o_{I,\mathfrak{D}},\mathcal{O}_{X^o_{I,\mathfrak{D}}})$$

is flat over $\operatorname{Spec}\Bbbk[P]/I$, and $X_G=\mathbb{V}_n\times\operatorname{Spec}\Bbbk[P]/G$.

Let $T\subset\operatorname{Spec}\Bbbk[P]/G$ be the Gross–Siebert locus, Definition 3.14. Note T determines open subschemes of the thickenings $\operatorname{Spec}\Bbbk[P]/I$, which we will shall denote by T_I.

We can describe the subscheme T_I of $\operatorname{Spec}\Bbbk[P]/I$ as follows. Let $E\subset P^{\mathrm{gp}}$ be the lattice generated by the face $P\setminus G$. Then as a subset of $\operatorname{Spec}\Bbbk[P]/G$, $T\cong\operatorname{Spec}\Bbbk[E]$. Furthermore, if we take the localization $P+E$ of P along the face $P\setminus G$, then T_I as a subscheme of $\operatorname{Spec}\Bbbk[P]/I$ is $\operatorname{Spec}\Bbbk[P+E]/(I+E)$.

Note that $\mathfrak{m}_{P+E}=(P+E)\setminus E$, and $G=P\cap\mathfrak{m}_{P+E}$, so we can write $\Bbbk[E]=\Bbbk[P+E]/\mathfrak{m}_{P+E}$.

We can now view φ as a multi-valued strictly $(P+E)$-convex function. Then we have the following obvious

Lemma 3.20. *Suppose* $\mathfrak{D}$ *is a consistent scattering diagram for the data* (B,Σ)*,* $P+E$*,* φ*,* $\mathfrak{m}_{P+E}$*. Then* $\mathfrak{D}$ *is consistent as a scattering diagram for* $(B,\Sigma),P,\varphi$ *and* G*. In particular, Theorem 3.18, (1) holds if* $\mathfrak{D}^{\mathrm{can}}$ *is consistent as a scattering diagram for* $P+E$*,* $\mathfrak{m}_{P+E}$*.*

For $\bar I\subset\mathfrak{m}_{P+E}$ an ideal with $\sqrt{\bar I}=\mathfrak{m}_{P+E}$, and $I=\bar I\cap P$, then $X^o_{I,\mathfrak{D}}$, which is flat over $\operatorname{Spec}\Bbbk[P]/I$, when restricted to the open set $\operatorname{Spec}\Bbbk[P+E]/\bar I$ gives the flat family $X^o_{\bar I,\mathfrak{D}}$.

We now replace P by $P+E$ and J by $\mathfrak{m}_{P+E}$ in what follows.

Now consider the affine manifold $\bar B := B_{(\bar Y,\bar D)}$. By Example 1.2, $\bar B$ in fact has no singularity at the origin, and is affine isomorphic to $M_{\mathbb{R}}=\mathbb{R}^2$ (with $M=\mathbb{Z}^2$), while $\bar\Sigma$ is precisely the fan for $\bar Y$. In order to distinguish between constructions on (Y,D) and $(\bar Y,\bar D)$, we decorate all existing notation with bars. For example, if $\tau\in\Sigma$, denote the corresponding cone of $\bar\Sigma$ by $\bar\tau$. Let $\bar\varphi$ be the multi-valued $P^{\mathrm{gp}}_{\mathbb{R}}$-valued function on

$\bar{B}$ such that

$$p_{\bar{\rho},\bar{\varphi}} = p^*[\bar{D}_{\bar{\rho}}].$$

Note that by Lemma 1.14, we can assume $\bar{\varphi}$ is in fact a single-valued fuction on $M_{\mathbb{R}}$. This single-valuedness will be important to be able to apply the method of Kontsevich and Soibelman, Lemma 3.24. We then have the associated $P^{\mathrm{gp}}_{\mathbb{R}}$-bundle $\overline{\pi} : \overline{\mathbb{P}}_0 \to \bar{B}_0$, with convex section $\overline{\varphi} : \bar{B}_0 \to \overline{\mathbb{P}}_0$, by the construction of §2.1.

We now have sheaves $\bar{\mathcal{P}}$ on $\bar{B}_0$ and $\mathcal{P}$ on B_0, induced by the two functions $\bar{\varphi}$ and φ respectively.

Note that since $\bar{\varphi}$ is single-valued and $\bar{B}$ has no singularities, $\bar{\mathcal{P}}$ is the constant sheaf with fibre $P^{\mathrm{gp}} \oplus M$.

There is a canonical piecewise linear map

$$\nu : B \to \bar{B}$$

which restricts to an integral affine isomorphism $\nu|_\sigma : \sigma \to \bar{\sigma}$, where $\sigma \in \Sigma_{\max}$ and $\bar{\sigma} \in \bar{\Sigma}_{\max}$ is the corresponding cell of $\bar{\Sigma}$. Note this map identifies $B(\mathbb{Z})$ with $\bar{B}(\mathbb{Z})$.

Lemma 3.21. *There is a unique piecewise linear map*

$$\tilde{\nu} : \mathbb{P}_0 \to \bar{\mathbb{P}}_0$$

satisfying

- $\bar{\pi} \circ \tilde{\nu} = \nu \circ \pi$.
- $\tilde{\nu}$ *is equivariant for the* $P^{\mathrm{gp}}_{\mathbb{R}}$*-action.*
- $\tilde{\nu} \circ \varphi = \bar{\varphi} \circ \nu$.

Proof. Existence and uniqueness are clear, since both sides are bundles of P^{gp} torsors. □

For each maximal cone $\sigma \in \Sigma_{\max}$, the derivative ν_* of ν induces a canonical identification of $\Lambda_{B,\sigma}$ with $\Lambda_{\bar{B},\overline{\sigma}}$. This then gives an induced isomorphism of monoids:

$$\tilde{\nu}_\sigma : P_{\varphi_\sigma} \to P_{\bar{\varphi}_{\bar{\sigma}}} \tag{3.5}$$

given by

$$\varphi_\sigma(m) + p \mapsto \bar{\varphi}_{\bar{\sigma}}(\nu_*(m)) + p,$$

for $p \in P$ and $m \in \Lambda_\sigma$. This identifies the $\Bbbk[P]$-algebras $\Bbbk[P_{\varphi_\sigma}]$ and $\Bbbk[P_{\bar{\varphi}_{\bar{\sigma}}}]$, and the completions $\widehat{\Bbbk[P_{\varphi_\sigma}]}$ and $\widehat{\Bbbk[P_{\bar{\varphi}_{\bar{\sigma}}}]}$.

Because the map ν is only piecewise linear around rays $\rho \in \Sigma$, there is only a piecewise linear identification of P_{φ_ρ} with $P_{\bar{\varphi}_{\bar{\rho}}}$ and hence no identification of the corresponding rings. However, ν_* is still defined on the tangent space to ρ, and there is an identification

$$\tilde{\nu}_\rho : \{\varphi_\rho(m) + p \,|\, m \text{ is tangent to } \rho,\ p \in P\} \to \{\bar{\varphi}_{\bar{\rho}}(m) + p \,|\, m \text{ is tangent to } \bar{\rho},\ p \in P\}$$

given by

$$\varphi_\rho(m) + p \mapsto \bar{\varphi}_{\bar{\rho}}(\nu_*(m)) + p.$$

We now explain the Kontsevich-Soibelman lemma. This has to do with scattering diagrams on the smooth affine surface $M_{\mathbb{R}} = \mathbb{R}^2$ (such as $\bar{B} = B_{(\bar{Y},\bar{D})}$). For this general discussion, we fix the data of a monoid Q which comes along with a map $r : Q \to M$. Let $\mathfrak{m}_Q = Q \setminus Q^\times$, and let $\widehat{\Bbbk[Q]}$ denote the completion of $\Bbbk[Q]$ with respect to the monomial ideal $\mathfrak{m}_Q$. (In our application we take $Q = P_{\bar{\varphi}}$ as defined in (1.3).)

We can then consider a variant of the notion of scattering diagram:

Definition 3.22. We define a *scattering diagram for the pair Q, $r : Q \to M$*. This is a set

$$\mathfrak{D} = \{(\mathfrak{d}, f_{\mathfrak{d}})\}$$

where

- $\mathfrak{d} \subset M_{\mathbb{R}}$ is given by
  $$\mathfrak{d} = -\mathbb{R}_{\geq 0} m_0$$
  if $\mathfrak{d}$ is an *outgoing ray* and
  $$\mathfrak{d} = \mathbb{R}_{\geq 0} m_0$$
  if $\mathfrak{d}$ is an *incoming ray*, for some $m_0 \in M \setminus \{0\}$.
- $f_{\mathfrak{d}} \in \widehat{\Bbbk[Q]}$.
- $f_{\mathfrak{d}} \equiv 1 \mod \mathfrak{m}_Q$.
- $f_{\mathfrak{d}} = 1 + \sum_p c_p z^p$ for $c_p \in \Bbbk$, $r(p) \neq 0$ a positive multiple of m_0.
- For any $k > 0$, there are only a finite number of rays $(\mathfrak{d}, f_{\mathfrak{d}}) \in \mathfrak{D}$ with $f_{\mathfrak{d}} \not\equiv 1 \mod \mathfrak{m}_Q^k$.

Definition 3.23. Given a loop γ in $M_{\mathbb{R}}$ around the origin, we define the *path ordered product*

$$\theta_{\gamma,\mathfrak{D}} : \widehat{\Bbbk[Q]} \to \widehat{\Bbbk[Q]}$$

as follows. For each $k > 0$, let $\mathfrak{D}[k] \subset \mathfrak{D}$ be the subset of rays $(\mathfrak{d}, f_{\mathfrak{d}}) \in \mathfrak{D}$ with $f_{\mathfrak{d}} \not\equiv 1 \mod \mathfrak{m}_Q^k$. This set is finite. For $\mathfrak{d} \in \mathfrak{D}[k]$ with $\gamma(t_0) \in \mathfrak{d}$, define

$$\theta^k_{\gamma,\mathfrak{d}} : \Bbbk[Q]/\mathfrak{m}_Q^k \to \Bbbk[Q]/\mathfrak{m}_Q^k$$

by

$$\theta^k_{\gamma,\mathfrak{d}}(z^q) = z^q f_{\mathfrak{d}}^{\langle n_{\mathfrak{d}}, r(q)\rangle}$$

for $n_{\mathfrak{d}} \in M^*$ primitive satisfying, with m a non-zero tangent vector of $\mathfrak{d}$,

$$\langle n_{\mathfrak{d}}, m\rangle = 0, \quad \langle n_{\mathfrak{d}}, \gamma'(t_0)\rangle < 0.$$

Then, if γ crosses the rays $\mathfrak{d}_1, \ldots, \mathfrak{d}_n$ in order with $\mathfrak{D}[k] = \{\mathfrak{d}_1, \ldots, \mathfrak{d}_n\}$, we can define

$$\theta^k_{\gamma,\mathfrak{D}} = \theta^k_{\gamma,\mathfrak{d}_n} \circ \cdots \circ \theta^k_{\gamma,\mathfrak{d}_1}.$$

We then define $\theta_{\gamma,\mathfrak{D}}$ by taking the limit as $k \to \infty$.

The following is a slight generalisation of a result of Kontsevich and Soibelman which appeared in [KS06].

Theorem 3.24. *Let $\mathfrak{D}$ be a scattering diagram in the sense of Definition 3.22. Then there is another scattering diagram* $\mathrm{Scatter}(\mathfrak{D})$ *containing $\mathfrak{D}$ such that* $\mathrm{Scatter}(\mathfrak{D}) \setminus \mathfrak{D}$ *consists only of outgoing rays and $\theta_{\gamma,\mathrm{Scatter}(\mathfrak{D})}$ is the identity.*

For a proof of this theorem essentially as stated here, see [GPS09], Theorem 1.4. The result is unique if $\mathrm{Scatter}(\mathfrak{D}) \setminus \mathfrak{D}$ has at most one ray in each possible direction; we shall assume $\mathrm{Scatter}(\mathfrak{D})$ has been chosen to have this property. This can always be done.

We apply this in the following situation. We take Q to be the monoid $P_{\bar{\varphi}}$ which yields the Mumford degeneration associated to the data $(\bar{B}, \bar{\Sigma})$, $\bar{\varphi}$ (recalling $\bar{B} = M_{\mathbb{R}}$), defined by

$$P_{\bar{\varphi}} = \{(m, \bar{\varphi}(m) + p) \,|\, m \in M, p \in P\} \subset M \times P^{\mathrm{gp}}.$$

This comes with a canonical map $r : P_{\bar{\varphi}} \to M$ by projection.

Definition 3.25. Let $\mathfrak{D}$ be a scattering diagram for the data $(B, \Sigma), P, \varphi, \mathfrak{m}_P$. Suppose furthermore that $\mathfrak{D}$ has at most one outgoing ray in each possible direction, and that for $(\rho_i, f_{\rho_i}) \in \mathfrak{D}$ the unique outgoing ray with support ρ_i,

$$f_{\rho_i} = g_{\rho_i} \prod_{j=1}^{\ell_i} (1 + b_{ij} X_i^{-1})$$

for some $g_{\rho_i} \equiv 1 \mod G$ (so in particular $\mathfrak{D}^{\mathrm{can}}$ satisfies this hypothesis, by Lemma 3.15). We define a scattering diagram $\nu(\mathfrak{D})$ on $\bar{B}$ as follows. For every ray $(\mathfrak{d}, f_{\mathfrak{d}}) \in \mathfrak{D}$ not equal to (ρ_i, f_{ρ_i}) for some i, $\nu(\mathfrak{D})$ contains the ray $(\nu(\mathfrak{d}), \tilde{\nu}_{\tau_{\mathfrak{d}}}(f_{\mathfrak{d}}))$, and for each ray (ρ_i, f_{ρ_i}), $\nu(\mathfrak{D})$ contains two rays, $(\bar{\rho}_i, \tilde{\nu}_{\tau_{\mathfrak{d}}}(g_{\rho_i}))$ and $(\bar{\rho}_i, \prod_{j=1}^{\ell_i}(1 + b_{ij}^{-1} \bar{X}_i))$.

We note that $\nu(\mathfrak{D})$ may not actually be a scattering diagram in the sense of Definition 3.22, as it is possible that $f_{\mathfrak{d}} \notin \widehat{\Bbbk[P_{\bar{\varphi}}]}$: if $p \in P_{\varphi_\tau}$, then $\tilde{\nu}_\tau(p) \in P_{\overline{\varphi}_\tau}$ but need not lie in $P_{\overline{\varphi}}$.

In the case of $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$, we can use the Kontsevich-Soibelman lemma to describe $\nu(\mathfrak{D}^{\mathrm{can}})$. This will both show that $\nu(\mathfrak{D}^{\mathrm{can}})$ is a scattering diagram in the sense of Definition 3.22 and that it satisfies an important additional property which will allow us to prove consistency. Let

$$\bar{\mathfrak{D}}_0 = \{(\bar{\rho}_i, \prod_{j=1}^{\ell_i}(1 + b_{ij}^{-1}\bar{X}_i)) \mid 1 \leq i \leq n\}. \tag{3.6}$$

Let $\mathfrak{m}_{\bar{\varphi}} = P_{\bar{\varphi}} \setminus P_{\bar{\varphi}}^{\times}$ as usual. Then by the strict convexity of $\bar{\varphi}$, $\bar{X}_i \in \mathfrak{m}_{\bar{\varphi}}$ so that $\bar{\mathfrak{D}}_0$ is a scattering diagram for the pair $P_{\bar{\varphi}}$, r in the sense of Definition 3.22. Now define

$$\bar{\mathfrak{D}} := \mathrm{Scatter}(\bar{\mathfrak{D}}_0)$$

where we require $\bar{\mathfrak{D}} \setminus \bar{\mathfrak{D}}_0$ to have only one outgoing ray in each direction (and no incoming rays).

Proposition 3.26. $\bar{\mathfrak{D}} = \nu(\mathfrak{D}^{\mathrm{can}})$. *In particular,* $\nu(\mathfrak{D}^{\mathrm{can}})$ *is a scattering diagram in the sense of Definition 3.22 and* $\theta_{\gamma,\bar{\mathfrak{D}}} \equiv 1$ *for a loop* γ *around the origin.*

Proof. The result follows (after some work) from the main result of [GPS09]. We will explain this derivation in §3.3. The precise statement we need here is Theorem 3.38. □

Remark 3.27. As explained in §0.6.2 the *sign changes* between $\nu(\mathfrak{D}^{\mathrm{can}})$ and $\bar{\mathfrak{D}}$ are explained by the fact that on $\bar{B}$ we have moved the singularities to infinity. If instead they were at finite distance, then the input diagram, call it $\mathfrak{D}_0'$, would have for each singular point an additional outgoing ray, with attached function

$$\prod_{j=1}^{\ell_i}(1 + b_{ij}\bar{X}_i^{-1}).$$

(If we think in terms of $K3$ degenerations as in Remark 1.11, this is the only possible choice of functions that is symmetric with respect to flops and dependent only on the flopping curve.) Now when we take the limit as the singular point moves to the origin, the incoming ray disappears, and this outgoing ray remains.

Example 3.28. Continuing with Example 3.7, note that the pair (Y, D) can be obtained from the toric pair $(\bar{Y}, \bar{D})$ defined by the fan $\bar{\Sigma}$ with rays generated by $(1, 0)$, $(1, 1)$, $(0, 1)$, $(-1, 0)$ and $(0, -1)$, corresponding to $\bar{D}_1, \ldots, \bar{D}_5$, by blowing up one point on each of $\bar{D}_4$ and $\bar{D}_5$. This description determines $\bar{\mathfrak{D}}_0$ and hence $\bar{\mathfrak{D}}$. One can check this description agrees with that given in Example 3.7 for $\mathfrak{D}^{\mathrm{can}}$, see e.g. [GPS09], Example 1.6 for a similar computation.

Now suppose we have a scattering diagram $\mathfrak{D}$ for $(B,\Sigma), P, \varphi, \mathfrak{m}_P$ as in Definition 3.25, and suppose that $\mathfrak{D}$ satisfies the hypotheses of that definition, and that $\nu(\mathfrak{D})$ is a scattering diagram in the sense of Definition 3.22. (For example, by Proposition 3.26, $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$ satisfies these assumptions.) For $I \subset P$ an ideal with $\sqrt{I} = J$, we now have deformations $X^o_{I,\mathfrak{D}}$ and $\bar{X}^o_{I,\nu(\bar{\mathfrak{D}})}$. The latter scheme is glued from open sets

$$\overline{U}_{\bar{\rho},I} = \operatorname{Spec} \overline{R}_{\bar{\rho},I}$$

along open sets identified with $\operatorname{Spec} \overline{R}_{\bar{\sigma},\bar{\sigma},I}$. Here we are decorating the rings coming from the data on $\bar{B}$ with bars as before, while we maintain the notation $R_{\rho,I}$, etc., for those rings coming from the data on B.

Lemma 3.29. *With $\mathfrak{D}$ satisfying the hypotheses of Definition 3.25 and $\nu(\mathfrak{D})$ a scattering diagram in the sense of Definition 3.22, there are isomorphisms*

$$p_i : R_{\rho_i,I} \to \overline{R}_{\bar{\rho}_i,I}$$

and

$$p_{i-1,i} : R_{\sigma_{i-1,i},\sigma_{i-1,i},I} \to \overline{R}_{\bar{\sigma}_{i-1,i},\bar{\sigma}_{i-1,i},I}$$

for all i such that the diagrams

$$\begin{tikzcd} R_{\rho_i,I} \arrow[r] \arrow[d, "p_i"'] & R_{\sigma_{i-1,i},\sigma_{i-1,i},I} \arrow[d, "p_{i-1,i}"] \\ \overline{R}_{\bar{\rho}_i,I} \arrow[r] & \overline{R}_{\bar{\sigma}_{i-1,i},\bar{\sigma}_{i-1,i},I} \end{tikzcd} \qquad \begin{tikzcd} R_{\rho_i,I} \arrow[r] \arrow[d, "p_i"'] & R_{\sigma_{i,i+1},\sigma_{i,i+1},I} \arrow[d, "p_{i,i+1}"] \\ \overline{R}_{\bar{\rho}_i,I} \arrow[r] & \overline{R}_{\bar{\sigma}_{i,i+1},\bar{\sigma}_{i,i+1},I} \end{tikzcd}$$

and

$$\begin{tikzcd} R_{\sigma_{i-1,i},\sigma_{i-1,i},I} \arrow[r, "\theta_{\gamma,\mathfrak{D}}"] \arrow[d, "p_{i-1,i}"'] & R_{\sigma_{i-1,i},\sigma_{i-1,i},I} \arrow[d, "p_{i-1,i}"] \\ \overline{R}_{\bar{\sigma}_{i-1,i},\bar{\sigma}_{i-1,i},I} \arrow[r, "\theta_{\bar{\gamma},\nu(\mathfrak{D})}"'] & \overline{R}_{\bar{\sigma}_{i-1,i},\bar{\sigma}_{i-1,i},I} \end{tikzcd}$$

are commutative.

Consequently, the maps p_i and $p_{i-1,i}$ induce an isomorphism

$$p : X^o_{I,\mathfrak{D}} \to \bar{X}^o_{I,\nu(\mathfrak{D})}$$

over $\operatorname{Spec} \Bbbk[P]/I$.

Proof. To describe p_i, use the representations $R'_{\rho,I}$ and $\overline{R}'_{\bar{\rho},I}$ given by Lemma 2.19, so that

$$R'_{\rho,I} = \frac{(\Bbbk[P]/I)[X_{i-1}, X_i^{\pm}, X_{i+1}]}{(X_{i-1}X_{i+1} - z^{\eta([D_i])}X_i^{-D_i^2} g_{\rho_i} \prod_{j=1}^{\ell_i}(1 + b_{ij}X_i^{-1}))},$$
$$\overline{R}'_{\rho,I} = \frac{(\Bbbk[P]/I)[\bar{X}_{i-1}, \bar{X}_i^{\pm}, \bar{X}_{i+1}]}{(\bar{X}_{i-1}\bar{X}_{i+1} - z^{\eta(p^*[\bar{D}_i])}\bar{X}_i^{-\bar{D}_i^2} \bar{g}_{\rho_i} \prod_{j=1}^{\ell_i}(1 + b_{ij}^{-1}\bar{X}_i))}.$$

We simply define p_i to be the identity on $\Bbbk[P]/I$ and $p_i(X_j) = \bar{X}_j$. This makes sense since $D_i^2 = \bar{D}_i^2 - \ell_i$ and $[D_i] = p^*[\bar{D}_i] - \sum_{j=1}^{\ell_i} E_{ij}$, so that

$$\begin{aligned} p_i\left(z^{\eta([D_i])}X_i^{-D_i^2}\prod_{j=1}^{\ell_i}(1 + b_{ij}X_i^{-1})\right) &= z^{\eta(p^*[\bar{D}_i])}\bar{X}_i^{-\bar{D}_i^2}\left(\prod_{j=1}^{\ell_i} b_{ij}^{-1}\bar{X}_i\right)\left(\prod_{j=1}^{\ell_i}(1 + b_{ij}\bar{X}_i^{-1})\right) \\ &= z^{\eta(p^*[\bar{D}_i])}\bar{X}_i^{-\bar{D}_i^2}\prod_{j=1}^{\ell_i}(1 + b_{ij}^{-1}\bar{X}_i). \end{aligned}$$

The map $p_{i-1,i}$ is induced by $\tilde{\nu}_{\sigma_{i-1,i}}$ defined in (3.5). It is then straightforward to check the commutativity of the three diagrams. □

Lemma 3.30. *Suppose $\mathfrak{D}$ is a scattering diagram satisfying the hypotheses of Definition 3.25 and $\nu(\mathfrak{D})$ a scattering diagram in the sense of Definition 3.22. For $Q \in \sigma_{i-1,i}$, we distinguish between*

$$\mathrm{Lift}_Q(q) \in R_{\sigma_{i-1,i},\sigma_{i-1,i},I}$$

for the lift of $q \in B_0(\mathbb{Z})$ and

$$\mathrm{Lift}_{\nu(Q)}(\nu(q)) \in \bar{R}_{\bar{\sigma}_{i-1,i},\bar{\sigma}_{i-1,i}}$$

the lift of $\nu(q)$. Then

1. $p_{i-1,i}(\mathrm{Lift}_Q(q)) = \mathrm{Lift}_{\nu(Q)}(\nu(q))$.
2. *Under the natural identifications $(P_{\bar{\varphi}_\tau})^{\mathrm{gp}} = (P_{\bar{\varphi}})^{\mathrm{gp}}$, for $\tau \in \bar{\Sigma} \setminus \{0\}$, $P_{\bar{\varphi}} \subset P_{\bar{\varphi}_\tau}$, and for any broken line γ for q, $\mathrm{Mono}(\gamma) \in \Bbbk[P_{\bar{\varphi}}]$.*
3. *ν induces a bijection between broken lines: If $\gamma : (-\infty, 0] \to B_0$ is a broken line in B_0, then $\nu \circ \gamma$ is a broken line in $\bar{B}_0$, and conversely, if $\bar{\gamma} : (-\infty, 0] \to \bar{\mathcal{P}}$ is a broken line in $\bar{B}_0$, then $\nu^{-1} \circ \bar{\gamma}$ is a broken line in B_0.*

Proof. (3) implies (1). For (3), clearly it is enough to compare bending and attached monomials of broken lines near a ray ρ_i.

Consider a broken line γ in B_0 passing from $\sigma_{i-1,i}$ to $\sigma_{i,i+1}$, and let cz^q be the monomial attached to the broken line before it crosses over ρ_i, so that $q \in P_{\varphi_{\sigma_{i-1,i}}}$. Let

θ_{ρ_i}, $\bar{\theta}_{\bar{\rho}_i}$ be defined by

$$\begin{aligned}\theta_{\rho_i}(z^p) &:= z^p f_{\rho_i}^{\langle n, r(p)\rangle}\\ \bar{\theta}_{\bar{\rho}_i}(z^p) &:= z^p \left(\bar{g}_{\rho_i}\prod_{j=1}^{\ell_i}(1+b_{ij}^{-1}\bar{X}_i)\right)^{\langle \bar{n},\bar{r}(p)\rangle}\end{aligned}$$

where $(\bar{\rho}_i, \bar{g}_{\rho_i}) \in \nu(\mathfrak{D})$ is the outgoing ray with support $\bar{\rho}_i$. Then we need to show that

$$p_{i,i+1}(\theta_{\rho_i}(cz^q)) = \bar{\theta}_{\bar{\rho}_i}(p_{i-1,i}(cz^q)) \tag{3.7}$$

to get the correspondence between broken lines.

Note that

$$p_{i-1,i}(X_{i-1}) = \bar{X}_{i-1}, \quad p_{i,i-1}(X_i) = \bar{X}_i, \quad p_{i,i+1}(X_i) = \bar{X}_i,$$

but to compute $p_{i,i+1}(X_{i-1})$, we need to use the relation

$$X_{i-1}X_{i+1} = z^{\eta([D_i])}X_i^{-D_i^2}$$

in $\Bbbk[P_{\varphi_{\rho_i}}]$ to write

$$X_{i-1} = z^{\eta([D_i])}X_i^{-D_i^2}X_{i+1}^{-1}.$$

On the other hand, one has the relation

$$\bar{X}_{i-1}\bar{X}_{i+1} = z^{\eta(p^*[\bar{D}_i])}\bar{X}_i^{-\bar{D}_i^2}$$

in $\Bbbk[P_{\bar{\varphi}_{\bar{\rho}_i}}]$, so

$$\begin{aligned}p_{i,i+1}(X_{i-1}) &= z^{\eta([D_i]-p^*[\bar{D}_i])}\bar{X}_i^{-D_i^2+\bar{D}_i^2}\bar{X}_{i-1}\\ &= \bar{X}_i^{\ell_i}\bar{X}_{i-1}\prod_{j=1}^{\ell_i} b_{ij}^{-1}.\end{aligned}$$

Thus

$$\begin{aligned}p_{i,i+1}(\theta_{\rho_i}(X_{i-1})) &= p_{i,i+1}(X_{i-1}f_{\rho_i})\\ &= \bar{X}_i^{\ell_i}\bar{X}_{i-1}\left(\prod_{j=1}^{\ell_i} b_{ij}^{-1}\right)\left(\prod_{j=1}^{\ell_i}(1+b_{ij}\bar{X}_i^{-1})\right)\bar{g}_{\rho_i}\\ &= \bar{X}_{i-1}\bar{g}_{\rho_i}\prod_{j=1}^{\ell_i}(1+b_{ij}^{-1}\bar{X}_i)\\ &= \bar{\theta}_{\bar{\rho}_i}(p_{i-1,i}(X_{i-1}))\end{aligned}$$

as desired. Also,

$$p_{i,i+1}(\theta_{\rho_i}(X_i)) = \bar{X}_i = \bar{\theta}_{\bar{\rho}_i}(p_{i-1,i}(\bar{X}_i)).$$

Thus (3.7) holds. This shows (3).

For (2), the statement that $P_{\bar\varphi}\subset P_{\bar\varphi_\tau}$ is obvious. For $q\in\sigma\in\Sigma$, by definition the monomial attached to the first domain of linearity of a broken line for q is $z^{\varphi_\sigma(q)}$, which is identified under $\tilde\nu_\sigma$ with $z^{(\nu(q),\bar\varphi(q))}\in\Bbbk[P_{\bar\varphi}]$. For any $(\mathfrak{d},f_{\mathfrak{d}})\in\nu(\mathfrak{D})$, $f_{\mathfrak{d}}\in\widehat{\Bbbk[P_{\bar\varphi}]}$ by assumption, and hence all monomials associated to broken lines in $\bar B_0$ lie in $\Bbbk[P_{\bar\varphi}]$, hence (2). $\square$

Definition 3.31. Suppose $\mathfrak{D}$ is a scattering diagram satisfying the hypotheses of Definition 3.25 and $\nu(\mathfrak{D})$ is a scattering diagram in the sense of Definition 3.22. Let $\mathfrak{m}_{P_{\bar\varphi}}=P_{\bar\varphi}\setminus P_{\bar\varphi}^\times$, and let $I\subset P_{\bar\varphi}$ be an ideal with $\sqrt{I}=\mathfrak{m}_{P_{\bar\varphi}}$. We define for $q\in\bar B_0(\mathbb{Z})$ and $Q\in\bar B_0$,

$$\overline{\mathrm{Lift}}_Q(q)=\sum\mathrm{Mono}(\gamma)\in\Bbbk[P_{\bar\varphi}]/I$$

where the sum is over all broken lines γ for q with endpoint Q in $\bar B_0$ with respect to the scattering diagram $\nu(\mathfrak{D})$. One sees easily as in Lemma 2.30 that this is a finite sum.

The last reduction comes down to the following theorem, to be proved in the next section:

Theorem 3.32. *Assume $\mathfrak{D}$ satisfies the hypotheses of Definition 3.25 and $\nu(\mathfrak{D})$ is a scattering diagram in the sense of Definition 3.22. Suppose furthermore that $\theta_{\gamma,\nu(\mathfrak{D})}\equiv 1$ for a loop γ around the origin. (These hypotheses hold for $\mathfrak{D}=\mathfrak{D}^{\mathrm{can}}$ by Proposition 3.26.) Fix an ideal $I\subset P_{\bar\varphi}$ with $\sqrt{I}=\mathfrak{m}_{P_{\bar\varphi}}$ and $q\in\bar B_0(\mathbb{Z})$. If $Q,Q'\in M_{\mathbb{R}}\setminus\mathrm{Supp}(\nu(\mathfrak{D})_I)$ are general, and γ is a path connecting Q and Q' for which $\theta_{\gamma,\nu(\mathfrak{D})_I}$ is defined, then*

$$\overline{\mathrm{Lift}}_{Q'}(q)=\theta_{\gamma,\nu(\mathfrak{D})_I}(\overline{\mathrm{Lift}}_Q(q))$$

as elements of $\Bbbk[P_{\bar\varphi}]/I$.

Proof of Theorem 3.18, (1), hence of Theorem 3.8. Note that if we want to check an equality

$$\mathrm{Lift}_{Q'}(q)=\theta_{\gamma,\mathfrak{D}^{\mathrm{can}}}(\mathrm{Lift}_Q(q))$$

for $Q,Q'\in\sigma_{i-1,i}$, by Proposition 3.26, Lemma 3.29, and Lemma 3.30, it is sufficient to show that

$$\mathrm{Lift}_{\nu(Q')}(\nu(q))=\theta_{\bar\gamma,\bar{\mathfrak{D}}}(\mathrm{Lift}_{\nu(Q)}(\nu(q))).$$

On the other hand, to check this equality we can compare coefficients of monomials, and given any monomial z^p appearing on the left- or right-hand sides, we can apply Theorem 3.32 with $I=\mathfrak{m}_{P_{\bar\varphi}}^k$ for sufficiently large k so that $p\notin I$. The same strategy applies if Q, Q' lie on opposite sides of a ray ρ. $\square$

3.2. **The proof of Theorem 3.32.** The proof of Theorem 3.32 is just a variation of the argument given in [G09], §4, or rather, a generalization and simplification of this argument due to Carl, Pumperla and Siebert [CPS]. (See also [G11], §5.4.)

Suppose $\mathfrak{D}$ satisfies the hypotheses of Theorem 3.32. Recall that the sheaf $\overline{\mathcal{P}}$ is trivial over $\bar{B} \setminus \{0\}$ so we can extend it trivially over $\bar{B} = M_{\mathbb{R}}$. Recall also that all monomials attached to broken lines for the scattering diagram $\bar{\mathfrak{D}} := \nu(\mathfrak{D})$ in $\bar{B}$ lie in $\Bbbk[P_{\overline{\varphi}}]$ (Lemma 3.30, (2)). We then consider families of broken lines in $M_{\mathbb{R}}$ with respect to the scattering diagram $\bar{\mathfrak{D}}$:

Definition 3.33. A *family of broken lines* consists of the data:

- A continuous map $\Gamma : (-\infty, 0] \times \mathfrak{I} \to M_{\mathbb{R}}$ with $\mathfrak{I} \subset \mathbb{R}$ an interval such that for each $s \in \mathfrak{I}$, $\Gamma_s := \Gamma|_{(-\infty,0]\times\{s\}}$ is a piecewise integral affine map. Furthermore, if L_s is a maximal domain of linearity for Γ_s, there exists a closed set $L \subset (-\infty, 0] \times \mathfrak{I}$ such that $L \cap ((-\infty, 0] \times \{s\}) = L_s$ and $L \cap ((-\infty, 0] \times \{s'\})$ is non-empty and a maximal domain of linearity for $\Gamma_{s'}$ for all $s' \in \mathfrak{I}$.
- For each set $L \subset (-\infty, 0] \times \mathfrak{I}$ as above, a monomial $m_L \in \Bbbk[P_{\overline{\varphi}}]$, such that for each $s' \in \mathfrak{I}$, $\Gamma_{s'}$ along with this data is an ordinary broken line.

We say $\Gamma_{s'}$ is a *deformation* of Γ_s for $s, s' \in \mathfrak{I}$.

The basic idea is that broken lines deform freely as we deform the endpoint, unless, as we deform the endpoint, the image of the broken line passes through the origin of $\bar{B} = M_{\mathbb{R}}$. We then have to show that we can pass through the origin. This may not happen in a continuous fashion, but we will show that the total contribution from such lines remains unchanged.

To carry out this analysis, fixing the ideal I with $\sqrt{I} = \mathfrak{m}_{P_{\overline{\varphi}}}$, let

$$S_I = \{-r(p) \,|\, p \in P_{\bar{\varphi}} \setminus I\}.$$

It is easy to see that this is a finite set. Then set

$$\mathfrak{U}_I := \mathrm{Supp}_I(\bar{\mathfrak{D}}) \cup \bigcup_{m \in S_I} \mathbb{R}_{\geq 0} m.$$

Suppose that we want to deform a broken line γ by deforming its endpoint. We can do this continuously as follows. If L is the unique unbounded domain of linearity, we can translate the unbounded segment $\gamma(L)$ of γ. Inductively, this deforms all the remaining segments of γ. As long as the endpoint of γ is not deformed through a ray along which γ bends or the image of one of the bending points does not reach the origin of $M_{\mathbb{R}}$, each bending point remains inside exactly the same set of rays in $\bar{\mathfrak{D}}$, and therefore the deformed broken line can bend in exactly the same way as γ. If γ contributes to $\overline{\mathrm{Lift}}_Q(q)$, then the monomial cz^p associated to the last line segment of

γ must satisfy $p \in P_{\bar{\varphi}} \setminus I$. Since the image of this last line segment in $M_{\mathbb{R}}$ points in the direction $-r(p)$, we see that we can continue to deform γ freely as long as the endpoint Q doesn't pass through a ray of $\bar{\mathfrak{D}}$ at which γ bends or passes through the ray $-\mathbb{R}_{\geq 0} r(p)$.

From this it is clear that as long as the endpoint of γ stays within one connected component of $M_{\mathbb{R}} \setminus \mathfrak{U}_I$, γ can be deformed continuously. More precisely, if we consider a path $\zeta : [0,1] \to \mathfrak{u}$, for $\mathfrak{u}$ a connected component of $M_{\mathbb{R}} \setminus \mathfrak{U}_I$, and γ is a broken line with endpoint $\zeta(0)$, then there is a continuous deformation Γ with $\gamma = \Gamma_0$ and with $\Gamma_s(0) = \zeta(s)$, $0 \leq s \leq 1$.

From this one sees that $\overline{\mathrm{Lift}}_q(Q)$ is independent of the choice of Q provided Q stays within $\mathfrak{u}$.

We will now analyze carefully how broken lines change if their endpoint passes between different connected components of $M_{\mathbb{R}} \setminus \mathfrak{U}_I$. So now consider two connected components $\mathfrak{u}_1$ and $\mathfrak{u}_2$ of $M_{\mathbb{R}} \setminus \mathfrak{U}_I$. Let $\ell = \overline{\mathfrak{u}}_1 \cap \overline{\mathfrak{u}}_2$, and assume $\dim \ell = 1$. Let Q_1 and Q_2 be general points in $\mathfrak{u}_1$ and $\mathfrak{u}_2$, near ℓ, positioned on opposite sides of ℓ. Let $\zeta : [0,1] \to M_{\mathbb{R}}$ be a short general path connecting Q_1 and Q_2 crossing ℓ precisely once. Let $s_0 \in (0,1)$ be the unique point such that $\zeta(s_0) \in \ell$. We may assume that $\gamma(s_0) \neq 0$.

Let $\mathfrak{B}(Q_i)$ be the set of broken lines with endpoint Q_i. Let $n_0 \in N$ be the primitive vector annihilating the tangent space to ℓ such that $\langle n_0, Q_2 \rangle > 0$. We can decompose $\mathfrak{B}(Q_i)$ into three sets $\mathfrak{B}^+(Q_i)$, $\mathfrak{B}^-(Q_i)$, and $\mathfrak{B}^0(Q_i)$ as follows. For $\gamma \in \mathfrak{B}(Q_i)$, let $m_\gamma \in M_{\mathbb{R}}$ be the projection $r(q_\gamma)$, where $\mathrm{Mono}(\gamma) = c_\gamma z^{q_\gamma} \in \Bbbk[P_{\overline{\varphi}}]$. Then $\gamma \in \mathfrak{B}^+(Q_i), \mathfrak{B}^-(Q_i)$, or $\mathfrak{B}^0(Q_i)$ depending on whether $\langle n_0, m_\gamma \rangle > 0$, $\langle n_0, m_\gamma \rangle < 0$, or $\langle n_0, m_\gamma \rangle = 0$. This gives decompositions

$$\begin{aligned}
\overline{\mathrm{Lift}}_{Q_1}(q) &= \overline{\mathrm{Lift}}^-_{Q_1}(q) + \overline{\mathrm{Lift}}^0_{Q_1}(q) + \overline{\mathrm{Lift}}^+_{Q_1}(q) \\
\overline{\mathrm{Lift}}_{Q_2}(q) &= \overline{\mathrm{Lift}}^-_{Q_2}(q) + \overline{\mathrm{Lift}}^0_{Q_2}(q) + \overline{\mathrm{Lift}}^+_{Q_2}(q)
\end{aligned}$$

We will show

$$\theta_{\zeta,\bar{\mathfrak{D}}}(\overline{\mathrm{Lift}}^-_{Q_1}(q)) = \overline{\mathrm{Lift}}^-_{Q_2}(q), \tag{3.8}$$

$$\theta^{-1}_{\zeta,\bar{\mathfrak{D}}}(\overline{\mathrm{Lift}}^+_{Q_2}(q))) = \overline{\mathrm{Lift}}^+_{Q_1}(q), \tag{3.9}$$

$$\overline{\mathrm{Lift}}^0_{Q_2}(q) = \overline{\mathrm{Lift}}^0_{Q_1}(q). \tag{3.10}$$

From this follows the desired identity

$$\theta_{\zeta,\bar{\mathfrak{D}}}(\overline{\mathrm{Lift}}_{Q_1}(q)) = \overline{\mathrm{Lift}}_{Q_2}(q),$$

as $\theta_{\zeta,\bar{\mathfrak{D}}}$ is necessarily the identity on $\overline{\mathrm{Lift}}^0_{Q_1}(q)$. One then uses this inductively to see that this holds for any path ζ with endpoints in $M_{\mathbb{R}} \setminus \mathfrak{U}_I$ for which $\theta_{\zeta,\bar{\mathfrak{D}}}$ is defined.

Proof of (3.8) *and* (3.9). If γ is a broken line with endpoint Q_1 and L the last domain of linearity of γ (i.e., $0 \in L$), then $\gamma(L) \cap \ell = \emptyset$ if $\langle n_0, m_\gamma \rangle \leq 0$, while $\gamma(L) \cap \ell \neq \emptyset$ if $\langle n_0, m_\gamma \rangle > 0$. (Here we are using Q_1 very close to ℓ.) On the other hand, if γ has endpoint Q_2, then $\gamma(L) \cap \ell = \emptyset$ if $\langle n_0, m_\gamma \rangle \geq 0$ and $\gamma(L) \cap \ell \neq \emptyset$ if $\langle n_0, m_\gamma \rangle < 0$.

To see, say, (3.8), we proceed as follows. Let $\gamma \in \mathfrak{B}^-(Q_1)$. By the previous paragraph, $\gamma(L) \cap \ell = \emptyset$. Write $\theta_{\gamma,\bar{\mathfrak{D}}}(\mathrm{Mono}(\gamma))$ as a sum of monomials $\sum_{i=1}^s d_i z^{q_i'}$ as in Definition 2.22, (3). We can then deform γ continuously along ζ to time s_0. Indeed, γ can only converge to something passing through the origin if $\zeta(s_0) \in -\mathbb{R}_{\geq 0} r(q_\gamma)$ for $\mathrm{Mono}(\gamma) = c_\gamma z^{q_\gamma}$. But this only happens if $\gamma \in \mathfrak{B}^0(Q_1)$.

Let γ' be the deformation of γ with endpoint $\zeta(s_0)$. For $1 \leq i \leq s$, we then get a broken line γ_i' by adding a short line segment to γ' in the direction $-r(q_i')$, with attached monomial $d_i z^{q_i'}$. This new broken line has endpoint in $\mathfrak{u}_2$, and hence can be deformed to a broken line $\gamma_i'' \in \mathfrak{B}^-(Q_2)$. We note that the line may not actually bend at ℓ if $d_i z^{q_i'}$ is the term $\mathrm{Mono}(\gamma)$ appearing in $\theta_{\zeta,\bar{\mathfrak{D}}}(\mathrm{Mono}(\gamma))$.

Conversely, any broken line $\gamma \in \mathfrak{B}^-(Q_2)$ clearly arises in this way.

From this, (3.8) becomes clear. (3.9) is identical. □

Proof of (3.10). We will show that there are partitions $\mathfrak{B}^0(Q_1) = \coprod_{i=1}^s \mathfrak{B}_i^1$ and $\mathfrak{B}^0(Q_2) = \coprod_{i=1}^s \mathfrak{B}_i^2$ such that for each i, the contributions to $\overline{\mathrm{Lift}}_{Q_1}(q)$ and $\overline{\mathrm{Lift}}_{Q_2}(q)$ from $\mathfrak{B}_i^1$ and $\mathfrak{B}_i^2$ are the same.

Let $\gamma_1 \in \mathfrak{B}^0(Q_1)$. If γ_1 deforms continuously to a broken line γ_2 in $\mathfrak{B}^0(Q_2)$ without ever passing through the origin, then γ_1 and γ_2 will each appear in one-element sets in the partition, say $\gamma_1 \in \mathfrak{B}_i^1$, $\gamma_2 \in \mathfrak{B}_i^2$, and clearly both these sets contribute the same term to $\overline{\mathrm{Lift}}_{Q_1}(q)$ and $\overline{\mathrm{Lift}}_{Q_2}(q)$.

We need to partition the remaining elements of $\mathfrak{B}^0(Q_1)$ and $\mathfrak{B}^0(Q_2)$ to show they make the same contribution to $\overline{\mathrm{Lift}}_Q(q)$.

To do so, recall by assumption that

$$\theta_{\eta,\bar{\mathfrak{D}}} = \mathrm{Id} \tag{3.11}$$

for η a loop in $M_\mathbb{R}$ around the origin.

Define a *trajectory* in $M_\mathbb{R}$ to be a pair $(\mathfrak{t}, cz^q)$ where $cz^q \in \Bbbk[P_{\bar{\varphi}}]$ is a monomial and either

1. $\mathfrak{t} = \mathbb{R}_{\geq 0} r(q)$, in which case we say $(\mathfrak{t}, cz^q)$ is an *incoming trajectory*, or
2. $\mathfrak{t} = -\mathbb{R}_{\geq 0} r(q)$, in which case we say $(\mathfrak{t}, cz^q)$ is an *outgoing trajectory.*

As usual, $\mathfrak{t}$ is called the *support* of $(\mathfrak{t}, cz^q)$.

Claim 3.34. *Given an incoming trajectory* $\mathfrak{t}_0 = (\mathbb{R}_{\geq 0} r(q_0), c_0 z^{q_0})$*, there is a set of outgoing trajectories* $\{\mathfrak{t}_i\}$ *with* $\mathfrak{t}_i = (-\mathbb{R}_{\geq 0} r(q_i), c_i z^{q_i})$ *with the following property. For* *each* $\mathfrak{t}_i$ *(including* $i=0$*) let* x_i *be a point in a connected component of* $M_{\mathbb{R}} \setminus \mathrm{Supp}_I(\bar{\mathfrak{D}})$ *whose closure contains the support of* $\mathfrak{t}_i$*. Let* ζ_i *be a path from* x_i *to* x_0 *not passing through* 0*; by* (3.11)*,* $\theta_i := \theta_{\zeta_i,\bar{\mathfrak{D}}}$ *is independent of this choice of path. Then*

$$c_0 z^{q_0} = \sum_i \theta_i(c_i z^{q_i}) \mod I. \tag{3.12}$$

Furthermore, the set of trajectories is unique in the sense that given a ray $\mathbb{R}_{\geq 0} m \subset M_{\mathbb{R}}$*, the sum* $\sum_i c_i z^{m_i}$ *over all* $i \neq 0$ *with the support of* $\mathfrak{t}_i$ *being* $\mathbb{R}_{\geq 0} m$ *is uniquely determined by* $(\mathfrak{t}_0, c_0 z^{q_0})$*.*

Proof. First note that (3.12) does not depend on the precise choice of x_i or x_0: if $\mathfrak{t}_i$ (including the case $i=0$) is contained in a ray of $\bar{\mathfrak{D}}$, then $c_i z^{q_i}$ is invariant under the corresponding automorphism.

We now construct the set $\{\mathfrak{t}_i\}$ by induction. At the kth step, we will find a set $\mathfrak{T}_k$ of outgoing trajectories for which (3.12) holds modulo the ideal $\mathfrak{m}^k_{P_{\bar{\varphi}}}$. For $k=1$, we take

$$\mathfrak{T}_1 = \{(-\mathbb{R}_{\geq 0} r(q_0), c_0 z^{q_0})\},$$

which works since $\theta_i \equiv \mathrm{Id} \mod \mathfrak{m}_{P_{\bar{\varphi}}}$.

Assume now we have constructed $\mathfrak{T}_k = \{\mathfrak{t}_i \,|\, i \in I\}$. Then by the induction hypothesis

$$c_0 z^{q_0} - \sum_i \theta_i(c_i z^{q_i}) = \sum c_j z^{q_j} \mod \mathfrak{m}^{k+1}_{P_{\bar{\varphi}}}$$

with $q_j \in \mathfrak{m}^k_{P_{\bar{\varphi}}}$. Then take

$$\mathfrak{T}_{k+1} = \mathfrak{T}_k \cup \{(-\mathbb{R}_{\geq 0} r(q_j), c_j z^{q_j})\};$$

(3.12) now holds modulo $\mathfrak{m}^{k+1}_{P_{\bar{\varphi}}}$.

Since $\mathfrak{m}^k_{P_{\bar{\varphi}}} \subset I$ for large k, this process terminates. It is also clear that at each step there are no choices to be made, hence the uniqueness. □

The point of this set of outgoing trajectories is that it tells us precisely what terms broken lines can produce. Recall we are trying to understand $\overline{\mathrm{Lift}}_Q(q)$. To do this, pick a general point $x \in M_{\mathbb{R}}$ contained in a connected component of $M_{\mathbb{R}} \setminus \mathrm{Supp}_I(\bar{\mathfrak{D}})$ whose closure contains $\mathbb{R}_{\geq 0} q$. Consider a slight modification of the notion of broken line which are maps $\gamma : \mathbb{R} \to M_{\mathbb{R}}$ which are piecewise integral affine together with attached monomials $c_L z^{q_L}$ as usual, and with $\gamma(0) = x$, otherwise satisfying the conditions of Definition 2.22 except for the requirement that $\gamma(0) = Q$. The only difference is that these broken lines have no endpoints and are constrained to pass through x at time 0 instead.

Let $\mathfrak{B}$ be the set of such modified broken lines γ such that the monomial associated with the first segment is $z^{(q,\overline{\varphi}(q))} \in \Bbbk[P_{\bar{\varphi}}]$, and $\gamma|_{(-\infty,0]}$ is linear. For $\gamma_i \in \mathfrak{B}$, let $c_i z^{q_i}$ be the monomial attached to the last line segment of γ_i, and let ζ_i be a path from a point $x_i = \gamma_i(t)$ for $t \gg 0$ to x. Let $\theta_i = \theta_{\zeta_i,\bar{\mathfrak{D}}}$.

Claim 3.35. $\sum_i \theta_i(c_i z^{q_i}) = z^{q_0}$.

Proof. Note that if $\bar{\mathfrak{D}}$ just consisted of one line, then the broken lines in $\mathfrak{B}$ would either not bend or bend at one point, the intersection of $x - \mathbb{R}_{\geq 0} q$ with the unique element of $\bar{\mathfrak{D}}$. It then follows immediately from condition (3) of Definition 2.22 that the claim is true in this case.

Using this, the general case then follows easily if we take, for the path ζ_i, the segment of γ_i running from x_i backwards to x. □

As a consequence, by the uniqueness statement of Claim 3.34, if we fix an outgoing direction $-\mathbb{R}_{\geq 0} m$, the sum $\sum_i c_i z^{q_i}$ where i runs over all indices such that $-r(q_i) \in -\mathbb{R}_{\geq 0} m$ is well-defined, irrespective of the choice of the point x.

We can now complete the proof in this case. Take the set $\mathfrak{B}^1 \subset \mathfrak{B}^0(Q_1)$ to be the subset of all broken lines in $\mathfrak{B}^0(Q_1)$ which do not deform continuously to broken lines with endpoint Q_2 (because they degenerate to broken lines passing through the origin). We define $\mathfrak{B}^2$ similarly to be the set of all broken lines in $\mathfrak{B}^0(Q_2)$ which do not deform continuously to broken lines with endpoint Q_1. We will be done once we show these sets yield the same contribution to $\overline{\mathrm{Lift}}(q)$.

Recalling that we are deforming the endpoints of the broken lines in $\mathfrak{B}^0(Q_1)$ and $\mathfrak{B}^0(Q_2)$ along a path ζ, we define for $s \neq s_0$ the set $\mathfrak{B}_s$ of broken lines with endpoint $\zeta(s)$ which again fail to deform through time s_0, so that each broken line in $\mathfrak{B}_s$ for $s < s_0$ is a deformation of a unique broken line in $\mathfrak{B}^1$ and each broken line in $\mathfrak{B}_s$ for $s > s_0$ is a deformation of a unique broken line in $\mathfrak{B}^2$.

Let $x_0 \in \mathbb{R}_{\geq 0} r(q)$. Locally near x_0, $\mathbb{R}_{\geq 0} r(q)$ splits $M_{\mathbb{R}}$ into two connected components, and broken lines in $\mathfrak{B}_s$ for $s < s_0$ are locally contained in one of these connected components, and broken lines in $\mathfrak{B}_s$ for $s > s_0$ are locally contained in the other. Fix points x, x' on either side of $\mathbb{R}_{\geq 0} r(q)$ near x_0, chosen so that some element of $\mathfrak{B}_s$ for some $s < s_0$ passes through, say, x. For each broken line $\gamma \in \mathfrak{B}^1$, one can find at least one $s < s_0$ such that γ can be deformed to a $\gamma_s \in \mathfrak{B}_s$ which passes through x. Similarly, for each broken line $\gamma \in \mathfrak{B}^2$, one can find at least one $s > s_0$ such that γ can be deformed to a $\gamma_s \in \mathfrak{B}_s$ which passes through x'. For each $\gamma \in \mathfrak{B}^1$, we make a choice of one of these γ_s's passing through x, and let $\mathfrak{B}_x$ be the set of these choices. We define $\mathfrak{B}_{x'}$ similarly, so that $\mathfrak{B}_x$ is in one-to-one correspondence with $\mathfrak{B}^1$ and $\mathfrak{B}_{x'}$ is in one-to-one correspondence with $\mathfrak{B}^2$. In particular, if $c_\gamma z^{q_\gamma}$ denotes the monomial attached to the last segment of a broken line γ, we just need to show that

$$\sum_{\gamma\in\mathfrak{B}_x} c_\gamma z^{q_\gamma} = \sum_{\gamma\in\mathfrak{B}_{x'}} c_\gamma z^{q_\gamma}.$$

But this follows from Claim 3.35 and the uniqueness statement of Claim 3.34.

3.3. **The proof of Proposition 3.26: The connection with** [GPS09]**.** Here we derive Proposition 3.26 from the main result of [GPS09]. We will need to review one form of this result, which gives an enumerative interpretation for the output of the Kontsevich-Soibelman lemma.

Fix $M \cong \mathbb{Z}^2$ as usual. Suppose we are given positive integers $\ell_1, \ldots, \ell_n$ and primitive vectors $m_1, \ldots, m_n \in M$. Let $\ell = \sum_{i=1}^n \ell_i$ and $Q = M \oplus \mathbb{N}^\ell$, with $r : Q \to M$ the projection. Denote the variables in $\Bbbk[Q]$ corresponding to the generators of $\mathbb{N}^\ell$ as t_{ij}, for $1 \le i \le n$ and $1 \le j \le \ell_i$. Consider the scattering diagram for the data $r : Q \to M$ (in the sense of Definition 3.22)

$$\mathfrak{D} = \{(\mathbb{R}_{\ge 0} m_i, \prod_{j=1}^{\ell_i}(1 + t_{ij} z^{m_i}))\,|\, 1 \le i \le n\}.$$

We wish to interpret $(\mathfrak{d}, f_{\mathfrak{d}}) \in \mathrm{Scatter}(\mathfrak{D}) \setminus \mathfrak{D}$. Choose a complete fan $\Sigma_{\mathfrak{d}}$ in $M_{\mathbb{R}}$ which contains the rays $\mathbb{R}_{\ge 0} m_1, \ldots, \mathbb{R}_{\ge 0} m_n$ as well as the ray $\mathfrak{d}$ (which may coincide with one of the other rays). Let $X_{\mathfrak{d}}$ be the corresponding toric surface, and let $D_1, \ldots, D_n, D_{\mathrm{out}}$ be the divisors corresponding to the above rays. Choose general points $x_{i1}, \ldots, x_{i\ell_i} \in D_i$, and let

$$\nu : \tilde{X}_{\mathfrak{d}} \to X_{\mathfrak{d}}$$

be the blow-up of all the points $\{x_{ij}\}$. Let $\tilde{D}_1, \ldots, \tilde{D}_n, \tilde{D}_{\mathrm{out}}$ be the proper transforms of the divisors $D_1, \ldots, D_n, D_{\mathrm{out}}$ and E_{ij} the exceptional curve over x_{ij}.

Now introduce the additional data of $\mathbf{P} = (\mathbf{P}_1, \ldots, \mathbf{P}_n)$, where $\mathbf{P}_i$ denotes a sequence $p_{i1}, \ldots, p_{i\ell_i}$ of ℓ_i non-negative numbers. We will use the notation $\mathbf{P}_i = p_{i1} + \cdots + p_{i\ell_i}$ and call $\mathbf{P}_i$ an *ordered partition.* Define

$$|\mathbf{P}_i| = \sum_{j=1}^{\ell_i} p_{ij}.$$

We shall restrict attention to those $\mathbf{P}$ such that

$$-\sum_{i=1}^n |\mathbf{P}_i| m_i = k_{\mathbf{P}} m_{\mathfrak{d}} \tag{3.13}$$

where $m_{\mathfrak{d}} \in M$ is a primitive generator of $\mathfrak{d}$ and $k_{\mathbf{P}}$ is a positive integer.

Given this data, consider the class $\beta \in H_2(X_{\mathfrak{d}}, \mathbb{Z})$ specified by the requirement that, if D is a toric divisor of $X_{\mathfrak{d}}$ with $D \notin \{D_1, \ldots, D_n, D_{\text{out}}\}$, then $D \cdot \beta = 0$; if $D_{\text{out}} \notin \{D_1, \ldots, D_n\}$,

$$D_i \cdot \beta = |\mathbf{P}_i|, \qquad D_{\text{out}} \cdot \beta = k_{\mathbf{P}};$$

while if $D_{\text{out}} = D_j$ for some j, then

$$D_i \cdot \beta = \begin{cases} |\mathbf{P}_i| & i \neq j, \\ |\mathbf{P}_i| + k_{\mathbf{P}} & i = j. \end{cases}$$

That such a class exists follows easily from (3.13) and the first part of Lemma 3.37 below. It is also unique. We can then define

$$\beta_{\mathbf{P}} = \nu^*(\beta) - \sum_{i=1}^{p} \sum_{j=1}^{\ell_i} p_{ij}[E_{ij}] \in H_2(\tilde{X}_{\mathfrak{d}}, \mathbb{Z}).$$

We define $N_{\mathbf{P}} := N_{\beta_{\mathbf{P}}}$ as in Definition 3.1, using $(\tilde{Y}, \tilde{D}) = (\tilde{X}_{\mathfrak{d}}, \tilde{D})$, where $\tilde{D}$ is the proper transform of the toric boundary of $X_{\mathfrak{d}}$, and using $C = \tilde{D}_{\text{out}}$. Then one of the main theorems of [GPS09] states

Theorem 3.36.

$$\log f_{\mathfrak{d}} = \sum_{\mathbf{P}} k_{\mathbf{P}} N_{\mathbf{P}} t^{\mathbf{P}} z^{-k_{\mathbf{P}} m_{\mathfrak{d}}}, \tag{3.14}$$

where the sum is over all $\mathbf{P}$ *satisfying* (3.13) *and* $t^{\mathbf{P}}$ *denotes the monomial* $\prod_{ij} t_{ij}^{p_{ij}}$.

We can adapt this theorem for our purposes as follows. Fix a fan Σ in $M_{\mathbb{R}}$ defining a complete non-singular toric surface $\bar{Y}$, with $\bar{D} = D_1 + \cdots + D_n$ the toric boundary. Choose points $x_{i1}, \ldots, x_{i\ell_i} \in D_i$, and define a new surface Y as the blow-up $\nu : Y \to \bar{Y}$ at the points $\{x_{ij}\}$. Let E_{ij} be the exceptional curve over x_{ij}.

Let $P = \mathrm{NE}(\bar{Y})$; because $\bar{Y}$ is toric, this is a finitely generated monoid with $P^\times = \{0\}$. Let $\bar{\varphi} : M_{\mathbb{R}} \to P^{\mathrm{gp}}_{\mathbb{R}}$ be the Σ-piecewise linear strictly P-convex function given by Lemma 1.14.

We will need the following lemma:

Lemma 3.37. *Let* T_Σ *be the free abelian group generated by the rays of* Σ, *with generator* t_ρ *corresponding to a ray* $\rho \in \Sigma$. *Let*

$$s : T_\Sigma \to M$$

be the map defined by $s(t_\rho) = m_\rho$, *for* m_ρ *a primitive generator of* ρ. *Then*

$$H_2(\bar{Y}, \mathbb{Z}) \cong \ker s.$$

Furthermore, if $\sum_\rho a_\rho t_\rho \in \ker s$*, then the corresponding element of* $H_2(\bar{Y}, \mathbb{Z})$ *is*

$$\sum_\rho a_\rho \bar{\varphi}(m_\rho) \in P^{\mathrm{gp}}.$$

Proof. The isomorphism is standard, given by

$$H_2(\bar{Y}, \mathbb{Z}) \ni \beta \mapsto \sum_\rho (D_\rho \cdot \beta) t_\rho.$$

To see the second statement, first note that the claim is not affected by adding a linear function to $\bar{\varphi}$, since $\sum_\rho a_\rho m_\rho = 0$ by assumption. Letting $p = \sum_\rho a_\rho \bar{\varphi}(m_\rho)$, we need to show that $p \cdot D_\rho = a_\rho$ for each ρ.

Label the rays of Σ cyclically as $\rho_1, \ldots, \rho_n$, with $\sigma_{i,i+1}$ containing ρ_i, ρ_{i+1} as usual. We can modify $\bar{\varphi}$ by adding a linear function so that $\bar{\varphi}|_{\sigma_{n,1}} = 0$. Once we've done this, then with m_i the primitive generator of ρ_i and $n_i \in N$ primitive, annihilating m_i, with $\langle n_i, m_{i+1} \rangle > 0$, we have

$$\bar{\varphi}|_{\sigma_{i,i+1}} = \sum_{k=1}^{i} n_k \otimes p_{\rho_k, \bar{\varphi}}.$$

Note that $\langle n_k, m_i \rangle = -\langle n_i, m_k \rangle$ and $p_{\rho_k, \bar{\varphi}} = [D_k]$, so

$$\bar{\varphi}(m_i) = -\sum_{k=1}^{i} \langle n_i, m_k \rangle [D_k] = -\sum_{k=1}^{i-1} \langle n_i, m_k \rangle [D_k].$$

Thus on the surface $\bar{Y} \setminus \bigcup_{k=i+1}^{n} D_k$, $\bar{\varphi}(m_i)$ is in fact the divisor of zeroes and poles of z^{-n_i}, so $D_k \cdot \bar{\varphi}(m_i) = 0$ for $2 \le k \le i-1$. Also, clearly $D_k \cdot \bar{\varphi}(m_i) = 0$ for $i+1 \le k \le n-1$. Next, note that if $i < n$, (recalling that $\bar{Y}$ is non-singular)

$$D_i \cdot \bar{\varphi}(m_i) = -\langle n_i, m_{i-1} \rangle = 1$$

and

$$\begin{aligned} D_n \cdot \bar{\varphi}(m_i) &= -\langle n_i, m_1 \rangle - \delta_{i,n} \langle n_i, m_{n-1} \rangle \\ D_1 \cdot \bar{\varphi}(m_i) &= \begin{cases} 0 & i = 1 \\ -\langle n_i, m_1 \rangle D_1^2 - \langle n_i, m_2 \rangle & 2 \le i \le n \end{cases} \end{aligned}$$

Thus we see that for $i \ne 1, n$,

$$D_i \cdot p = a_{\rho_i},$$

as desired, and

$$\begin{aligned}D_n\cdot p &= -\sum_{i=1}^{n}\langle a_{\rho_i}n_i, m_1\rangle + a_{\rho_n} = a_{\rho_n},\\ D_1\cdot p &= -\sum_{i=1}^{n}\langle a_{\rho_i}n_i, m_1\rangle D_1^2 - \sum_{i=2}^{n}\langle a_{\rho_i}n_i, m_2\rangle\\ &= a_{\rho_1}\langle n_1, m_2\rangle = a_{\rho_1}\end{aligned}$$

since $\sum a_{\rho_i}n_i = 0$. □

Let $E \subset H_2(Y,\mathbb{Z})$ be the lattice spanned by the classes of the exceptional curves of ν, so that $H_2(Y,\mathbb{Z}) = \nu^* H_2(\bar{Y},\mathbb{Z}) \oplus E$. We then obtain a map

$$\varphi = \nu^* \circ \bar{\varphi} : M_{\mathbb{R}} \to \nu^* P^{\mathrm{gp}} \oplus E.$$

Let

$$Q = \{(m,p) \in M \oplus H_2(Y,\mathbb{Z}) \,|\, \exists p' \in \nu^* P \oplus E \text{ such that } p = p' + \varphi(m)\}.$$

There is an obvious projection $r : Q \to M$, and by strict convexity of $\bar{\varphi}$, $Q^\times = E$.

We consider the scattering diagram, $\bar{\mathfrak{D}}_0$, over $\widehat{\Bbbk[Q]}$ given by

$$\bar{\mathfrak{D}}_0 = \{(\mathbb{R}_{\geq 0} m_i, \prod_{j=1}^{\ell_i}(1 + z^{(m_i,\varphi(m_i)-E_{ij})})) \,|\, 1 \leq i \leq n\}.$$

Then we have

Theorem 3.38. *Let* $(\mathfrak{d}, f_{\mathfrak{d}}) \in \mathrm{Scatter}(\bar{\mathfrak{D}}_0) \setminus \bar{\mathfrak{D}}_0$, *assuming that there is at most one ray of* $\mathrm{Scatter}(\bar{\mathfrak{D}}_0) \setminus \bar{\mathfrak{D}}_0$ *in each possible outgoing direction. (Note by definition of* $\mathrm{Scatter}(\bar{\mathfrak{D}}_0)$, $(\mathfrak{d}, f_{\mathfrak{d}})$ *cannot be incoming.) Then, following the notation of Definition 3.1 and 3.3,*

$$\log f_{\mathfrak{d}} = \sum_{\beta} k_\beta N_\beta z^{(-k_\beta m_{\mathfrak{d}}, \pi_*(\beta) - \varphi(k_\beta m_{\mathfrak{d}}))}. \tag{3.15}$$

Here $\pi : \tilde{Y} \to Y$ *is the toric blow-up of* Y *determined by* $\mathfrak{d}$ *and* $C \subset \tilde{Y}$ *is the component of the boundary determined by* $\mathfrak{d}$. *If* $\mathfrak{d}$ *is not one of the rays* $\mathbb{R}_{\geq 0} m_i$, *then we sum over all* $\mathbb{A}^1$*-classes* $\beta \in H_2(\tilde{Y},\mathbb{Z})$ *satisfying* (3.1), *and if* $\mathfrak{d} = \mathbb{R}_{\geq 0} m_i$ *we sum over all such classes* **except for** *classes given by multiple covers of one of the exceptional divisors* E_{ij}.

Proof. Let Q' be the submonoid of $M \oplus \mathbb{N}^\ell$ generated by elements of the form (m_i, d_{ij}), where d_{ij} is the (i,j)-th generator of $\mathbb{N}^\ell$. Note that Q' itself is freely generated by these elements. Thus we can define a map

$$\alpha : Q' \to Q$$

by $(m_i, d_{ij}) \mapsto (m_i, \varphi(m_i) - E_{ij})$. The scattering diagram

$$\mathfrak{D}' := \{(\mathbb{R}_{\geq 0} m_i, \prod_{j=1}^{\ell_i} (1 + z^{(m_i, d_{ij})})) \,|\, 1 \leq i \leq n\}$$

then has image under the map α (applying α to each $f_{\mathfrak{d}}$) the scattering diagram $\bar{\mathfrak{D}}_0$. Thus if we apply α to each element of $\mathrm{Scatter}(\mathfrak{D}')$, we must get $\mathrm{Scatter}(\bar{\mathfrak{D}}_0)$, as $\theta_{\gamma, \mathrm{Scatter}(\mathfrak{D}')}$ being the identity on $\widehat{\Bbbk[Q']}$ implies that $\theta_{\gamma, \alpha(\mathrm{Scatter}(\mathfrak{D}'))}$ is the identity on $\widehat{\Bbbk[Q]}$.

To obtain the result, we now note that the set of possible $\mathbb{A}^1$ classes in $\tilde{Y}$ occurring in the expression (3.15) are precisely the classes $\{\beta_{\mathbf{P}}\}$ where $\mathbf{P}$ runs over all partitions satisfying (3.13). Now applying α to a term appearing in (3.14) of the form

$$k_{\mathbf{P}} N_{\mathbf{P}} t^{\mathbf{P}} z^{-k_{\mathbf{P}} m_{\mathfrak{d}}} = k_{\beta_{\mathbf{P}}} N_{\beta_{\mathbf{P}}} \left(\prod t_{ij}^{p_{ij}} \right) z^{\sum_{i=1}^{p} |\mathbf{P}_i| m_i},$$

we get

$$k_{\beta_{\mathbf{P}}} N_{\beta_{\mathbf{P}}} z^{(-k_{\mathbf{P}} m_{\mathfrak{d}}, \sum_{i=1}^{p} |\mathbf{P}_i| \varphi(m_i) - \sum_{i,j} p_{ij} E_{ij})}.$$

But by Lemma 3.37 and (3.13),

$$\sum_{i=1}^{p} |\mathbf{P}_i| \varphi(m_i) - \sum_{i,j} p_{ij} E_{ij} = \pi_*(\beta_{\mathbf{P}}) - \varphi(k_{\mathbf{P}} m_{\mathfrak{d}}),$$

hence the result. □

A direct comparison of the formula of the above theorem and the formula in the definition of the canonical scattering diagram then yields Proposition 3.26.

3.4. **Smoothness: Around the Gross–Siebert locus.** Next we prove that our deformation of $\mathbb{V}_n$ is indeed a smoothing. As explained in §0.6.2 we do this by working over the Gross–Siebert locus (Definition 3.14). Here our deformation (when restricted to one-parameter subgroups associated to p^*A, A an ample divisor on $\bar{Y}$) agrees with the construction of [GS07]. This is important here because the deformations of [GS07] come with explicit charts that cover all of $\mathbb{V}_n$, from which it is clear that they give a smoothing. So conceptually, the smoothing claim is clear. Because we work with formal families the actual argument is a bit more delicate. First we make rigorous the notion of a *smooth generic fibre* for a formal family:

Definition-Lemma 3.39. Let $f : Z \to W$ be a flat finite type morphism of schemes or a flat morphism of complex analytic spaces, of relative dimension d. Then $\mathrm{Sing}(f) \subset Z$ is the closed embedding defined by the d^{th} Fitting ideal of $\Omega^1_{Z/W}$. $\mathrm{Sing}(f)$ is empty if and only if f is smooth. Formation of $\mathrm{Sing}(f)$ commutes with all base extensions of W.

Proof. For the definition of the Fitting ideal, see e.g., [E95], 20.4. The fact that it commutes with base-change follows from the fact that $\Omega^1_{Z/W}$ commutes with base-change and [E95], Cor. 20.5. That $\mathrm{Sing}(f)$ is empty if and only if f is smooth follows from [E95], Prop. 20.6 and the definition of smoothness. $\square$

Now for a formal family, *smoothness of the generic fibre* is measured by the fact that $\mathrm{Sing}(f)$ does not surject scheme-theoretically onto the base. More precisely, let S be a normal variety or complex analytic space, $V \subset S$ a connected closed subset, and $\mathfrak{S}$ the formal completion of S along V. Let $\mathfrak{f}\colon \mathfrak{X} \to \mathfrak{S}$ be an adic flat morphism of formal schemes or formal complex analytic spaces, and $\mathfrak{Z} \subset \mathfrak{X}$ the scheme theoretic singular locus of $\mathfrak{f}$. Then we say the *generic fiber of* $\mathfrak{f}$ *is smooth* if the map $\mathcal{O}_{\mathfrak{S}} \to \mathfrak{f}_*\mathcal{O}_{\mathfrak{Z}}$ is not injective. See Theorem 4.12.

We now continue with our usual setting of a surface (Y, D), and assume we have a toric model $p : (Y, D) \to (\bar{Y}, \bar{D})$, the blowup at distinct points along the boundary, and an inclusion $\eta : \mathrm{NE}(Y) \to P$ satisfying Assumptions 3.13, and let I be a monomial ideal with radical G. Let $\mathfrak{D}$ be a scattering diagram for this data such that $\mathfrak{D}$ satisfies the hypotheses of Definition 3.25, $\nu(\mathfrak{D})$ is a scattering diagram in the sense of Definition 3.22, and $\theta_{\gamma,\nu(\mathfrak{D})} \equiv 1$ for a loop γ around the origin. Thus by Theorem 3.32, $\mathfrak{D}$ is consistent. These hypotheses apply in particular when $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$.

Let T be the Gross–Siebert locus: using the notation of Remark 3.19, we have $T = \mathrm{Spec}\,\Bbbk[P + E]/\mathfrak{m}_{P+E}$. Consistency of $\mathfrak{D}$ gives a flat family

$$f_I : X_I \to \mathrm{Spec}\,\Bbbk[P]/I,$$

which restricts to a family

$$f_{\bar{I}} : X_{\bar{I}} \to \mathrm{Spec}\,\Bbbk[P + E]/\bar{I},$$

where $\bar{I} = I + E$.

On the other hand, letting $\bar{\Sigma}$ be the fan for $\bar{Y}$ in $\bar{B} = M_{\mathbb{R}}$, we have the piecewise linear function $\bar{\varphi} : \bar{B} \to P^{\mathrm{gp}}_{\mathbb{R}}$ with $p_{\bar{\rho},\bar{\varphi}} = p^*[\bar{D}_{\bar{\rho}}]$, as before. This now determines the Mumford family

$$\bar{f}_{\bar{I}} : \bar{X}_{\bar{I}} \to \mathrm{Spec}\,\Bbbk[P + E]/\bar{I}.$$

Our goal is to compare these two families. The restriction of either family to T is the trivial family $\mathbb{V}_n \times T \to T$. Thus either family contains a canonical copy of T, i.e., $\{0\} \times T$, where 0 is the vertex of $\mathbb{V}_n$.

In what follows, to simplify notation, replace P by $P + E$, G by $\mathfrak{m}_{P+E} = G + E$ and I by $\bar{I} = I + E$.

Proposition 3.40. *Fix an ideal I with $\sqrt{I}=G$. There are open affine sets $\bar{U}_I \subset \bar{X}_I$, $U_I \subset X_I$, both sets containing the canonical copy of T, and an isomorphism*

$$\mu_I : U_I \to \bar{U}_I$$

of families over $\operatorname{Spec} \Bbbk[P]/I$.

Moreover, there is a non-zero monomial $y \in \Bbbk[P]$ whose pullback to X_I is in the stalk at any point $x \in T \subset \mathbb{V}_n \times T$ of the ideal of $\operatorname{Sing}_{f_I}$ *for all I.*

Proof. The generic fibre of the Mumford family over $\operatorname{Spec} \Bbbk[P]$ is smooth: indeed the family is trivial over the open torus orbit of $\operatorname{Spec} \Bbbk[P]$, with fibre an algebraic torus. It follows that there is a non-zero monomial $y \in \Bbbk[P]$ in the ideal of $\operatorname{Sing}_{\bar{f}}$ for the global Mumford family

$$\bar{f} : \operatorname{Spec} \Bbbk[P_{\bar{\varphi}}] \to \operatorname{Spec} \Bbbk[P].$$

Of course its restriction then lies in the stalk at any point x of the ideal sheaf of $\operatorname{Sing}_{\bar{f}_I}$ for all I. Thus once we establish the claimed isomorphisms, the final statement follows.

Recall from §3.1 the construction of $\bar{\mathfrak{D}} := \nu(\mathfrak{D})$ and the scheme $\bar{X}^o_{I,\bar{\mathfrak{D}}}$ from $\bar{\mathfrak{D}}$. By Lemma 3.29, $X^o_{I,\mathfrak{D}} \cong \bar{X}^o_{I,\bar{\mathfrak{D}}}$, so we in fact have an isomorphism

$$X_I \cong \operatorname{Spec} \Gamma(\bar{X}^o_{I,\bar{\mathfrak{D}}}, \mathcal{O}_{\bar{X}^o_{I,\bar{\mathfrak{D}}}}) =: \bar{X}_{I,\bar{\mathfrak{D}}}.$$

So we can work with $\bar{X}_{I,\bar{\mathfrak{D}}}$ instead of X_I. On the other hand, the Mumford family $\bar{X}_I$ over $\operatorname{Spec} \Bbbk[P]/I$ can be described similarly. Using the empty scattering diagram instead of the scattering diagram $\bar{\mathfrak{D}}$, one has by Lemma 2.10

$$\bar{X}_I = \operatorname{Spec} \Gamma(\bar{X}^o_{I,\emptyset}, \mathcal{O}_{\bar{X}^o_{I,\emptyset}}).$$

Now define an ideal $I_0 \subset P_{\bar{\varphi}}$ as follows. For $\sigma \in \bar{\Sigma}_{\max}$, let $\bar{\varphi}_\sigma$ denote the linear extension of $\bar{\varphi}|_\sigma$. We set

$$I_0 := \{(m,p) \in P_{\bar{\varphi}} \mid p - \bar{\varphi}_\sigma(m) \in I \text{ for some } \sigma \in \bar{\Sigma}_{\max}\}.$$

Note that $\sqrt{I_0} = \mathfrak{m}_{P_{\bar{\varphi}}}$. By assumption, $\bar{\mathfrak{D}}$ is a scattering diagram for $P_{\bar{\varphi}}$, and hence there are only a finite number of $(\mathfrak{d}, f_\mathfrak{d}) \in \bar{\mathfrak{D}}$ for which $f_\mathfrak{d} \not\equiv 1 \mod I_0$. Furthermore, modulo I_0, each $f_\mathfrak{d}$ is a polynomial.

Let $\bar{\mathfrak{D}}_I$ be the scattering diagram obtained from $\bar{\mathfrak{D}}$ by, for each outgoing ray $(\mathfrak{d}, f_\mathfrak{d})$, truncating each $f_\mathfrak{d}$ by throwing out all terms which lie in I_0. The incoming rays remain unchanged. Thus $\bar{\mathfrak{D}}_I$ can be viewed as a finite scattering diagram. Let

$$h := \prod_{\mathfrak{d} \in \bar{\mathfrak{D}}_I} f_\mathfrak{d}.$$

This is an element of $\Bbbk[P_{\bar{\varphi}}]$. Note that necessarily $h \equiv 1 \mod \mathfrak{m}_{P_{\bar{\varphi}}}$. Thus $h \neq 0$ defines an open subset $\bar{U} \subset \operatorname{Spec} \Bbbk[P_{\bar{\varphi}}]/G\Bbbk[P_{\bar{\varphi}}] = \bar{X}_G = X_G = \mathbb{V}_n \times T$ which contains the canonical copy of T. Since $\bar{X}_I$ and X_I both have underlying topological space $\bar{X}_G$, this defines open sets $\bar{U}_I$ of $\bar{X}_I$ and U_I of X_I. We shall show these two sets are isomorphic.

Recall for $\tau_1, \tau_2 \in \Sigma$ with $\{0\} \neq \tau_1 \subset \tau_2$, we obtain from the ideal I the ring $R_{\tau_1,\tau_2,I} = \Bbbk[P_{\bar{\varphi}_{\tau_1}}]/I_{\tau_1,\tau_2}$. Since $h \in \Bbbk[P_{\bar{\varphi}}] \subset \Bbbk[P_{\bar{\varphi}_{\tau_1}}]$, h defines an element of $R_{\tau_1,\tau_2,I}$. Let

$$S_{\tau_1,\tau_2,I} := (R_{\tau_1,\tau_2,I})_h.$$

Given a path γ in $M_{\mathbb{R}} \setminus \{0\}$, note that by construction of h, $\theta_{\gamma,\bar{\mathfrak{D}}_I}$ makes sense as an automorphism of the localization $\Bbbk[P_{\bar{\varphi}}]_h$, since to define the automorphism associated with crossing a ray $(\mathfrak{d}, f_{\mathfrak{d}})$, we only need $f_{\mathfrak{d}}$ to be invertible. However since by construction h is divisible by $f_{\mathfrak{d}}$, $f_{\mathfrak{d}}$ is invertible. In particular, $\theta_{\gamma,\bar{\mathfrak{D}}_I}$ also makes sense as an automorphism of $S_{\tau_1,\tau_2,I}$ for any I.

Choose an orientation on $M_{\mathbb{R}}$, labelling the rays $\rho_1, \ldots, \rho_n$ of $\bar{\Sigma}$ in a counterclockwise order, with $\sigma_{i-1,i}$ as usual the maximal cone containing ρ_{i-1} and ρ_i. For two distinct points p, q on the unit circle in $M_{\mathbb{R}}$ not contained in $\mathrm{Supp}(\bar{\mathfrak{D}}_I)$, let $\gamma_{p,q}$ be a *counterclockwise* path from p to q, and write $\theta_{p,q}$ for $\theta_{\gamma_{p,q},\bar{\mathfrak{D}}_I}$ acting on any of the rings $S_{\tau_1,\tau_2,I}$.

For each ρ_i, let $p_{i,+}$ be a point on this unit circle contained in the connected component of $\sigma_{i,i+1} \setminus \mathrm{Supp}(\bar{\mathfrak{D}}_I)$ adjacent to ρ_i, and $p_{i,-}$ a point in this unit circle contained in the connected component of $\sigma_{i-1,i} \setminus \mathrm{Supp}(\bar{\mathfrak{D}}_I)$ adjacent to ρ_i.

Both $\bar{X}^o_{I,\emptyset}$ and $\bar{X}^o_{I,\bar{\mathfrak{D}}}$ are constructed by first constructing open sets $\bar{U}_{\rho_i,I}$, $U_{\rho_i,I}$, where both of these are given as spectra of fibred products

$$R_{\rho_i,\sigma_{i-1,i},I} \times_{(R_{\rho_i,\rho_i,I})_{f_{\rho_i}}} R_{\rho_i,\sigma_{i,i+1},I}.$$

In the Mumford case, the maps are just the canonical surjections, while in the case $\bar{X}^o_{I,\bar{\mathfrak{D}}}$, we can take $R_{\rho_i,\sigma_{i-1,i},I} \to (R_{\rho_i,\rho_i,I})_{f_{\rho_i}}$ to be the composition of the canonical map surjection followed by $\theta_{p_{i,-},p_{i,+}}$. Note that this automorphism is defined using the truncated scattering diagram $\bar{\mathfrak{D}}_I$ rather than $\bar{\mathfrak{D}}$. However, in fact this is the same automorphism. Indeed, given an outgoing ray $(\mathfrak{d}, f_{\mathfrak{d}}) \in \bar{\mathfrak{D}}$, any monomial z^p which appears in $f_{\mathfrak{d}}$ satisfies $-r(p) \in \tau_{\mathfrak{d}}$. One then applies the following claim to observe that modulo $I_{\tau_{\mathfrak{d}},\tau_{\mathfrak{d}}}$, the function $f_{\mathfrak{d}}$ and the truncated version coincide.

Claim 3.41. *Let $\tau \in \bar{\Sigma}$, and suppose $(m,p) \in P_{\bar{\varphi}}$ satisfies $-m \in \tau$. Then $(m,p) \in I_0$ if and only if $(m,p) \in I_{\tau,\tau}$.*

Proof of claim. Clearly $I_{\tau,\tau} \cap P_{\overline{\varphi}} \subset I_0$, so one implication is clear. Conversely, suppose that $(m,p) \in I_0$, so that $p - \overline{\varphi}_{\sigma}(m) \in I$ for some $\sigma \in \Sigma_{\max}$. If $\tau \subset \sigma' \in \mathscr{P}_{\max}$, let $\rho_1, \ldots, \rho_n$ be the sequence of rays traversed in passing from σ to σ', chosen so that all

$\rho_1, \ldots, \rho_n$ lie in a half-plane bounded by the line $\mathbb{R}m$. Then

$$\overline{\varphi}_{\sigma'}(m) = \overline{\varphi}_\sigma(m) + \sum_{i=1}^{n} \langle n_{\rho_i}, m\rangle p_{\rho_i,\overline{\varphi}}.$$

Note that since $-m \in \tau$, we must have $\langle n_{\rho_i}, m\rangle \leq 0$ for each i, and hence $p - \overline{\varphi}_{\sigma'}(m) = p - \overline{\varphi}_\sigma(m) + p'$ for some $p' \in P$. Hence $(m,p) \in I_{\tau,\tau}$. □

After localizing all these rings at h, we obtain open subsets $\bar{U}^h_{\rho_i,I}$ and $U^h_{\rho_i,I}$ of $\bar{U}_{\rho_i,I}$ and $U_{\rho_i,I}$ respectively as the spectra of the fibred products

$$S_{\rho_i,\sigma_{i-1,i},I} \times_{S_{\rho_i,\rho_i,I}} S_{\rho_i,\sigma_{i,i+1},I}$$

Choose a base-point q on the unit circle not in $\mathrm{Supp}(\bar{\mathfrak{D}}_I)$. We can easily write down an isomorphism between these two fibred products via the diagram

$$\begin{tikzcd}
S_{\rho_i,\sigma_{i-1,i},I} \arrow[rr, "\theta_{p_{i,-},q}"] \arrow[dr, "\theta_{p_{i,-},p_{i,+}}"] & & S_{\rho_i,\sigma_{i-1,i},I} \arrow[dr] & \\
& S_{\rho_i,\rho_i,I} \arrow[rr, "\theta_{p_{i,+},q}"] & & S_{\rho_i,\rho_i,I} \\
S_{\rho_i,\sigma_{i,i+1},I} \arrow[ur] \arrow[rr, "\theta_{p_{i,+},q}"] & & S_{\rho_i,\sigma_{i,i+1},I} \arrow[ur] &
\end{tikzcd}$$

Here unlabelled arrows are the canonical surjections, while decorated arrows are the canonical maps composed with the given automorphism. Since the diagram is clearly commutative, it induces an isomorphism between the respective fibred products, and hence an isomorphism $\bar{U}^h_{\rho_i,I} \to U^h_{\rho_i,I}$.

Next recall in the construction of $\bar{X}^o_{I,\emptyset}$, the open sets $\bar{U}_{\rho_i,I}$ and $\bar{U}_{\rho_{i+1},I}$ are glued together trivially along the common open set $\mathrm{Spec}\, R_{\sigma_{i,i+1},\sigma_{i,i+1},I}$, while in the construction of $\bar{X}^o_{I,\bar{\mathfrak{D}}}$, $U_{\rho_i,I}$ and $U_{\rho_{i+1},I}$ are glued together along the common open set $\mathrm{Spec}\, R_{\sigma_{i,i+1},\sigma_{i,i+1},I}$ using $\theta_{p_{i,+},p_{i+1,-}}$. On the other hand, we have an obviously commutative diagram

$$\begin{tikzcd}
S_{\rho_i,\sigma_{i,i+1},I} \arrow[rr, "\theta_{p_{i,+},q}"] \arrow[dr, "\theta_{p_{i,+},p_{i+1,-}}"] & & S_{\rho_i,\sigma_{i,i+1},I} \arrow[dr] & \\
& S_{\sigma_{i,i+1},\sigma_{i,i+1},I} \arrow[rr, "\theta_{p_{i+1,-},q}"] & & S_{\sigma_{i,i+1},\sigma_{i,i+1},I} \\
S_{\rho_{i+1},\sigma_{i,i+1},I} \arrow[ur] \arrow[rr, "\theta_{p_{i+1,-},q}"] & & S_{\rho_{i+1},\sigma_{i,i+1},I} \arrow[ur] &
\end{tikzcd}$$

This shows that the isomorphisms between $U^h_{\rho_i,I}$ and $\bar{U}^h_{\rho_i,I}$ are compatible with the gluings, hence giving an isomorphism between $U_I \setminus T$ and $\bar{U}_I \setminus T$.

Now $\mathbb{V}_n$ satisfies Serre's condition S_2. Since $\bar{X}_I$ and X_I are flat deformations of $\mathbb{V}_n \times T$, by Lemma 2.11 the above isomorphism extends across the codimension two set T, giving the desired isomorphism between U_I and $\bar{U}_I$. □

3.5. **The relative torus.** The flat deformations $X_{I,\mathfrak{D}^{\mathrm{can}}} \to \operatorname{Spec} \Bbbk[P]/I$ produced by the canonical scattering diagram have a useful special property: there is a natural torus action on the total space $X_{I,\mathfrak{D}^{\mathrm{can}}}$ compatible with a torus action on the base. The meaning of this action will be clarified in Part II, where we will prove that our family extends naturally, in the positive case, to a universal family of Looijenga pairs (Z, D) together with a choice of isomorphism $D \overset{\sim}{\to} D_*$, where D_* is a fixed n-cycle. The torus action then corresponds to changing the choice of isomorphism.

Fixing the pair (Y, D) as usual, $D = D_1 + \cdots + D_n$, let $\mathbb{A}^D = \mathbb{A}^n$ be the affine space with one coordinate for each component D_i. Let T^D be the diagonal torus acting on $\mathbb{A}^D$, i.e., the torus T^D whose character group

$$\chi(T^D) = \mathbb{Z}^D$$

is the free module with basis $e_{D_1}, \ldots, e_{D_n}$.

Definition 3.42. We have a canonical map $w : A_1(Y) \to \chi(T^D)$ given by

$$C \mapsto \sum (C \cdot D_i) e_{D_i}.$$

Suppose $P \subset A_1(Y)$ is a toric submonoid containing $\mathrm{NE}(Y)$. We then get an action of T^D on $\operatorname{Spec} \Bbbk[P]$, as well as on $\operatorname{Spec} \Bbbk[P]/I$ for any monomial ideal I, and hence also on $\operatorname{Spf}(\widehat{\Bbbk[P]})$ for any completion of $\Bbbk[P]$ with respect to a monomial ideal.

We can also define a unique piecewise linear map

$$w : B \to \chi(T^D) \otimes_{\mathbb{Z}} \mathbb{R}$$

with $w(0) = 0$ and $w(m_i) = e_{D_i}$, for m_i the primitive generator of the ray ρ_i.

Theorem 3.43. *Let $J \subset P$ be a radical ideal and I an ideal with $\sqrt{I} = J$ for which $X_{I,\mathfrak{D}^{\mathrm{can}}} \to \operatorname{Spec} \Bbbk[P]/I$ is defined. Then T^D acts equivariantly on $X_{I,\mathfrak{D}^{\mathrm{can}}} \to \operatorname{Spec} \Bbbk[P]/I$. Furthermore, each theta function ϑ_q, $q \in B(\mathbb{Z})$, is an eigenfunction of this action, with character $w(q)$.*

Proof. It's enough to check this on the open subset $X^o_{I,\mathfrak{D}^{\mathrm{can}}} \subset X_{I,\mathfrak{D}^{\mathrm{can}}}$. We have a cover of $X^o_{I,\mathfrak{D}^{\mathrm{can}}}$ by open sets the hypersurfaces

$$U_{i,I} \subset \mathbb{A}^2_{x_{i-1},x_{i+1}} \times (\mathbb{G}_m)_{x_i} \times \operatorname{Spec} \Bbbk[P]/I$$

given by the equation

$$x_{i-1}x_{i+1} = z^{[D_i]} x_i^{-D_i^2} f_{\rho_i},$$

where f_{ρ_i} is the function attached to the ray ρ_i in $\mathfrak{D}^{\mathrm{can}}$. If we act on x_j with weight $w(m_j)$ and z^p with weight $w(p)$ (for $p \in P$), then we note that for every $(\mathfrak{d}, f_{\mathfrak{d}}) \in \mathfrak{D}^{\mathrm{can}}$, every monomial in $f_{\mathfrak{d}}$ has weight zero by the explicit description of $f_{\mathfrak{d}}$ in Definition 3.3. In particular, the equation defining $U_{i,I}$ is clearly T^D equivariant, and each of the monomials is an eigenfunction.

Now $X^o_{I,\mathfrak{D}^{\mathrm{can}}}$ is obtained by gluing $U_{\rho_i,\sigma_{i,i+1},I} \subset U_{i,I}$ with $U_{\rho_{i+1},\sigma_{i,i+1}} \subset U_{i+1,I}$, using scattering automorphisms of in $\mathfrak{D}^{\mathrm{can}}$, and these open sets are naturally identified with $(\mathbb{G}_m)^2_{x_i,x_{i+1}} \times \operatorname{Spec} \Bbbk[P]/I$. The scattering automorphisms commute with the T^D action, by the fact that the scattering functions have weight zero. Thus T^D acts equivariantly on $X_{I,\mathfrak{D}^{\mathrm{can}}} \to \operatorname{Spec} \Bbbk[P]/I$.

Now we check our canonical global function ϑ_q is an eigenfunction, with character $w(q)$. By construction, given a broken line γ, the weights of monomials attached to adjacent domains of linearity are the same, since the functions in the scattering diagram are of weight zero. Thus the weight of $\operatorname{Mono}(\gamma)$ only depends on q. This weight can be determined by fixing the base point Q in a cone σ which contains q, in which case the broken line for q which doesn't bend and is wholly contained in σ yields the monomial $z^{\varphi_\sigma(q)}$, which has weight $w(q)$. Thus ϑ_q is an eigenfunction with weight $w(q)$. $\square$

4. LOOIJENGA'S CONJECTURE

In this section we apply our main theorem to give a proof of Looijenga's conjecture on smoothability of cusp singularities, Corollary 0.5. The simple conceptual idea is explained at the end of §0.6.2. Here we give the details. These are slightly involved. Indeed, the most natural construction of a cusp, the quotient construction of Example 1.9, is analytic, and we will have to deal with convergence issues to show that our construction extends to this analytic situation.

Let (Y, D) be a rational surface with anti-canonical cycle. Let $\sigma_P \subset A_1(Y, \mathbb{R})$ be a rational polyhedral cone containing $\operatorname{NE}(Y)$, $P = \sigma_P \cap A_1(Y, \mathbb{Z})$ the associated toric monoid, $J \subset P$ a radical monomial ideal, and I a monomial ideal such that $\sqrt{I} = J$. We write $S = \operatorname{Spec} \Bbbk[P]$ and $S_I = \operatorname{Spec} \Bbbk[P]/I$. In §4.1 and §4.2 below we use the field $\Bbbk = \mathbb{C}$.

4.1. **Cusp family.** Suppose that the intersection matrix $(D_i \cdot D_j)_{1 \le i,j \le n}$ is negative definite. Let $f\colon Y \to Y'$ be the contraction (in the analytic category) of $D \subset Y$ to a cusp singularity $q \in Y'$. We assume that f is the minimal resolution of Y', that is, $D_i^2 \le -2$ for all i. We further assume that $n \ge 3$. Let L be a nef divisor on Y such that

$$\operatorname{NE}(Y)_{\mathbb{R}_{\ge 0}} \cap L^{\perp} = \langle D_1, \ldots, D_n \rangle_{\mathbb{R}_{\ge 0}}.$$

Here the subscript $\mathbb{R}_{\geq 0}$ denotes the real cone in $H_2(Y,\mathbb{R})$ generated by the given elements or set. (If Y' is projective we can take $L = h^*A$ for A an ample divisor on Y'. In general, let A be an ample divisor on Y. There exist unique $a_i \in \mathbb{Q}$ such that $L := A + \sum a_i D_i$ is orthogonal to D_j for each j. By negative definiteness of $\langle D_1, \ldots, D_n \rangle$, we have $a_i > 0$ for each i. It follows that L is nef.) Assume that σ_P is strictly convex and $\sigma_P \cap L^\perp$ is a face of σ_P. Let $\mathfrak{m} = P \setminus \{0\}$ and $J = P \setminus P \cap L^\perp \subset P$, the radical monomial ideal associated to the face $\sigma_P \cap L^\perp$ of σ_P.

Theorem 4.1. *There exists an analytic open neighbourhood S'_J of $0 \in S_J$ and an analytic flat family $f_J \colon X_J \to S'_J$ together with a section $s \colon S'_J \to X_J$ satisfying the following properties:*

1. *The general fibre $X_{J,t}$ of f_J is a Stein analytic surface with a unique singularity $s(t) \in X_{J,t}$ isomorphic to the dual cusp to $q \in Y'$.*
2. *Fix $R > 1$. For each ray $\rho \in \Sigma$ there is an open analytic subset $V_{\rho,J} \subset X_J$ and open analytic embeddings*

   $$V_{\rho,J} \subset \{(X_-, X_+, X) \in U_{\rho,J} \mid |X_-| < R|X|,\, |X_+| < R|X|\} \subset U_{\rho,J}$$

   where

   $$U_{\rho,J} := V(X_-X_+ - z^{[D_\rho]}X^{-D_\rho^2}) \subset \mathbb{A}^2_{X_-,X_+} \times (\mathbb{G}_m)_X \times S_J$$

   such that
   - (a) $X^o_J := X_J \setminus s(S'_J) = \bigcup_{\rho\in\Sigma} V_{\rho,J}$
   - (b) $V_{\rho,J} \cap V_{\rho',J} = \emptyset$ *unless $\rho = \rho'$ or ρ and ρ' are the edges of a maximal cone $\sigma \in \Sigma$.*
3. *The restriction of X_J/S'_J to $S_{J+\mathfrak{m}^{N+1}}$ is identified with an analytic neighbourhood of the vertex in the restriction of the family $X_{\mathfrak{m}^{N+1}}/S_{\mathfrak{m}^{N+1}}$ given by Theorem 2.33, (1) with $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$, for each $N \geq 0$.*

Proof. (1) We use the notation of Example 1.9, so that we have an infinite fan $\tilde{\Sigma}$ in $M_\mathbb{R}$ with the primitive generators of the rays being the v_i for $i \in \mathbb{Z}$. We also have $T \in \mathrm{SL}(M)$ acting on the fan $\tilde{\Sigma}$. We have $B = |\tilde{\Sigma}|/\Gamma$, where Γ is the group generated by T.

Now let $\Xi \subset M_\mathbb{R}$ be the convex hull of the points $v_i \in M$. Thus Ξ is an infinite convex polytope. Let $\tilde{\Sigma}'$ be the subdivision of Ξ induced by $\tilde{\Sigma}$. We can build a Mumford degeneration Z/S_J with special fibre Z_0 the stable toric variety associated to $\tilde{\Sigma}'$ (that is, Z_0 is the union of the toric surfaces associated to the maximal polytopes in $\tilde{\Sigma}'$), as follows.

S_J is the affine toric variety associated to the face $\sigma_{\mathrm{bdy}} := \sigma_P \cap L^\perp$ of σ_P, and $P_{\mathrm{bdy}} := \sigma_{\mathrm{bdy}} \cap H_2(Y,\mathbb{Z})$ contains the classes of the components of the boundary of Y.

We define a piecewise linear convex function $\varphi\colon |\tilde{\Sigma}'| \to P_{\mathrm{bdy}}$ by

$$\varphi_{\sigma_+} - \varphi_{\sigma_-} = n_\rho \otimes p_\rho$$

where σ_+, σ_- are two polytopes with common edge ρ, $\varphi_{\sigma_\pm}$ are as usual the affine linear extensions of $\varphi|_{\sigma_\pm}$, and $n_\rho \in M^*$ is the primitive element annihilating ρ and positive on σ_+. Finally, $p_\rho = [D_{i \bmod n}] \in P$ if $\rho = v_i + \mathbb{R}_{\geq 0} \cdot v_i$. This determines φ modulo an integral affine function.

Then φ determines a Mumford degeneration: this is a slight generalization of §1.2. One defines

$$\tilde{\Xi} := \{(m, r) \,|\, m \in \Xi, r \in \varphi(m) + \sigma_{\mathrm{bdy}}\} \subset M_{\mathbb{R}} \oplus (P^{\mathrm{gp}}_{\mathrm{bdy}} \otimes_{\mathbb{Z}} \mathbb{R}).$$

Let $C(\tilde{\Xi})$ be the closure of

$$\{(sm, sr, s) \,|\, (m, r) \in \tilde{\Xi}, s \geq 0\} \subset M_{\mathbb{R}} \oplus (P^{\mathrm{gp}}_{\mathrm{bdy}} \otimes_{\mathbb{Z}} \mathbb{R}) \oplus \mathbb{R}.$$

Then $\mathbb{C}[C(\tilde{\Xi}) \cap (M \oplus P^{\mathrm{gp}}_{\mathrm{bdy}} \oplus \mathbb{Z})]$ has a natural grading given by the last coordinate, and the degree zero part of this ring is easily seen to contain $\mathbb{C}[P_{\mathrm{bdy}}]$. Thus we obtain the Mumford family determined by φ as

$$Z := \operatorname{Proj} \mathbb{C}[C(\tilde{\Xi}) \cap (M \oplus P^{\mathrm{gp}}_{\mathrm{bdy}} \oplus \mathbb{Z})] \to \operatorname{Spec} \mathbb{C}[P_{\mathrm{bdy}}] = S_J.$$

One sees easily that the fibre over $0 \in S_J$ of $Z \to S_J$ has infinitely many components indexed by the 2-cells of the subdivision $\tilde{\Sigma}'$ of Ξ, each of which is a copy of the blowup of $\mathbb{A}^2$ at the origin. The general fibre is a toric surface (only locally of finite type) containing an infinite chain of smooth rational curves, which specializes to the union of the exceptional curves of the blowups in Z_0. By construction Γ acts on Z over S_J (because φ is Γ-invariant modulo integral affine functions). Let $E \subset Z/S_J$ be the family of curves described above (the relative toric boundary). The group Γ acts properly discontinuously on a tubular neighbourhood N of $E \subset Z$ (cf. [AMRT75], p. 48). Let $p\colon (F \subset \tilde{X}) \to S_J$ denote the quotient of $(E \subset N) \to S_J$ by Γ.

The divisor $F \subset \tilde{X}$ is Cartier and the dual of its normal bundle is relatively ample over a neighbourhood of $0 \in S_J$. Indeed, the special fibre $\tilde{X}_0$ is a union of n irreducible components each isomorphic to a tubular neighbourhood of the exceptional locus in the blowup of $\mathbb{A}^2$, and $F_0 \subset \tilde{X}_0$ is the cycle of n smooth rational curves formed by the exceptional curves of the blowups. Hence the normal bundle of F_0 in $\tilde{X}_0$ has degree -1 on each component of F_0. Moreover, we have $R^1 p_*(\mathcal{N}^\vee_{F/\tilde{X}})^{\otimes k} = 0$ for each $k > 0$. Indeed, by cohomology and base change it suffices to show that $H^1((\mathcal{N}^\vee_{F_0/\tilde{X}_0})^{\otimes k}) = 0$, and this follows from Serre duality. Now by a relative version of Grauert's contractibility criterion [F75], Thm. 2, taking global sections of the structure sheaf defines a contraction $p\colon \tilde{X} \to X_J/S_J$ to a family of Stein analytic spaces with exceptional locus F. The general fibre $X_{J,t}$ of X_J/S_J is the dual cusp by Lemma 4.2 below.

We now show that X_J/S_J is flat and the special fibre is the neighbourhood of the n-vertex obtained by contracting $F_0 \subset \tilde{X}_0$. The key point is that $R^1 p_* \mathcal{O}_{\tilde{X}}$ is a locally free $\mathcal{O}_{S_J}$-module, cf. [W76], Theorem 1.4(b). Indeed, we have

$$R^1 p_* \mathcal{O}_{\tilde{X}}(-F) = 0$$

by cohomology and base change, the theorem on formal functions, and the vanishing $H^1((\mathcal{N}^\vee_{F_0/\tilde{X}_0})^{\otimes k}) = 0$ for $k > 0$ used above. So, pushing forward the exact sequence

$$0 \to \mathcal{O}_{\tilde{X}}(-F) \to \mathcal{O}_{\tilde{X}} \to \mathcal{O}_F \to 0$$

we obtain

$$R^1 p_* \mathcal{O}_{\tilde{X}} = R^1 p_* \mathcal{O}_F \simeq \mathcal{O}_{S_J}.$$

Recall that S_J is a toric variety, so in particular Cohen-Macaulay. Let $t_1, \ldots, t_r$ be a regular sequence at $0 \in S_J$ of length $\dim S_J$ and write

$$S_J^i = V(t_1, \cdots, t_i) \subset S_J,$$

$\tilde{X}^i = \tilde{X}|_{S_J^i}$, and let X_J^i/S_J^i be the family over S_J^i defined by $\mathcal{O}_{X_J^i} = p_* \mathcal{O}_{\tilde{X}^i}$. Arguing as above we find that $R^1 p_* \mathcal{O}_{\tilde{X}^i} \simeq \mathcal{O}_{S_J^i}$. Pushing forward the exact sequence

$$0 \to \mathcal{O}_{\tilde{X}^i} \xrightarrow{\cdot t_{i+1}} \mathcal{O}_{\tilde{X}^i} \longrightarrow \mathcal{O}_{\tilde{X}^{i+1}} \longrightarrow 0$$

we deduce that the natural map

$$\mathcal{O}_{X_J^i}/t_{i+1}\mathcal{O}_{X_J^i} \to \mathcal{O}_{X_J^{i+1}} \tag{4.1}$$

is an isomorphism. Hence by the local criterion of flatness [Ma89], Ex. 22.3, p. 178, it suffices to show that X_J^r/S_J^r is flat with special fibre the n-vertex. But S_J^r is the spectrum of an Artinian $\mathbb{C}$-algebra, so this follows from [W76], Theorem 1.4(b).

(2) Write $Z^o = Z \setminus E$, and $m_i = -D^2_{i \bmod n}$, $a_i = z^{[D_{i \bmod n}]}$ for $i \in \mathbb{Z}$. We have an open covering

$$Z^o = \bigcup_{i \in \mathbb{Z}} U_{i,J},$$

where

$$U_{i,J} = V(x_{i-1}x_{i+1} - a_i x_i^{m_i}) \subset \mathbb{A}^2_{x_{i-1},x_{i+1}} \times (\mathbb{G}_m)_{x_i} \times S_J.$$

Similarly, we have an open covering

$$Z = \bigcup_{i \in \mathbb{Z}} \bar{U}_{i,J},$$

where

$$\begin{aligned}\bar{U}_{i,J} &= V(x'_{i-1}x'_{i+1} - a_i x_i^{m_i-2}) \subset \mathbb{C}^3_{x'_{i-1},x_i,x'_{i+1}} \times S_J,\\ F \cap \bar{U}_{i,J} &= V(x_i) \subset \bar{U}_{i,J},\end{aligned}$$

and $\bar{U}_{i,J} \setminus F = U_{i,J}$ via $x_{i-1} = x_i x'_{i-1}$, $x_{i+1} = x_i x'_{i+1}$. (Note that $m_i = -D_i^2 \geq 2$ by assumption.)

Recall that the infinite cyclic group Γ acts on Z/S_J, there is a Γ-invariant tubular neighbourhood $N \subset Z$ of $E \subset Z$ on which the action is properly discontinuous, and $F \subset \tilde{X}$ is obtained as the quotient of $E \subset N$ by Γ. In terms of the open covering above the action is given by $U_{i,J} \to U_{i+n,J}$, $x_j \mapsto x_{j+n}$. Note that the map $U_{i,J} \cap N \to \tilde{X}$ is *not* an open embedding (because, for example, $U_{i,J}$ contains the general fibre of Z^o/S_J). Fix $R > 1$. We define $W_i \subset U_{i,J} \cap N$ by

$$W_i = \{(x_{i-1}, x_i, x_{i+1}) \in U_{i,J} \cap N \mid |x_{i-1}| < R|x_i|,\ |x_{i+1}| < R|x_i|\}$$

and similarly define $\overline{W}_i \subset \bar{U}_{i,J} \cap N$ by

$$\overline{W}_i = \{(x'_{i-1}, x_i, x'_{i+1}) \in \bar{U}_{i,J} \cap N \mid |x'_{i-1}| < R, |x'_{i+1}| < R\}.$$

Then $\overline{W}_i \setminus E = W_i$.

The $\overline{W}_i$ cover the special fibre E_0 of E/S_J (using $R > 1$). The open set $\bigcup \overline{W}_i \subset N$ is Γ-invariant, the quotient $(N, E) \to (\tilde{X}, F)$ by Γ is a covering map, and $p\colon \tilde{X} \to X_J$ is proper with exceptional locus F. Hence we may assume (passing to an analytic neighbourhood of $0 \in S_J$ and $s(0) \in X_J$) that $N = \bigcup \overline{W}_i$.

By Lemma 4.4 below there exists $\delta > 0$ such that

$$\overline{W}_i \cap \{(x_{i-1}, x_i, x_{i+1}) \,|\, |x_i| < \delta\} \subset \{|x_j| < 1 \quad \forall j\}$$

for each i. We replace $\overline{W}_i$ by $\overline{W}_i \cap \{|x_i| < \delta\}$, and modify W_i similarly. Then as above we may assume that $N = \bigcup \overline{W}_i$, and $N \subset \{|x_i| < 1 \quad \forall i\}$. We claim that $W_i \cap W_j = \emptyset$ for all $j > i+1$. It suffices to work on the general fibre of Z^o/S_J, which is an algebraic torus. The coordinate functions x_i are characters of this torus (up to a multiplicative constant). By construction we have $|x_i| < 1$ for each i on N. Hence, shrinking the base S_J, we may assume that $|a_i x_i^{m_i-2}| < 1/R^2$ for each i. The relation

$$x_{i-1}x_{i+1} = a_i x_i^{m_i}$$

gives the inequality

$$|x_{i+1}/x_i| < (1/R^2)|x_i/x_{i-1}|.$$

Combining such inequalities we obtain

$$|x_{j+1}/x_j| < (1/R^2)^{j-i}|x_{i+1}/x_i| \quad \text{for } j > i. \tag{4.2}$$

Now $|x_{i+1}/x_i| < R$ on W_i and $|x_{j-1}/x_j| < R$ on W_j, so $|x_j/x_{j-1}| > (1/R^2)|x_{i+1}/x_i|$ on $W_i \cap W_j$. For $j > i+1$ this contradicts the inequality (4.2), hence $W_i \cap W_j = \emptyset$ as claimed.

It follows that W_i embeds in X_J. Let V_i denote its image, with indices now understood modulo n. Thus $X_J^o = \bigcup V_i$ and the inverse image of V_i is the (disjoint) union of W_j such that $j \equiv i \bmod n$. We have $V_i \cap V_j = \emptyset$ for $j \neq i-1, i, i+1$ by our claim above. So, writing $V_{\rho,J} := V_i$ for ρ the ray of Σ corresponding to D_i, the condition (2)(b) is satisfied.

(3) We have the open covering $X_J^o = \bigcup V_i$ and open embeddings $V_i \subset U_{i,J}$, and an open covering $X^o_{\mathfrak{m}^{N+1}} = \bigcup U_{i,\mathfrak{m}^{N+1}}$. The restrictions of $U_{i,J}/S_J$ and $U_{i,\mathfrak{m}^{N+1}}/S_{\mathfrak{m}^{N+1}}$ to $S_{J+\mathfrak{m}^{N+1}}$ are identified, and the gluing maps coincide. It follows that the restriction of X_J/S_J is identified with a neighbourhood of the vertex in the restriction of $X_{\mathfrak{m}^{N+1}}/S_{\mathfrak{m}^{N+1}}$ using Lemma 2.11. □

Lemma 4.2. *Let $q \in Y'$ be a cusp singularity and $f\colon Y \to Y'$ the minimal resolution of $q \in Y'$, with exceptional locus $D = D_1 + \cdots + D_n$ a cycle of smooth rational curves. Define vectors $v_i \in \mathbb{Z}^2$ for $i \in \mathbb{Z}$ by $v_0 = \begin{pmatrix}1\\0\end{pmatrix}$, $v_1 = \begin{pmatrix}0\\1\end{pmatrix}$ and $v_{i-1}+v_{i+1} = (-D^2_{i \bmod n})v_i$ for each $i \in \mathbb{Z}$. Let $\Xi \subset \mathbb{R}^2$ be the convex hull of the v_i. Then Ξ is an unbounded convex polytope and the v_i are the integral points lying on the boundary of Ξ. Let $T \in \mathrm{SL}(2,\mathbb{Z})$ be the matrix such that $Tv_0 = v_n$ and $Tv_1 = v_{n+1}$. Then $\Gamma := \langle T \rangle \simeq \mathbb{Z}$ acts properly discontinuously on Ξ.*

Let Z be the quasiprojective toric variety (only locally of finite type) associated to Ξ. Let $E \subset Z$ be the toric boundary of Z, an infinite chain of smooth rational curves corresponding to the boundary of Ξ. Then there exists a tubular neighbourhood $E \subset N \subset Z$ such that the Γ action on Ξ induces a properly discontinuous Γ action on N. Let $F \subset \tilde{X}$ denote the quotient of $E \subset N$ by Γ. So F is a cycle of smooth rational curves. Then $F \subset \tilde{X}$ can be contracted to a singularity $p \in X$, which is a copy of the dual cusp to $q \in Y'$. Moreover, $\tilde{X}$ is obtained from the minimal resolution of $p \in X$ by contracting all the (-2)-curves.

Proof. Let Σ be the fan in $\mathbb{R}^2$ with maximal cones $\sigma_{i,i+1} = \langle v_i, v_{i+1}\rangle_{\mathbb{R}_{\geq 0}}$ for $i \in \mathbb{Z}$. Let $\overline{C}$ denote the closure of the support of Σ. Let Σ' be the normal fan for the polytope Ξ and $\overline{C}'$ the closure of its support. We observe that $\overline{C}'$ coincides with the dual of $\overline{C}$, together with the induced Γ-action. It follows that $\tilde{X}$ is a partial resolution of a copy of the dual cusp. See Example 1.9.

The surface $\tilde{X}$ has Du Val singularities of type A. Indeed, v_i is a vertex of Ξ iff $m_i := -D_i^2 > 2$. The corresponding point of Z is smooth if $m_i = 3$ and a singularity of type A_{m_i-3} if $m_i > 3$ (by direct calculation using $v_{i-1}+v_{i+1} = m_i v_i$). Also $K_{\tilde{X}}$ is relatively ample over X by Lemma 4.3 below. Indeed, the vectors $u_i := v_{i+1} - v_i$ are the primitive integral vectors in the direction of the edges of Ξ, and

$$u_i - u_{i-1} = v_{i+1} + v_{i-1} - 2v_i = (m_i - 2)v_i,$$

so the lines $v_i + \mathbb{R} \cdot (u_i - u_{i-1}) = \mathbb{R} \cdot v_i$ all meet at the origin. We deduce that $\tilde{X}$ is obtained from the minimal resolution of X by contracting all (-2)-curves as claimed. ☐

Lemma 4.3. *Let $P \subset \mathbb{R}^2$ be a rational convex polygon and X the associated toric surface. Fix an orientation and let e_0, e_1, e_2 denote consecutive edges of the boundary of P. Let $C \subset X$ denote the component of the toric boundary associated to the bounded edge $e_1 \subset P$. Let $v_0 = e_0 \cap e_1$ and $v_1 = e_1 \cap e_2$ be the vertices of e_1. Let $u_0, u_1, u_2 \in \mathbb{Z}^2$ denote the primitive integral vectors in the direction of e_0, e_1, e_2. Then $K_X \cdot C > 0$ iff the lines $v_0 + \mathbb{R}(u_1 - u_0)$ and $v_1 + \mathbb{R}(u_2 - u_1)$ meet on the opposite side of e_1 to P.*

Proof. This is an elementary toric calculation. When rewritten in terms of the normal fan of P it is the criterion of [R83], 4.3. ☐

Lemma 4.4. *We use the notation of the proof of Theorem 4.1(2). Let x_j be the coordinate functions on $Z^o = \bigcup U_{i,J}$. There exists $\delta > 0$ such that on each open set $U_{i,J}$ if $|x_i| < \delta$, $|x_{i-1}| < R|x_i|$, $|x_{i+1}| < R|x_i|$, and $|z^p| < 1$ for all $p \in P_{\mathrm{bdy}}$ then $|x_j| < 1$ for all j.*

Proof. The points v_{i-1}, v_i, and v_{i+1} are consecutive integral points on the boundary of the infinite convex polytope Ξ with asymptotic directions w_1, w_2. It follows that $w_1 = \alpha_{i1} v_i + \beta_{i1}(v_{i-1} - v_i)$ and $w_2 = \alpha_{i2} v_i + \beta_{i2}(v_{i+1} - v_i)$ for some $\alpha_{i1}, \beta_{i1}, \alpha_{i2}, \beta_{i2} \in \mathbb{R}_{>0}$. Note that the ratios $\beta_{i1}/\alpha_{i1}, \beta_{i2}/\alpha_{i2}$ only depend on i modulo n (because w_1, w_2 are eigenvectors of T and $T(v_i) = v_{i+n}$). Let μ be the maximum of the ratios $\beta_{i1}/\alpha_{i1}, \beta_{i2}/\alpha_{i2}$ for $i = 1, \ldots, n$. Let $\delta = R^{-\mu}$. If $j > i$ then $v_j = \alpha v_i + \beta(v_{i+1} - v_i)$ with $\beta/\alpha < \beta_{i2}/\alpha_{i2}$. The coordinate function x_j can be written as $z^p x_i^{\alpha}(x_{i+1}/x_i)^{\beta}$ on V_i, where $p \in P_{\mathrm{bdy}}$. Thus $|x_j| < \delta^{\alpha} R^{\beta} < 1$ for $|x_i| < \delta$ and $|z^p| < 1$. The same is true for $j < i$ by symmetry. ☐

4.2. **Thickening of cusp family.**

Theorem 4.5. *Let $p_J \colon X_J \to S'_J$ be the analytic family of Theorem 4.1. Possibly after replacing S'_J by a smaller neighbourhood of $0 \in S_J$ and X_J by a smaller neighbourhood of $s(S'_J) \subset X_J$, the following holds. Let $I \subset P$ be a monomial ideal such that $\sqrt{I} = J$ and let $S'_I \subset S_I$ denote the induced thickening of $S'_J \subset S_J$. There is an infinitesimal deformation $p_I \colon X_I \to S'_I$ of $p_J \colon X_J \to S'_J$ such that for each $N > 0$ the restriction to $\operatorname{Spec} \Bbbk[P]/(I + \mathfrak{m}^{N+1})$ is identified with an analytic neighbourhood of the vertex in* *the restriction of the family* $X_{\mathfrak{m}^{N+1}}/S_{\mathfrak{m}^{N+1}}$ *given by Theorem 2.33, (1) applied with* $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$.

As usual, $\mathfrak{D}^{\mathrm{can}}$ is the canonical scattering diagram on B associated to the pair (Y, D). Let $\mathfrak{D}$ be the scattering diagram obtained by reducing $\mathfrak{D}^{\mathrm{can}}$ modulo I as follows: For each ray $\mathfrak{d}$ of $\mathfrak{D}^{\mathrm{can}}$, we truncate the attached function $f_{\mathfrak{d}}$ by removing monomial terms lying in $I \cdot \Bbbk[P_{\varphi_{\tau_{\mathfrak{d}}}}]$, and we discard the ray if the truncated function equals 1. Thus $\mathfrak{D}$ has only finitely many rays $\mathfrak{d}$ and the attached functions $f_{\mathfrak{d}}$ are finite sums of monomials.

We use the scattering diagram $\mathfrak{D}$ to define a complex analytic space X_I^o/S_I' as follows. Recall from Lemma 2.19 the description of the schemes $U_{\rho,I}/S_I$ for $\rho \in \Sigma$: if $\sigma_-, \sigma_+ \in \Sigma$ are the maximal cones containing ρ,

$$U_{\rho,I} = V(X_-X_+ - z^{[D_\rho]}X^{-D_\rho^2}f_\rho) \subset \mathbb{A}^2_{X_-,X_+} \times (\mathbb{G}_m)_X \times S_I.$$

We have open subsets $U_{\rho,\sigma_\pm,I} \subset U_{\rho,I}$ defined by $X_\pm \neq 0$. For $\sigma \in \Sigma$ a maximal cone with edges ρ, ρ' we have canonical identifications

$$U_{\rho,\sigma,I} = U_{\sigma,\sigma,I} = U_{\rho',\sigma,I} \tag{4.3}$$

where

$$U_{\sigma,\sigma} = (\mathbb{G}_m)^2_{X,X'} \times S_I.$$

Recall that we have an open covering $X_J^o = \bigcup_{J\in\Sigma} V_{\rho,J}$ and open analytic embeddings $V_{\rho,J} \subset U_{\rho,J}$. For a maximal cone $\sigma \in \Sigma$ with edges ρ, ρ', write

$$V_{\sigma,\sigma,J} := (V_{\rho,J} \cap U_{\rho,\sigma,J}) \cap (V_{\rho',J} \cap U_{\rho',\sigma,J}) \subset U_{\sigma,\sigma,J}$$

where we use the identification (4.3). Let $V_{\rho,\sigma,J} \subset V_{\rho,J}$, $V_{\rho',\sigma,J} \subset V_{\rho',J}$ denote the open subsets corresponding to $V_{\sigma,\sigma,J}$ under (4.3). Let $V_{\rho,I}$, $V_{\rho,\sigma,I}$, etc., be the infinitesimal thickenings of these open sets determined by the thickenings $U_{\rho,I}$ of $U_{\rho,J}$. Let $\sigma \in \Sigma$ be a maximal cone with edges ρ, ρ'. Let $\theta_{\gamma,\mathfrak{D}} \colon U_{\rho',\sigma,I} \to U_{\rho,\sigma,I}$ be the gluing isomorphism defined as in §2.2. Note that the canonical scattering diagram $\mathfrak{D}^{\mathrm{can}}$ is trivial modulo J (because $\langle D_1, \ldots, D_n\rangle_{\mathbb{Q}}$ does not contain any $\mathbb{A}^1$-classes). Hence $\theta_{\gamma,\mathfrak{D}}$ restricts to the identification (4.3) modulo J, and thus restricts to an isomorphism

$$V_{\rho',\sigma,I} \to V_{\rho,\sigma,I}.$$

Gluing the $V_{\rho,I}$ via these isomorphisms we obtain an infinitesimal deformation X_I^o/S_I' of X_J^o/S_J'. Note that there are no triple overlaps of the $V_{\rho,J}$ by Theorem 4.1(2)(b), hence no compatibility condition for the gluing automorphisms. It is clear from the construction that the families X_I^o/S_I' and $X_{\mathfrak{m}^{N+1}}/S_{\mathfrak{m}^{N+1}}$ are compatible.

Proof of Theorem 4.5. We define sections $\vartheta_q \in \Gamma(X_I^o, \mathcal{O}_{X_I^o})$ for $q \in B(\mathbb{Z})$, compatible with the sections of Theorem 2.33, (2). We proceed as in the algebraic case: we first define a local section $\mathrm{Lift}_Q(q)$ for each choice of basepoint $Q \in B_0 \setminus \mathrm{Supp}(\mathfrak{D})$ on a corresponding open patch of X_I^o using the broken lines construction. The new difficulty here is that the functions $\mathrm{Lift}_Q(q)$ are not algebraic, even over the unthickened locus S'_J. Indeed, by definition $\mathrm{Lift}_Q(q) = \sum \mathrm{Mono}(\gamma)$ is a formal sum of monomials corresponding to broken lines γ for q with endpoint Q. This sum has infinitely many terms in the present case and so we must prove convergence. This is done in §4.2.1, see Propositions 4.8 and 4.9.

Once this convergence is proved, we observe that these patch to give well defined global sections. This follows from the consistency of $\mathfrak{D}^{\mathrm{can}}$ and compatibility of X_I^o/S'_I with $X^o_{\mathfrak{m}^{N+1}}/S_{\mathfrak{m}^{N+1}}$ for $N \geq 0$.

We define an infinitesimal thickening X_I/S'_I of X_J/S'_J by $\mathcal{O}_{X_I} = i_*\mathcal{O}_{X_I^o}$ where $i\colon X_J^o \subset X_J$ is the inclusion. Then X_I/S'_I is flat by Lemma 2.34 and the existence of the lifts ϑ_q. □

4.2.1. *Convergence of Lifts.* Let $C \subset M_{\mathbb{R}}$ be the closure of the support $|\tilde{\Sigma}|$ of the fan $\tilde{\Sigma}$, a closed convex cone. Let w_1, w_2 be generators of C. Then w_1, w_2 are eigenvectors of T with eigenvalues λ^{-1}, λ for some $\lambda \in \mathbb{R}$. We may assume that $\lambda > 1$. Let $\pi\colon \tilde{B}_0 \to B_0$ denote the universal cover of B_0. So $\tilde{B}_0$ is identified with the interior $\mathrm{Int}(C)$ of C, with deck transformations given by the action of $\Gamma = \langle T \rangle$ on C. Let $\tilde{\mathcal{P}}$, $\tilde{\varphi}$, and $\tilde{B}_0(\mathbb{Z})$ denote the pullbacks of $\mathcal{P}$, φ, and $B_0(\mathbb{Z})$. Let $\tilde{\mathfrak{D}}$ denote the scattering diagram on $\tilde{B}_0$ induced by $\mathfrak{D}$. We fix a trivialization of $\tilde{\mathcal{P}}$ as the constant sheaf with fibre $P^{\mathrm{gp}} \oplus M$.

The behaviour of the broken lines γ is best studied by passing to the universal cover $\tilde{B}_0$ of B_0. Let $Q \in B_0$, $q \in B_0(\mathbb{Z})$, and choose lifts $\tilde{Q} \in \tilde{B}_0$, $\tilde{q} \in \tilde{B}_0(\mathbb{Z})$. Then a broken line γ on B_0 for q with endpoint Q lifts uniquely to a broken line $\tilde{\gamma}$ on $\tilde{B}_0$ for $T^N(\tilde{q})$ with endpoint $\tilde{Q}$, for some $N \in \mathbb{Z}$, and the attached monomials are identified via $\tilde{\mathcal{P}} = \pi^*\mathcal{P}$. Note that $T^N(\tilde{q})$ approaches $\mathbb{R}_{\geq 0} \cdot w_2$ as $N \to \infty$ and $\mathbb{R}_{\geq 0} \cdot w_1$ as $N \to -\infty$.

If γ is a broken line for a point $\tilde{q} \in \tilde{B}_0(\mathbb{Z})$ with endpoint $\tilde{Q} \in \tilde{B}_0$ then $\gamma\colon (-\infty, 0] \to \tilde{B}_0$ is a piecewise linear path in $\tilde{B}_0 = \mathrm{Int}\, C$ with initial direction $-\tilde{q}$, ending at $\tilde{Q}$, and crossing all the rays between $\mathbb{R}_{\geq 0}\tilde{q}$ and $\mathbb{R}_{\geq 0}\tilde{Q}$ in order. Let $t_1, \ldots, t_l \in (-\infty, 0)$ denote the points where γ is not affine linear. Each point $\gamma(t_i)$ lies on a ray $\mathfrak{d}_i$ of the scattering diagram $\tilde{\mathfrak{D}}$ and the change $\gamma'(t_i + \epsilon) - \gamma'(t_i - \epsilon)$ in the direction of γ as it crosses $\mathfrak{d}_i$ is an integral multiple of the primitive generator of $\mathfrak{d}_i$. Moreover this multiple is positive because each ray of the canonical scattering diagram is an outgoing ray in the terminology of Definition 2.17. So the path γ is "convex when viewed from the origin".

It is convenient for the convergence calculation to decompose the monomials for broken lines as follows. Let $\gamma\colon (-\infty, 0] \to \tilde{B}_0$ be a broken line, $t \in (-\infty, 0]$ a point such that γ is affine linear near t and $\gamma(t)$ lies in the interior of a maximal cone σ of

$\tilde{\Sigma}$, and cz^q the monomial attached to the domain of linearity of γ containing t. Here $c \in \Bbbk$ and $q \in P_{\tilde{\varphi}_\sigma} \subset P^{\mathrm{gp}} \oplus M$. We write $cz^q = az^{\tilde{\varphi}_\sigma(m)}$, where $m = r(q) \in M$ and $a = cz^{q-\tilde{\varphi}_\sigma(m)}$ is a monomial in $\Bbbk[P]$. We also use the same decomposition for the monomials occuring in the scattering functions $f_{\mathfrak{d}}$ for $\mathfrak{d} \in \tilde{\mathfrak{D}}$. Let $\mathfrak{d}$ be a ray in $\tilde{\mathfrak{D}}$ with primitive integral generator $m \in M$ and $\tau = \tau_{\mathfrak{d}}$ the smallest cone of $\tilde{\Sigma}$ containing $\mathfrak{d}$. Then $f_{\mathfrak{d}} - 1$ is a sum of monomials cz^q where $c \in \Bbbk$ and $q \in P_{\tilde{\varphi}_\tau}$, $0 \neq -r(q) \in \mathfrak{d}$. We write $cz^q = az^{\tilde{\varphi}_\tau(m)}$ where $m = r(q)$ and $a = cz^{q-\tilde{\varphi}_\tau(m)}$ is a monomial in $\Bbbk[P]$.

The scattering diagram $\mathfrak{D}$ on B is finite (because we have reduced modulo I), and $\tilde{\mathfrak{D}}$ is its inverse image under $\pi\colon \tilde{B}_0 \to B_0$. Thus $\tilde{\mathfrak{D}}$ has only finitely many Γ-orbits of rays. Moreover, the Γ-action on the scattering functions $f_{\mathfrak{d}}$ is induced by the given action on $M_{\mathbb{R}}$ and the trivial action on $\Bbbk[P]$ as follows: writing $f_{\mathfrak{d}} = 1 + \sum a_m z^{\tilde{\varphi}_\tau(m)}$ as above, $f_{T(\mathfrak{d})} = 1 + \sum a_m z^{\tilde{\varphi}_{T(\tau)}(T(m))}$.

Lemma 4.6. *Let $\tilde{Q} \in \tilde{B}_0 \setminus \mathrm{Supp}(\tilde{\mathfrak{D}})$ be a point contained in the interior of a maximal cone $\sigma \in \tilde{\Sigma}$. We consider broken lines γ on $\tilde{B}_0$ for $T^N(\tilde{q})$, some $N \in \mathbb{Z}$, with endpoint $\tilde{Q}$, such that* $\mathrm{Mono}(\gamma) \notin I \cdot \Bbbk[P_{\tilde{\varphi}_\sigma}]$. *Let $k \geq 0$ be such that $(k+1)J \subset I$.*

1. *The number of bends of γ is at most k.*
2. *The number of broken lines for $T^N(\tilde{q})$ with endpoint $\tilde{Q}$ is $O(|N|^k)$.*
3. *For γ a broken line from $T^N(\tilde{q})$ with endpoint $\tilde{Q}$ write* $\mathrm{Mono}(\gamma) = a_\gamma z^{\tilde{\varphi}_\sigma(m_\gamma)}$ *where a_γ is a monomial in $\Bbbk[P]$. Assume that $|z^p| < \epsilon < 1$ for all $p \in P$. Then*
$$|a_\gamma| = O\Big(\prod_{\substack{\rho\in\Sigma\\ \dim\rho=1}} |z^{[D_\rho]}|^{|N|}\Big).$$

Proof. (1) At a bend $t_i \in (-\infty, 0]$ of γ the attached monomial $c_i z^{q_i}$ is replaced by the monomial $c_{i+1}z^{q_{i+1}} = cz^q \cdot c_i z^{q_i}$ where cz^q is a term in a positive power of the scattering function $f_{\mathfrak{d}}$ associated to the ray $\mathfrak{d}$ containing $\gamma(t_i)$. In particular $cz^q \in J \cdot \Bbbk[P_{\tilde{\varphi}_{\tau_{\mathfrak{d}}}}]$. Since $\mathrm{Mono}(\gamma) \notin I \cdot \Bbbk[P_{\tilde{\varphi}_\sigma}]$ and $(k+1)J \subset I$ it follows that there are at most k bends.

(2) Such a broken line crosses $O(|N|)$ scattering rays. If γ is a broken line for $T^N(\tilde{q})$ then the initial attached monomial is specified, equal to $z^{\tilde{\varphi}(T^N(\tilde{q}))}$. At a scattering ray $\mathfrak{d}$, let u denote the primitive generator of $\mathfrak{d}$, $f = f_{\mathfrak{d}}$ the attached function, and let cz^q be the monomial attached to the incoming segment of the broken line. Then the possible continuations of the broken line past $\mathfrak{d}$ correspond to the monomial terms in f^d, where $d = |r(q) \wedge u|$ is the index of the sublattice of M generated by $r(q)$ and u. Note that since $f \equiv 1 \mod J \cdot \Bbbk[P_{\tilde{\varphi}_{\tau_{\mathfrak{d}}}}]$ the number of monomial terms in f^d not lying in $I \cdot \Bbbk[P_{\tilde{\varphi}_{\tau_{\mathfrak{d}}}}]$ is bounded independent of d. Further, since there are a finite number of Γ-orbits of scattering rays, and the Γ-action preserves monomials, there is a bound on the number of monomial terms independent of the ray $\mathfrak{d}$. Now by (1) the total number of broken lines is $O(|N|^k)$.

(3) By symmetry we may assume that $N \geq 0$. Let $\mathfrak{d} \in \tilde{\mathfrak{D}}$ be a scattering ray, $f = f_{\mathfrak{d}}$ the attached function, and γ a broken line that crosses $\mathfrak{d}$. Suppose first that $\mathfrak{d}$ is contained in the interior of a maximal cone σ of $\tilde{\Sigma}$. Let $az^{\tilde{\varphi}_\sigma(m)}$ be the monomial attached to the incoming segment of γ near $\mathfrak{d}$. Let u be the primitive generator of $\mathfrak{d}$. Then the outgoing monomial $a'z^{\tilde{\varphi}_\sigma(m')}$ is obtained from the incoming monomial by multiplication by a monomial term in f^d, where $d = |m \wedge u|$. Write $f = 1 + f_1 + \cdots + f_r$, a sum of monomials. Since $f \equiv 1 \mod J \cdot \Bbbk[P_{\varphi_\sigma}]$ we have

$$f^d \equiv \sum_{i_1+\cdots+i_r \leq k} \binom{d}{i_1, \ldots, i_r} f_1^{i_1} \cdots f_r^{i_r} \mod I \cdot \Bbbk[P_{\varphi_\sigma}]. \tag{4.4}$$

The multinomial coefficient

$$\binom{d}{i_1, \ldots, i_r} := \frac{d!}{i_1! \cdots i_r!(d - i_1 - \cdots - i_r)!}$$

is bounded by d^k. The direction of the scattering ray $\mathfrak{d}$ is $u = T^s(\beta)$ where $0 \leq s \leq N$ and $\beta \in M$ is chosen from a finite set. The vector $m \in M$ is of the form

$$m = T^N(\tilde{q}) - \sum_{i=1}^{l} T^{s_i} \alpha_i$$

where $l \leq k$, $0 \leq s_i \leq N$ for each i, and the $\alpha_i \in M$ are chosen from a finite set. Indeed, for the monomial terms cz^q occurring in the powers f^d of the function $f = f_{\mathfrak{d}}$ attached to a given scattering ray $\mathfrak{d}$, only finitely many exponents $q \in P_{\tilde{\varphi}_{\tau_{\mathfrak{d}}}}$ occur (working modulo $I \cdot \Bbbk[P_{\tilde{\varphi}_{\tau_{\mathfrak{d}}}}]$). So there are only finitely many possible changes of exponent q for the attached monomial cz^q of a broken line at a scattering ray modulo the action of Γ. Now identify $M = \mathbb{Z}^2$ and let $\|\cdot\|$ denote the standard norm on $M_{\mathbb{R}} = \mathbb{R}^2$. Then

$$d = |m \wedge u| \leq \|m\| \cdot \|u\| = O(\lambda^{2N}).$$

So, the coefficient $a' \in \Bbbk[P]$ of the outgoing monomial is given by $a' = c \cdot z^p \cdot a$ where $c \in \Bbbk$, $p \in P$, and $c = O(\lambda^{2kN})$. Thus $|a'| = O(\lambda^{2kN}) \cdot |a|$ for $|z^p| < 1$.

Second, let $\rho \in \tilde{\Sigma}$ be a ray and σ_+, σ_- the maximal cones containing ρ. Suppose γ is a broken line that crosses ρ, travelling from σ_- to σ_+. Let $a_- z^{\tilde{\varphi}_{\sigma_-}(m)}$ be the monomial attached to the incoming segment of γ near ρ and $a_+ z^{\tilde{\varphi}_{\sigma_+}(m')}$ the monomial attached to the outgoing segment. By the definition of $\tilde{\varphi}$,

$$z^{\tilde{\varphi}_{\sigma_-}(m)} = \left(z^{[D_\rho]}\right)^{-\langle n_\rho, m\rangle} z^{\tilde{\varphi}_{\sigma_+}(m)},$$

where $n_\rho \in N$ is primitive, annihilates ρ, and is positive on σ_+. Write $d := -\langle n_\rho, m\rangle$; note $d = |u \wedge m| > 0$ where $u \in M$ is the primitive generator of ρ. If γ does not bend at ρ then $a_+ = (z^{[D_\rho]})^d \cdot a_-$. In general $a_+ = (z^{[D_\rho]})^d \cdot a'_-$ where $a'_- z^{\tilde{\varphi}_{\sigma_-}(m')}$ is obtained from $a_- z^{\tilde{\varphi}_{\sigma_-}(m)}$ as above (by applying the scattering automorphism associated to ρ and selecting a monomial term).

We need to show the exponent $d = |u \wedge m| > 0$ of $z^{[D_\rho]}$ in the previous paragraph is large for some rays ρ. This will allow us to absorb the $O(\lambda^{2N})$ factors coming from bends of γ and obtain the estimate (3). Let γ be a broken line for $T^N(\tilde{q})$, then

$$m_\gamma = T^N(\tilde{q}) - \sum_{i=1}^{l} T^{s_i}\alpha_i$$

as above, where $0 \leq l \leq k$, $N \geq s_1 \geq \cdots \geq s_l \geq 0$, and the α_i lie in a finite set. Write $s_0 = N$ and $s_{l+1} = 0$. Choose j such that $s_j - s_{j+1} \geq N/(k+1)$. Now consider the exponent $d = |u \wedge m|$ for $m = T^N(\tilde{q}) - \sum_{i=1}^{j} T^{s_i}\alpha_i$ given by the monomial attached to the segment of the broken line between bends j and $j+1$ and $u = T^{s_{j+1}}\beta$ the primitive generator of a ray of $\tilde{\Sigma}$ near bend $j+1$. Then $|u \wedge m| = |\beta \wedge T^{s_j - s_{j+1}} m'|$ where $m' = T^{-s_j}(m)$. Write $m' = \mu'_1 w_1 + \mu'_2 w_2$, then $|\mu'_1|$ is bounded and μ'_2 is bounded away from zero by Lemma 4.7. Now

$$u \wedge m = \beta \wedge T^{s_j - s_{j+1}} m' = \mu'_1(\beta \wedge w_1)\lambda^{-(s_j - s_{j+1})} + \mu'_2(\beta \wedge w_2)\lambda^{s_j - s_{j+1}},$$

where $s_j - s_{j+1} > N/(k+1)$, so

$$|u \wedge m| > c \cdot \lambda^{N/(k+1)}$$

for some constant $c > 0$.

Combining our results now gives the estimate

$$a_\gamma = O((\lambda^{2kN})^k \cdot \prod_{\substack{\rho \in \Sigma \\ \dim \rho = 1}} |z^{[D_\rho]}|^{c \cdot \lambda^{N/(k+1)}}).$$

where the first factor bounds the contribution associated to bends of γ and the second factor bounds the contribution associated to rays ρ of the fan crossed by γ, as described in the preceding two paragraphs. This implies the estimate (3) in the statement. Indeed, the above expression is of the form

$$\lambda^{aN} \cdot x^{c \cdot \lambda^{bN}}$$

where $a, b, c > 0$ and $\lambda > 1$ are constants, and $x = \prod |z^{[D_\rho]}|$. This is bounded by Cx^N for $0 \leq x < \epsilon < 1$, for some constant C (depending on ϵ). □

Lemma 4.7. *Let $A \subset \mathbb{R}$ be a finite set and $\lambda \in \mathbb{R}$, $\lambda > 1$. For $k \in \mathbb{N}$ let $S_k \subset \mathbb{R}$ be the set of real numbers s of the form $s = \sum_{i=1}^{l} c_i \lambda^{n_i}$ where $l \leq k$ and $c_i \in A$, $n_i \in \mathbb{Z}_{\geq 0}$ for each i. Then S_k is discrete for each k.*

Proof. Proof by induction on k. We have $S_0 = \{0\}$. Suppose S_k is discrete. We have $S_{k+1} = \bigcup_{n \geq 0} \lambda^n (S_k + A)$. Since $\lambda > 1$ we deduce that S_{k+1} discrete. □

For Propositions 4.8 and 4.9 below, the assertions hold after possibly replacing X_J by a smaller neighbourhood of $s(0) \in X_J$ (independent of I, Q and q).

Proposition 4.8. *Let $Q \in B_0 \setminus \mathrm{Supp}(\mathfrak{D})$ be a point contained in the interior of a maximal cone σ of Σ. Each term of the formal sum $\mathrm{Lift}_Q(q) = \sum \mathrm{Mono}(\gamma)$ is an analytic function on $V_{\sigma,\sigma,I}$ and the sum defines an analytic function on $V_{\sigma,\sigma,I}$.*

Proof. Recall that $V_{\sigma,\sigma,I}$ is an infinitesimal thickening of the reduced complex analytic space $V_{\sigma,\sigma,J}$. Write

$$V_{\sigma,\sigma} := \{(X_1, X_2) \mid |X_1| < R|X_2|, |X_2| < R|X_1|\} \subset (\mathbb{G}_m)^2_{X_1,X_2} \times S.$$

Then $V_{\sigma,\sigma}$ is a reduced complex analytic space containing $V_{\sigma,\sigma,I}$ as a locally closed subspace. We show that the sum $\mathrm{Lift}_Q(q)$ converges (uniformly on compact sets) to an analytic function on a neighbourhood of $V_{\sigma,\sigma,I}$ in $V_{\sigma,\sigma}$.

Let $\tilde{Q} \in \tilde{B}_0$ be a lift of Q and $\tilde{\sigma}$ the lift of σ containing $\tilde{Q}$. Let u_1, u_2 be the primitive generators of $\tilde{\sigma}$ (a basis of M) such that the orientation of u_1, u_2 agrees with that of w_1, w_2. Let $X_i = z^{\tilde{\varphi}_{\tilde{\sigma}}(u_i)}$, $i = 1, 2$, be the associated coordinate functions on $V_{\sigma,\sigma,I}$, so that

$$V_{\sigma,\sigma,I} \subset \{(X_1, X_2) \mid |X_1| < R|X_2|, |X_2| < R|X_1|\} \subset (\mathbb{G}_m)^2_{X_1,X_2} \times S'_I.$$

For $m \in M$, writing $m = \alpha_1 u_1 + \alpha_2 u_2$, we have $z^{\tilde{\varphi}_{\tilde{\sigma}}(m)} = X_1^{\alpha_1} X_2^{\alpha_2}$.

As already noted, broken lines γ on B_0 for q with endpoint Q lift uniquely to broken lines on $\tilde{B}_0$ for $T^N(\tilde{q})$ with endpoint $\tilde{Q}$, for some $N \in \mathbb{Z}$, and the attached monomials are identified. Write $\mathrm{Mono}(\gamma) = a_\gamma z^{\tilde{\varphi}_{\tilde{\sigma}}(m_\gamma)}$ and $m_\gamma = \alpha_1 u_1 + \alpha_2 u_2$. Clearly $\mathrm{Mono}(\gamma) = a_\gamma X_1^{\alpha_1} X_2^{\alpha_2} \in \Bbbk[P][X_1^{\pm 1}, X_2^{\pm 1}]$ is an analytic function on $V_{\sigma,\sigma}$. Also write $m_\gamma = \mu_1 w_1 + \mu_2 w_2$. By Lemma 4.10, (1), μ_1 and μ_2 are bounded below. The points u_1 and u_2 are adjacent integral points on the boundary of the infinite convex polytope Ξ with asymptotic directions w_1, w_2. It follows that $w_1 = \beta_1 u_2 + \gamma_1(u_1 - u_2)$ and $w_2 = \beta_2 u_1 + \gamma_2(u_2 - u_1)$, for some $\beta_1, \beta_2, \gamma_1, \gamma_2 > 0$. Hence

$$|z^{\tilde{\varphi}_{\tilde{\sigma}}(m_\gamma)}| = |X_1^{\alpha_1} X_2^{\alpha_2}| = (|X_2|^{\beta_1} |X_1/X_2|^{\gamma_1})^{\mu_1} (|X_1|^{\beta_2} |X_2/X_1|^{\gamma_2})^{\mu_2}.$$

Now $|X_1/X_2| < R$, $|X_2/X_1| < R$ on $V_{\sigma,\sigma}$. Thus, for $0 < \delta < \min(R^{-\gamma_1/\beta_1}, R^{-\gamma_2/\beta_2})$ and any $\delta' > 0$, $|z^{\tilde{\varphi}_{\tilde{\sigma}}(m_\gamma)}|$ is bounded for $0 < \delta' < |X_1|, |X_2| < \delta$. (Note that δ' is required because μ_1, μ_2 may be negative — we only obtain uniform convergence of the series $\mathrm{Lift}_Q(q)$ on compact subsets of $V_{\sigma,\sigma}$.) By Lemma 4.6, (3), if $|z^p| < \epsilon < 1$ for all $p \in P$ we have $|a_\gamma| = O(\epsilon^{|N|})$. By Lemma 4.6, (2), the number of broken lines for $T^N(\tilde{q})$ is $O(|N|^k)$. Combining, we deduce that $\mathrm{Lift}_q(Q) = \sum \mathrm{Mono}(\gamma)$ is convergent on the open analytic subset $V'_{\sigma,\sigma}$ of $V_{\sigma,\sigma}$ defined by $|X_1|, |X_2| < \delta$ and $|z^p| < 1$ for all $p \in P$, for some $\delta > 0$ (independent of I and q). After replacing X_J by an analytic neighbourhood of the vertex $s(0) \in X_J$, we may assume that $V_{\sigma,\sigma,I} \subset V'_{\sigma,\sigma}$. $\square$

Proposition 4.9. *Let* $Q \in B_0 \setminus \operatorname{Supp}(\mathfrak{D})$ *be a point contained in the interior of a maximal cone* σ *of* Σ *and let* ρ *be an edge of* σ*. Consider the formal sum* $\operatorname{Lift}_Q(q) = \sum \operatorname{Mono}(\gamma)$*. For* Q *sufficiently close to* ρ *each term of the sum is an analytic function on* $V_{\rho,I}$ *and the sum defines an analytic function on* $V_{\rho,I}$*.*

Proof. Write $\sigma_+ = \sigma$ and let $\sigma_- \in \Sigma$ be the other maximal cone containing ρ. Let $\tilde{Q}, \tilde{\rho}, \tilde{\sigma}_+, \tilde{\sigma}_-$ be compatible lifts to $\tilde{B}_0$. Let u, u_-, u_+ be the primitive generators of $\tilde{\rho}$ and the remaining edges of $\tilde{\sigma}_-$ and $\tilde{\sigma}_+$, and write X, X_-, X_+ for the corresponding coordinates on $V_{\rho,I}$. So

$$V_{\rho,I} \subset \{(X_-, X_+, X) \,|\, |X_-| < R|X|,\ |X_+| < R|X|\} \subset V(X_-X_+ - z^{[D_\rho]}X^{-D_\rho^2}f_\rho) \subset \mathbb{A}^2_{X_-,X_+} \times (\mathbb{G}_m)_X \times S'_I.$$

Define

$$V_\rho = \{(X_-, X_+, X) \,|\, |X_-| < R|X|,\ |X_+| < R|X|\} \subset V(X_-X_+ - z^{[D_\rho]}X^{-D_\rho^2}f_\rho) \subset \mathbb{A}^2_{X_-,X_+} \times (\mathbb{G}_m)_X \times S.$$

We assume that the orientation of u_-, u_+ is the same as that of w_1, w_2.

We first consider broken lines γ lying in the cone generated by u and w_2. Write $\operatorname{Mono}(\gamma) = a_\gamma z^{\tilde{\varphi}_{\tilde{\sigma}}(m_\gamma)}$, and $m_\gamma = \alpha u + \alpha_+ u_+ = \mu_1 w_1 + \mu_2 w_2$. By Lemma 4.10,(1), $|\mu_1|$ is bounded, and $\mu_2 > 0$ for all but finitely many γ. By Lemma 4.10,(2), $\alpha_+ \geq 0$, so $\operatorname{Mono}(\gamma) = a_\gamma X^\alpha X_+^{\alpha_+} \in \Bbbk[P][X^{\pm 1}, X_-, X_+]$ is analytic on V_ρ for each γ. In particular we may assume in what follows (discarding finitely many terms $\operatorname{Mono}(\gamma)$) that $\mu_2 > 0$. Writing $w_1 = -\beta_1 u_+ + \gamma_1 u$ and $w_2 = \beta_2 u + \gamma_2(u_+ - u)$, we have $\beta_1, \beta_2, \gamma_1, \gamma_2 > 0$ and

$$|z^{\tilde{\varphi}_{\tilde{\sigma}}(m_\gamma)}| = (|X_+|^{-\beta_1}|X|^{\gamma_1})^{\mu_1}(|X|^{\beta_2}|X_+/X|^{\gamma_2})^{\mu_2}.$$

Recall that $|X_+/X|, |X_-/X| < R$ on V_ρ. Also, the equation for V_ρ gives

$$|z^{[D_\rho]}X_+^{-1}| = |X_-| \cdot |X|^{D_\rho^2} \cdot |f_\rho|^{-1}.$$

The function f_ρ on V_ρ restricts to the constant function 1 over S_J. Hence we may impose the condition $|f_\rho| > \delta'$ for any small $\delta' > 0$. We see that for $0 < \delta < R^{-\gamma_2/\beta_2}$, and any $\delta' > 0$, if $\delta' < |X| < \delta$ and $|f_\rho| > \delta'$ then

$$|z^{\tilde{\varphi}_{\tilde{\sigma}}(m_\gamma)}| \cdot |z^{[D_\rho]}|^c$$

is bounded, where $c = \beta_1 \cdot \sup(\{\mu_1\})$ is a constant. Now by Lemma 4.6, (3), if $|z^p| < \epsilon < 1$ for all $p \in P$ then

$$|\operatorname{Mono}(\gamma)| = |a_\gamma z^{\tilde{\varphi}_{\tilde{\sigma}}(m_\gamma)}| = O(\epsilon^{|N|}).$$

Recall that the number of broken lines for $T^N\tilde{q}$ is $O(|N|^k)$ (Lemma 4.6, (2)). We deduce that the sum $\sum \operatorname{Mono}(\gamma)$ over broken lines γ lying in $\langle u, w_2\rangle_{\mathbb{R}_{\geq 0}}$ is uniformly convergent on compact sets for $|X| < \delta$, $f_\rho \neq 0$, and $|z^p| < 1$ for all $p \in P$, where $\delta > 0$ is independent of I and q.

Finally, we consider broken lines lying in $\langle u, w_1 \rangle_{\mathbb{R}_{\geq 0}}$. We write

$$\mathrm{Lift}_Q(q) = \mathrm{Lift}^+_Q(q) + \mathrm{Lift}^0_Q(q) + \mathrm{Lift}^-_Q(q)$$

where the terms are sums over broken lines γ with $\langle n_\rho, m_\gamma \rangle$ positive, zero, and negative respectively, where as before n_ρ is positive on σ_+. Then $\mathrm{Lift}^+_Q(q) + \mathrm{Lift}^0_Q(q)$ is the sum over broken lines lying in $\langle u, w_2 \rangle_{\mathbb{R}_{\geq 0}}$ and so defines an analytic function on $V_{\rho,I}$ as proved above. Let Q' be a point close to Q contained in σ_-. Then

$$\mathrm{Lift}^-_Q(q) = \theta(\mathrm{Lift}^-_{Q'}(q))$$

where θ is the scattering automorphism attached to ρ, see the proof of (3.8). Because the scattering automorphism is incorporated into the definition of $V_{\rho,I}$, this means that $\mathrm{Lift}^-_Q(q)$ and $\mathrm{Lift}^-_{Q'}(q)$ are identified as formal sums of functions on $V_{\rho,I}$. Now, $\mathrm{Lift}^-_{Q'}(q)$ is a sum over broken lines on $\tilde{B}_0$ lying in $\langle u, w_1 \rangle_{\mathbb{R}_{\geq 0}}$. Hence by symmetry $\mathrm{Lift}^-_{Q'}(q)$ defines an analytic function on $V_{\rho,I}$. Combining we deduce that $\mathrm{Lift}_Q(q)$ defines an analytic function on $V_{\rho,I}$. □

Lemma 4.10. *Let $\tilde{Q} \in \tilde{B}_0 \setminus \mathrm{Supp}(\tilde{\mathfrak{D}})$ be a point contained in the interior of a maximal cone $\tilde{\sigma} \in \tilde{\Sigma}$. Consider broken lines γ on $\tilde{B}_0$ for $T^N(\tilde{q})$ with endpoint $\tilde{Q}$, for all $N \in \mathbb{Z}$ such that $T^N(\tilde{q}) \in \langle \tilde{Q}, w_2 \rangle_{\mathbb{R}_{\geq 0}}$. Write $\mathrm{Mono}(\gamma) = a_\gamma z^{\tilde{\varphi}_{\tilde{\sigma}}(m_\gamma)}$, and $m_\gamma = \mu_1 w_1 + \mu_2 w_2$.*

1. *$|\mu_1|$ is bounded, and μ_2 is positive for all but finitely many γ. In particular, μ_1, μ_2 are bounded below.*
2. *Let u_1, u_2 be generators of $\tilde{\sigma}$ with the same orientation as w_1, w_2. Then for $\tilde{Q}$ sufficiently close to $\tilde{\rho} := \mathbb{R}_{\geq 0} \cdot u_1$, m_γ lies in the half space $\mathbb{R} \cdot u_1 + \mathbb{R}_{\geq 0} \cdot u_2$ for each γ.*

Proof. (1) Note that the rays spanned by w_1, w_2 are irrational so $\mu_1, \mu_2 \neq 0$. Suppose for a contradiction that there is an infinite sequence of broken lines γ such that $m_\gamma = \mu_1 w_1 + \mu_2 w_2$ with $\mu_2 < 0$. Each broken line has at most k bends and there are a finite number of Γ-orbits of possible changes $\alpha \in M$ of the derivative of γ at a bend; see Lemma 4.6 and its proof. So, passing to a subsequence, we may assume that the bends (in order of increasing $t \in (-\infty, 0]$) of each γ are of types $T^{s_1}\alpha_1, \ldots, T^{s_l}\alpha_l$ for some fixed $\alpha_1, \ldots, \alpha_l \in M$, $l \leq k$, and $N \geq s_1 \geq s_2 \geq \cdots \geq s_l \geq 0$ (depending on γ). Say $N - s_i$ is bounded for $i \leq l'$ and unbounded otherwise. Passing to a subsequence, we may assume that $N - s_i$ is constant for $i \leq l'$. Let γ' be the broken line obtained by truncating γ after the first l' bends, and moving by a homothety so that (extending its final line segment) γ' ends at $\tilde{Q}$. Then $m_{\gamma'} = T^N(m)$ for some fixed $m \in M$. Moreover, since $Tw_2 = \lambda w_2$, $\lambda > 1$, it follows that we must have $m = \nu_1 w_1 + \nu_2 w_2$ with $\nu_2 < 0$. This is a contradiction because $m_{\gamma'}$ lies in the half-space $\mathbb{R} \cdot \tilde{Q} + \mathbb{R}_{\geq 0} \cdot w_2$.

To see that $|\mu_1|$ is bounded, recall that $m_\gamma = T^N(\tilde{q}) - \sum_{i=1}^{l} T^{s_i}\alpha_i$ where $0 \leq l \leq k$, $0 \leq s_i \leq N$, and the α_i are selected from a finite set. Now since $Tw_1 = \lambda^{-1} w_1$ it follows that $|\mu_1|$ is bounded.

(2) Let $\mathfrak{u}$ be the connected component of $\tilde{B}_0 \backslash \mathrm{Supp}_I(\tilde{\mathfrak{D}})$ contained in $\tilde{\sigma}$ and containing $\tilde{\rho}$ in its closure. Let $\tilde{Q}' \in \mathfrak{u}$ be a point such that $\tilde{Q}' \in \langle \tilde{\rho}, \tilde{Q} \rangle_{\mathbb{R}_{\geq 0}}$. Then if $T^N(\tilde{q}) \in \langle \tilde{Q}, w_2 \rangle_{\mathbb{R}_{\geq 0}}$ and γ' is a broken line for $T^N(\tilde{q})$ with endpoint $\tilde{Q}'$, we obtain a broken line γ for $T^N(\tilde{q})$ with endpoint $\tilde{Q}$ and $m_\gamma = m_{\gamma'}$ as follows: First apply a homothety to obtain a broken line passing through $\tilde{Q}$, then truncate at $\tilde{Q}$.

Now suppose γ is a broken line for $T^N(\tilde{q})$ with endpoint $\tilde{Q}$ and m_γ not lying in the half-space $\mathbb{R} \cdot u_1 + \mathbb{R}_{\geq 0} \cdot u_2$. Since m_γ lies in the half-space $\mathbb{R} \cdot \tilde{Q} + \mathbb{R}_{\geq 0} \cdot w_2$ we find m_γ lies in the cone generated by $-\tilde{Q}, -u_1$. In particular, m_γ does not lie in the half-space $\mathbb{R} \cdot w_1 + \mathbb{R}_{\geq 0} w_2$, so by (1) there are only finitely many such γ. Now by the above construction it follows that for $\tilde{Q}$ sufficiently close to $\tilde{\rho}$ there are none. □

4.3. **Smoothness.**

Assumptions 4.11. *We consider the following three cases:*

1. *Every $\mathbb{A}^1$-class lies in J, and for any monomial ideal I such that $\sqrt{I} = J$ there are only a finite number of $\mathbb{A}^1$-classes not lying in I. Furthermore there are at least 2 rays ρ of Σ such that $z^{[D_\rho]} \in J$.*
2. *There is a toric model $p \colon Y \to \bar{Y}$. For some ample divisor $\bar{H}$ on $\bar{Y}$, $\sigma_P \cap (p^*\bar{H})^\perp$ is a face of σ_P, and $J = P \setminus P \cap (p^*\bar{H})^\perp$.*
3. *The field $\Bbbk = \mathbb{C}$. The intersection matrix $(D_i \cdot D_j)_{1 \leq i,j \leq n}$ is negative definite, $D_i^2 \leq -2$ for each i, and $n \geq 3$. The cone σ_P is strictly convex. For some nef divisor L on Y such that $L^\perp \cap \mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}} = \langle D_1, \ldots, D_n \rangle_{\mathbb{R}_{\geq 0}}$, $\sigma_P \cap L^\perp$ is a face of σ_P and $J = P \setminus P \cap L^\perp$.*

Let I be a monomial ideal such that $\sqrt{I} = J$. In cases (1) and (2) we have an algebraic flat family $f_I \colon X_I \to S_I$ by Theorem 2.33, (1), consistency of $\mathfrak{D}^{\mathrm{can}}$, and Remark 3.19. In case (3) we have an analytic flat family $f_I \colon X_I \to S_I'$ by Theorem 4.5, where S_I' is an analytic neighbourhood of $0 \in S_I$. In the remainder of this section we will abuse notation and denote S_I' by S_I.

Let $\mathfrak{f}_J \colon \mathfrak{X}_J \to \mathfrak{S}_J$ denote the formal deformation determined by the deformations $X_{J^{N+1}} \to S_{J^{N+1}}$ for $N \geq 0$. Thus in the algebraic cases $\mathfrak{S}_J = \mathrm{Spf}(\varprojlim \Bbbk[P]/J^{N+1})$ is the formal spectrum of the J-adic completion of $\Bbbk[P]$, $\mathfrak{X}_J$ is a formal scheme, and $\mathfrak{X}_J \to \mathfrak{S}_J$ is an adic flat morphism of formal schemes. In the analytic case, $\mathfrak{S}_J$ is the formal complex analytic space obtained as the completion of S along S_J, $\mathfrak{X}_J$ is a formal complex analytic space, and $\mathfrak{X}_J \to \mathfrak{S}_J$ is an adic flat morphism. We refer to [G60] and [B78] for background on formal schemes and formal complex analytic spaces.

Let $Z_I := \mathrm{Sing}(f_I) \subset X_I$ denote the singular locus of $f_I \colon X_I \to S_I$. Thus $Z_I \subset X_I$ is a closed embedding of schemes or complex analytic spaces. Since the singular locus is compatible with base-change, the singular loci $Z_{J^n} \subset X_{J^n}$ determine a closed embedding $\mathfrak{Z}_J \subset \mathfrak{X}_J$ which we refer to as the singular locus of $\mathfrak{f}_J \colon \mathfrak{X}_J \to \mathfrak{S}_J$.

We shall prove the following main smoothing theorems in this section:

Theorem 4.12. *Assume σ_P is strictly convex and $J = \mathfrak{m} := P \setminus \{0\}$ satisfies the conditions 4.11, (1). Then the map $\mathcal{O}_{\mathfrak{S}_J} \to \mathfrak{f}_{J*}\mathcal{O}_{\mathfrak{Z}_J}$ is not injective. In particular, for $I = \mathfrak{m}^{N+1}$ and $N \gg 0$, the map $\Bbbk[P]/I \to f_{I*}\mathcal{O}_{\mathrm{Sing}(f_I)}$ is not injective.*

Theorem 4.13. *(Looijenga's conjecture) Suppose that $\Bbbk = \mathbb{C}$ and the intersection matrix $(D_i \cdot D_j)_{1 \le i,j \le n}$ is negative definite, so that $D \subset Y$ can be contracted to a cusp singularity $q \in Y'$. Then the dual cusp to $q \in Y'$ is smoothable.*

We have a section $s \colon S_J \to X_J$ such that, for $t \in S_J$ general, the point $s(t) \in X_{J,t}$ on the fibre is the vertex in the algebraic case and the cusp in the analytic case. We write $X_J^o := X_J \setminus s(S_J) \subset X_J$ and $X_I^o \subset X_I$, $\mathfrak{X}_J^o \subset \mathfrak{X}_J$ for the induced open embeddings.

Lemma 4.14. *There exists $0 \neq g \in \Bbbk[P]$ such that $\mathrm{Supp}(g \cdot \mathcal{O}_{\mathfrak{Z}_J})$ is contained in $s(S_J)$. In particular, $\mathfrak{f}_{J*}(g \cdot \mathcal{O}_{\mathfrak{Z}_J})$ is a coherent sheaf on $\mathfrak{S}_J$.*

Proof. We first recall explicit open coverings of $\mathfrak{X}_J^o$. In the algebraic cases (1) and (2), $\mathfrak{X}_J^o$ is a union of open subschemes $\mathfrak{U}_{i,J}$, $i = 1, \ldots, n$, given as follows. Write $a_i = z^{[D_i]}$ and $m_i = -D_i^2$. In case (1),

$$\mathfrak{U}_{i,J} = V(x_{i-1}x_{i+1} - a_i x_i^{m_i}) \subset \mathbb{A}^2_{x_{i-1},x_{i+1}} \times (\mathbb{G}_m)_{x_i} \times \mathfrak{S}_J. \tag{4.5}$$

In case (2), let E_{ij} be the exceptional curves of $p \colon (Y, D) \to (\bar{Y}, \bar{D})$ over points of $\bar{D}_i$, and write $b_{ij} = z^{[E_{ij}]}$. Then

$$\mathfrak{U}_{i,J} = V(x_{i-1}x_{i+1} - a_i x_i^{m_i} \prod (1 + b_{ij} x_i^{-1})) \subset \mathbb{A}^2_{x_{i-1},x_{i+1}} \times (\mathbb{G}_m)_{x_i} \times \mathfrak{S}_J.$$

In the analytic case (3), let $\mathfrak{U}_{i,J}$ be defined as in (4.5) above. Then $\mathfrak{X}_J^o$ is a union of open subspaces $\mathfrak{V}_{i,J}$, $i = 1, \ldots, n$, such that $\mathfrak{V}_{i,J}$ is an analytic open subspace of $\mathfrak{U}_{i,J}$ for each i.

We now use the charts $\mathfrak{U}_{i,J}$ to compute the singular locus explicitly. In case (1) the singular locus $\mathfrak{Z}_{i,J}$ of $\mathfrak{U}_{i,J}/\mathfrak{S}_J$ is given by

$$\mathfrak{Z}_{i,J} = V(x_{i-1}, x_{i+1}, a_i) \subset \mathfrak{U}_{i,J}.$$

Hence if we define $g = a_1 \cdots a_n$ then $\mathrm{Supp}(g \cdot \mathcal{O}_{\mathfrak{Z}_J})$ is contained in $s(S_J)$. The same is true in the analytic case (3).

Similarly, in case (2) the structure sheaf of the singular locus of $\mathfrak{U}_{i,J}$ is annihilated by $g_i := a_i \prod_{j \neq k}(b_{ij} - b_{ik})$. (Here $\prod_{j \neq k}(b_{ij} - b_{ik})$ is the discriminant of the polynomial $f(x_i) := \prod(x_i + b_{ij})$. It is a linear combination of f and f' with coefficients in $\Bbbk[\{b_{ij}\}][x_i]$. See [L02], p. 200–204.) So we can take $g = g_1 \cdots g_n$.

The support of $g \cdot \mathcal{O}_{\mathfrak{Z}_J}$ is a closed subset of $s(S_J)$, hence proper over S_J. It follows that $\mathfrak{f}_{J*}(g \cdot \mathcal{O}_{\mathfrak{Z}_J})$ is coherent by [B78], 3.1 and [G61], 3.4.2. □

Let $u(J)$ denote the natural map

$$u(J)\colon \mathcal{O}_{\mathfrak{S}_J} \to \mathfrak{f}_{J*}(\mathcal{O}_{\mathfrak{Z}_J}).$$

Let $\mathfrak{m}$ denote the monomial ideal $P \setminus P^\times$ of P as usual. So, for any monomial ideal $J \subsetneq P$, we have $J \subset \mathfrak{m}$.

Lemma 4.15. *$u(J)$ is injective if and only if $u(\mathfrak{m})$ is injective.*

Proof. Let $0 \neq g \in \Bbbk[P]$ be the element given by Lemma 4.14. Let $\mathcal{K}_J$ be the kernel of $u(J)$ and $\mathcal{K}'_J$ the kernel of $g \cdot u(J)$. Thus $\mathcal{K}_J, \mathcal{K}'_J$ are ideal sheaves in $\mathcal{O}_{\mathfrak{S}_J}$ and $g \cdot \mathcal{K}'_J \subset \mathcal{K}_J \subset \mathcal{K}'_J$. The local rings of $\mathfrak{S}_J$ are domains by Lemma 4.16, so $\mathcal{K}_J = 0$ if and only if $\mathcal{K}'_J = 0$. The sheaf $\mathcal{K}'_J$ is coherent because the image of $g \cdot u(J)$ is contained in the coherent subsheaf $\mathfrak{f}_{J*}(g \cdot \mathcal{O}_{\mathfrak{Z}_J}) \subset \mathfrak{f}_{J*}\mathcal{O}_{\mathfrak{Z}_J}$.

We claim that the natural map

$$\mathcal{K}'_J \otimes_{\mathcal{O}_{\mathfrak{S}_J}} \mathcal{O}_{\mathfrak{S}_\mathfrak{m}} \to \mathcal{K}'_\mathfrak{m}$$

is an isomorphism. Let $z \in S_\mathfrak{m}$ be a point and let $\hat{\mathcal{O}}_{S,z}$ denote the completion of $\mathcal{O}_{S,z}$ at its maximal ideal. Note that $\hat{\mathcal{O}}_{S,z}$ coincides with the completions of $\mathcal{O}_{\mathfrak{S}_\mathfrak{m},z}$ and $\mathcal{O}_{\mathfrak{S}_J,z}$ at their maximal ideals. It suffices to show that the map

$$\mathcal{K}'_{J,z} \otimes_{\mathcal{O}_{\mathfrak{S}_J,z}} \hat{\mathcal{O}}_{S,z} \to \mathcal{K}'_{\mathfrak{m},z} \otimes_{\mathcal{O}_{\mathfrak{S}_\mathfrak{m},z}} \hat{\mathcal{O}}_{S,z}$$

is an isomorphism for each z (because $\mathcal{O}_{\mathfrak{S}_\mathfrak{m},z} \subset \hat{\mathcal{O}}_{S,z}$ is faithfully flat). We have an exact sequence of coherent sheaves

$$0 \to \mathcal{K}'_J \to \mathcal{O}_{\mathfrak{S}_J} \to \mathfrak{f}_{J*}(g \cdot \mathcal{O}_{\mathfrak{Z}_J})$$

and so an exact sequence of $\hat{\mathcal{O}}_{S,z}$-modules

$$0 \to \mathcal{K}'_{J,z} \otimes \hat{\mathcal{O}}_{S,z} \to \hat{\mathcal{O}}_{S,z} \to \mathfrak{f}_{J*}(g \cdot \mathcal{O}_{\mathfrak{Z}_J})_z \otimes \hat{\mathcal{O}}_{S,z}.$$

Now

$$\mathfrak{f}_{J*}(g \cdot \mathcal{O}_{\mathfrak{Z}_J})_z \otimes \hat{\mathcal{O}}_{S,z} = (g \cdot \mathcal{O}_{\mathfrak{Z}_J})_{s(z)} \otimes \hat{\mathcal{O}}_{S,z} = (\widehat{g \cdot \mathcal{O}_{\mathfrak{Z}_J}})_{s(z)} = g \cdot \hat{\mathcal{O}}_{\mathfrak{Z}_J,s(z)}$$

where the hats denote completion with respect to the maximal ideal of $\mathcal{O}_{\mathfrak{S}_J,z}$. Thus $\mathcal{K}'_{J,z} \otimes \hat{\mathcal{O}}_{S,z}$ is the kernel of the map

$$\hat{\mathcal{O}}_{S,z} \to g \cdot \hat{\mathcal{O}}_{\mathfrak{Z}_J,s(z)}.$$

By the base-change property for the singular locus, this map coincides with the corresponding map for $\mathfrak{m}$. This proves the claim.

The support of the ideal sheaf $\mathcal{K}'_J$ is either empty or S_J (because the local rings of $\mathfrak{S}_J$ are domains and S_J is connected). So $\mathcal{K}'_J = 0$ if and only if $\mathcal{K}'_{\mathfrak{m}} = 0$ by the claim. ☐

Lemma 4.16. *The local rings of* $\mathfrak{S}_J$ *are integral domains.*

Proof. The completion of the local ring of $\mathfrak{S}_J$ at a point $z \in S_J$ is identified with the completion of the local ring of the toric variety S at z. By Serre's criterion for normality, the completion of a normal Noetherian ring at a maximal ideal is a local normal Noetherian ring; in particular, it is a domain. Since $\mathcal{O}_{\mathfrak{S}_J,z}$ is a local Noetherian ring, it is contained in its completion. We deduce that $\mathcal{O}_{\mathfrak{S}_J,z}$ is a domain. ☐

Lemma 4.17. *Suppose we are in case (2) of Assumptions 4.11. Then* $u(J)$ *is not injective.*

Proof. By Lemma 4.14 there exists $0 \neq g \in \Bbbk[P]$ such that $\operatorname{Supp}(g \cdot \mathcal{O}_{\mathfrak{Z}_J}) \subset s(S_J)$. By Proposition 3.40 there exists $0 \neq h \in \Bbbk[P]$ such that $\operatorname{Supp}(h \cdot \mathcal{O}_{\mathfrak{Z}_J})$ is disjoint from $s(S_J)$. Thus $gh \cdot \mathcal{O}_{\mathfrak{Z}_J} = 0$, that is, gh lies in the kernel of $u(J)$. ☐

Proof of Theorem 4.12. By Proposition 1.19 there exists a toric blowup $\pi\colon (\tilde{Y}, \tilde{D}) \to (Y, D)$ such that $(\tilde{Y}, \tilde{D})$ has a toric model $p\colon (\tilde{Y}, \tilde{D}) \to (\bar{Y}, \bar{D})$.

Let $\bar{H}$ be an ample divisor on $\bar{Y}$. Let $\sigma_{\tilde{P}} \subset A_1(\tilde{Y}, \mathbb{R})$ be the rational polyhedral cone given by

$$\sigma_{\tilde{P}} := (\pi_*)^{-1}\sigma_P \cap \{\beta \in H_2(\tilde{Y}, \mathbb{R}) \mid \beta \cdot (p^*\bar{H}) \geq 0\}$$

and $\tilde{P} := \sigma_{\tilde{P}} \cap A_1(\tilde{Y}, \mathbb{Z})$ the associated toric monoid. Let $F \subset A_1(\tilde{Y}, \mathbb{Z})$ be the free abelian group generated by the π-exceptional divisors. Since σ_P is strictly convex there exists an ample divisor H on Y such that $\sigma_P \subset (H \geq 0)$ and $\sigma_P \cap H^\perp = \{0\}$. Then $\sigma_{\tilde{P}} \cap (\pi^*H)^\perp$ is a face of $\sigma_{\tilde{P}}$ contained in $F \otimes_{\mathbb{Z}} \mathbb{R}$. Define radical monomial ideals $\tilde{\mathfrak{m}}, \tilde{J}_1, \tilde{J}_2 \subset \tilde{P}$ by

$$\begin{aligned} \tilde{\mathfrak{m}} &= \tilde{P} \setminus \tilde{P}^\times \\ \tilde{J}_1 &= \tilde{P} \setminus (p^*\bar{H})^\perp \\ \tilde{J}_2 &= \tilde{P} \setminus (\pi^*H)^\perp \end{aligned}$$

By Lemma 4.17, $u(\tilde{J}_1)$ is not injective. Hence $u(\tilde{\mathfrak{m}})$ and $u(\tilde{J}_2)$ are not injective by Lemma 4.15. Let $\tilde{\mathfrak{X}}_{\tilde{J}_2}/\tilde{\mathfrak{S}}_{\tilde{J}_2}$ be the family given by Theorem 2.33, (1) with $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$ over the formal thickening of $\tilde{S}_{\tilde{J}_2} := \operatorname{Spec} \Bbbk[\tilde{P}]/\tilde{J}_2$.

Let $T = \operatorname{Hom}(F, \mathbb{G}_m)$ be the big torus for $\tilde{S}_{\tilde{J}_2}$. Write $\tilde{D} = \tilde{D}_1 + \cdots + \tilde{D}_r$. Let $T_{\tilde{Y}} = \operatorname{Hom}(A_1(\tilde{Y}), \mathbb{G}_m)$ be the big torus for $\tilde{S} = \operatorname{Spec} \Bbbk[\tilde{P}]$ and $T^{\tilde{D}} := \mathbb{G}_m^r \to T_{\tilde{Y}}$ the homomorphism of tori given by the map of character lattices

$$A_1(\tilde{Y}) \to \mathbb{Z}^r, \quad \beta \mapsto (\beta \cdot \tilde{D}_i)_{i=1}^r.$$

The family $\tilde{\mathfrak{X}}_{\tilde{J}_2}/\tilde{\mathfrak{S}}_{\tilde{J}_2}$ is $T^{\tilde{D}}$-equivariant, as discussed in §3.5. The composition $F \subset A_1(\tilde{Y}) \to \mathbb{Z}^r$ is a primitive embedding (because $F \subset \langle \tilde{D}_1, \ldots, \tilde{D}_r \rangle$ and its intersection matrix is unimodular, congruent to $-I$). So the corresponding composition $T^{\tilde{D}} \to T_{\tilde{Y}} \to T$ admits a splitting $T \to T^{\tilde{D}}$. By T-equivariance, the restriction of the family $\tilde{\mathfrak{X}}_{\tilde{J}_2}/\tilde{\mathfrak{S}}_{\tilde{J}_2}$ to the open subscheme of $\tilde{\mathfrak{S}}_{\tilde{J}_2}$ defined by $T \subset \tilde{S}_{\tilde{J}_2}$ is isomorphic to the direct product of $\mathfrak{X}_{\mathfrak{m}}/S_{\mathfrak{m}}$ with T. Thus $u(\mathfrak{m})$ is not injective because $u(\tilde{J}_2)$ is not injective. □

Proof of Theorem 4.13. Let $f\colon Y \to Y'$ be the contraction of $D \subset Y$. We may assume f is the minimal resolution of Y'. We may further assume $n \geq 3$. Indeed, the embedding dimension of the dual cusp equals $\max(n, 3)$ by [N80], Corollary 7.8, p. 232 and [KM98], Theorem 4.57, p. 143. So in particular for $n \leq 3$ the dual cusp is a hypersurface and thus smoothable. Let L be a nef divisor on Y such that $\mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}} \cap L^{\perp} = \langle D_1, \ldots, D_n \rangle_{\mathbb{R}_{\geq 0}}$. Let $\sigma_P \subset H_2(Y, \mathbb{R})$ be a strictly rational polyhedral cone containing $\mathrm{NE}(Y)$ such that $\sigma_P \cap L^{\perp}$ is a face of σ_P. Let $P = \sigma_P \cap H_2(Y, \mathbb{Z})$ and $J = P \setminus P \cap L^{\perp}$. By Lemma 4.15 and Theorem 4.12, $u(J)$ is not injective.

Let $P \in S_J$ be a point lying in the interior of the toric variety S_J and $h \in \mathcal{O}_{\mathfrak{S}_J, P}$ a nonzero element of the kernel of $u(J)$ near P. By Lemma 4.18 there is a morphism

$$v\colon \operatorname{Spec} \mathbb{C}[t]/(t^{N+1}) \to \mathfrak{S}_J$$

such that $0 \mapsto P$ and $0 \neq v^*(h) \in \mathbb{C}[t]/(t^{N+1})$. Let $\mathcal{Y}/\operatorname{Spec}(\mathbb{C}[t]/(t^{N+1}))$ be the pullback of $\mathfrak{X}_J/\mathfrak{S}_J$ by v and $\mathcal{Z} \subset \mathcal{Y}$ its singular locus. Then $\mathcal{Y}/\operatorname{Spec}(\mathbb{C}[t]/(t^{N+1}))$ is a deformation of the dual cusp singularity and $\mathcal{O}_{\mathcal{Z}}$ is annihilated by t^N. By [A76], Theorem 5.1, there is an algebraic finite type deformation $\mathcal{Y}'/\operatorname{Spec} \mathbb{C}[\![t]\!]$ whose restriction to $\operatorname{Spec}(\mathbb{C}[t]/(t^{N+1}))$ is locally analytically isomorphic to $\mathcal{Y}/\operatorname{Spec}(\mathbb{C}[t]/(t^{N+1}))$. Let $\mathcal{Z}' \subset \mathcal{Y}'$ denote the singular locus of $\mathcal{Y}'/\operatorname{Spec} \mathbb{C}[\![t]\!]$. Then $\mathcal{O}_{\mathcal{Z}'}$ is a finite $\mathbb{C}[\![t]\!]$-module because the fibre $\mathcal{Y}'_0$ has an isolated singularity (using [Ma89], Theorem 8.4, p. 58). Now $\mathcal{O}_{\mathcal{Z}} = \mathcal{O}_{\mathcal{Z}'}/t^{N+1}\mathcal{O}_{\mathcal{Z}'}$ and $t^N \mathcal{O}_{\mathcal{Z}} = 0$, so $t^N \mathcal{O}_{\mathcal{Z}'} = t^{N+1} \mathcal{O}_{\mathcal{Z}'}$ and thus $t^N \mathcal{O}_{\mathcal{Z}'} = 0$ by Nakayama's lemma. Hence the general fibre of $\mathcal{Y}'/\operatorname{Spec} \mathbb{C}[\![t]\!]$ is smooth, and $\mathcal{Y}'/\operatorname{Spec} \mathbb{C}[\![t]\!]$ is a smoothing of the dual cusp. □

Lemma 4.18. *Let A be the completion of a finitely generated normal Cohen-Macaulay $\mathbb{C}$-algebra at a maximal ideal. Let $0 \neq a \in A$. Then there exists $N \geq 0$ and a $\mathbb{C}$-algebra map $f : A \to \mathbb{C}[t]/(t^{N+1})$ such that $f(a) \neq 0$.*

Proof. Extend a to a regular sequence $a, t_1, \ldots, t_r$ of length $\dim A$. Then the normalization of $A/(t_1, \ldots, t_r)$ is a finite direct sum of copies of $\mathbb{C}[\![t]\!]$. Now the result is clear. $\square$

5. Extending the family over boundary strata

Here we prove Theorem 0.1 and Theorem 0.2. Note that Theorem 0.1 holds for $n \geq 3$ by Theorems 2.33 and 3.8. We use Theorem 0.2 for $n \geq 3$ to prove Theorem 0.1 for $n < 3$.

As usual, let P be the toric monoid associated to a strictly convex rational polyhedral cone $\sigma_P \subset A_1(Y)_{\mathbb{R}}$ which contains the Mori cone $\mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}}$. We have $\mathfrak{m} = P \setminus \{0\}$. For a monomial ideal $I \subset P$ we define

$$A_I := \bigoplus_{q \in B(\mathbb{Z})} R_I \cdot \vartheta_q$$

where $R_I = \Bbbk[P]/I$. We take throughout $\mathfrak{D} = \mathfrak{D}^{\mathrm{can}}$.

Assumptions 5.1. *For any monomial ideal I with $\sqrt{I} = \mathfrak{m}$, the multiplication rule of Theorem 2.38 defines an R_I-algebra structure on A_I, so that $A_I \otimes_{R_I} R_{\mathfrak{m}} = H^0(\mathbb{V}_n, \mathcal{O}_{\mathbb{V}_n})$.*

Note we have already shown that Assumptions 5.1 hold if $n \geq 3$.

Let $G \subset B(\mathbb{Z})$ be a finite collection of integer points such that the corresponding functions ϑ_q generate the $\Bbbk$-algebra $H^0(\mathbb{V}_n, \mathcal{O}_{\mathbb{V}_n})$. (Then the ϑ_q, $q \in G$ generate A_I as an R_I-algebra if $\sqrt{I} = \mathfrak{m}$ and Assumptions 5.1 hold.) Note for $n \geq 3$ we can take for G the points $\{v_i\}$, and for $n = 1, 2$, one can make a simple choice for G, see §5.1.

Lemma 5.2. *For any monomial ideal $J \subset P$, $\bigcap_{k>0}(J + \mathfrak{m}^k) = J$.*

Proof. The inclusion $\supset$ is obvious. For the other direction, as the intersection is a monomial ideal, it's enough to consider a monomial in the intersection. But notice that a monomial is in $J + \mathfrak{m}^k$ iff it is either in J or in $\mathfrak{m}^k$. The result follows since $\bigcap \mathfrak{m}^k = 0$. $\square$

Assume 5.1. We let $\mathcal{A}$ be the collection of monomial ideals $J \subset P$ with the following properties:

1. There is an R_J-algebra structure on A_J such that the canonical isomorphism of R-modules $A_J \otimes_{R_J} R_{I+J} = A_{I+J}$ is an algebra isomorphism, for all $\sqrt{I} = \mathfrak{m}$.
2. $\vartheta_q, q \in G$ generate A_J as an R_J-algebra.

By the lemma, the algebra structure in (1) is unique if it exists. The algebra structure on all A_I determines such a structure on $\hat{A} := \varprojlim_{\sqrt{I}=\mathfrak{m}} A_I$, $\hat{A}_J := \varprojlim_{\sqrt{I}=\mathfrak{m}} A_{I+J}$. Also, there are canonical inclusions

$$\hat{A} \subset \prod_{q \in B(\mathbb{Z})} \hat{R} \cdot \vartheta_q$$

$$\hat{A}_J \subset \prod_{q \in B(\mathbb{Z})} \hat{R}_J \cdot \vartheta_q$$

where $\hat{R}, \hat{R}_J$ are the completions of R, R_J at $\mathfrak{m}, J + \mathfrak{m}$ (here the direct products are viewed purely as $\hat{R}, \hat{R}_J$ modules). We can also view

$$A_J := \bigoplus_{q \in B(\mathbb{Z})} R_J \cdot \vartheta_q \subset \prod_{q \in B(\mathbb{Z})} \hat{R}_J \cdot \vartheta_q.$$

It is clear that $A_J \subset \hat{A}_J$ (as submodules of the direct product). Thus (1) holds if and only if the following, (1′), holds: For each $p, q \in B(\mathbb{Z})$, at most finitely many $z^C \vartheta_s$ with $C \notin J$ appear in the product expansion of Theorem 2.38 for $\vartheta_p \cdot \vartheta_q \in \hat{A}_J$.

Lemma 5.3. *If $J \in \mathcal{A}$ and $J \subset J'$, then $J' \in \mathcal{A}$. In addition, $\mathcal{A}$ is closed under finite intersections.*

Proof. The first statement is clear. Now assume $J_1, J_2 \in \mathcal{A}$. It's clear that (1′) holds for $J_1 \cap J_2$, so $A_{J_1 \cap J_2}$ is an algebra. Moreover we have an exact sequence of $\mathbb{k}$-modules

$$0 \to A_{J_1 \cap J_2} \to A_{J_1} \times A_{J_2} \to A_{J_1 + J_2} \to 0$$

exhibiting $A_{J_1 \cap J_2}$ as the fibred product $A_{J_1} \times_{A_{J_1+J_2}} A_{J_2} =: A_1 \times_B A_2 =: A$. We now show this fibred product is a finitely generated $\mathbb{k}$-algebra. Indeed, note that since the maps $A_1, A_2 \to B$ are surjective, so are the maps $A \to A_i$. Let $\{u_i\}$ be a generating set for the ideal $\ker(A_2 \to B)$. Since A_i is Noetherian, one can find a finite such set. Note that $\tilde{u}_i = (0, u_i) \in A$. In addition, choose finite sets $\{x_i\}$, $\{y_j\}$ generating A_1 and A_2 as $\mathbb{k}$-algebras. For each of these elements, choose a lift to A, giving a finite set of lifts $\{\tilde{u}_i, \tilde{x}_i, \tilde{y}_i\}$, which we claim generate A. Indeed, given $(x, y) \in A$, one can subtract a polynomial in the $\tilde{x}_i$'s to obtain $(0, y')$. Necessarily $y' \in \ker(A_2 \to B)$, and hence we can write $y' = \sum f_i u_i$ with f_i a polynomial in the y_i's. Let $\tilde{f}_i$ be the same polynomial in the $\tilde{y}_i$'s. Then $\sum \tilde{f}_i \tilde{u}_i = (0, y')$, showing generation.

Thus $A_{J_1 \cap J_2}$ is a also a finitely generated $R_{J_1 \cap J_2}$-algebra. Now the generation statement follows from Lemma 5.4. ☐

Lemma 5.4. *Let $I, J \subset R$ be ideals in a Noetherian ring, with $I \cdot J = 0$, and let S be a finitely generated R-algebra, and $R[T_1, \ldots, T_m] \to S$ an R-algebra map which is surjective modulo I and J. Then the map is surjective.*

Proof. The associated map $\operatorname{Spec} S \to \mathbb{A}^m \times \operatorname{Spec} R$ is proper, and thus S is a finite $R[T_1, \ldots, T_m]$-module. Now we can apply Nakayama's lemma. $\square$

Proposition 5.5. *There is a unique minimal radical monomial ideal $I_{\min} \subset P$ such that (1) and (2) hold for any monomial ideal J with $I_{\min} \subset \sqrt{J}$.*

Proof. Certainly any ideal J with $\mathfrak{m} \subset \sqrt{J}$ lies in $\mathcal{A}$. Note that a radical monomial ideal is the complement of a union of faces of P, so there are only a finite number of such ideals. Suppose I_1, I_2 are two radical ideals such that $J_i \in \mathcal{A}$ for any J_i with $I_i \subset \sqrt{J_i}$. Note that any ideal J with $I_1 \cap I_2 \subset \sqrt{J}$ can be written as $J_1 \cap J_2$, with $I_i \subset \sqrt{J_i}$. Thus by Lemma 5.3, $J \in \mathcal{A}$. This shows the existence of $I_{\min}$. $\square$

Proposition 5.6. *Assume 5.1.*

1. *Suppose the intersection matrix $(D_i \cdot D_j)$ is not negative semi definite. Then $I_{\min} = (0) \subset \Bbbk[P]$.*
2. *Suppose $F \subset \sigma_P$ is a face such that F does not contain the class of some component of D. Then $I_{\min} \subset P \setminus F$.*

Proof. We prove both cases simultaneously, writing $F := P$ in case (1). We claim there exists an effective divisor W with support D such that $W \cdot D_j > 0$ for all D_j contained in F. For case (1), see Lemma 5.9. In case (2), say $[D_1] \notin F$. Then we can take $W = \sum a_i D_i$ where $a_1 \gg a_2 \gg \cdots \gg a_n > 0$.

The algebra structure depends only on the deformation type of (Y, D). By the local Torelli theorem for Looijenga pairs, see [L81], II.2.5, we may replace (Y, D) by a deformation equivalent pair such that any irreducible curve $C \subset Y$ intersects D.

Let $\overline{\mathrm{NE}}(Y)_{\mathbb{R}_{\geq 0}} \subset A_1(Y, \mathbb{R})$ denote the closure of $\mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}}$. Let $F' := \overline{\mathrm{NE}}(Y)_{\mathbb{R}_{\geq 0}} \cap F$, a face of $\overline{\mathrm{NE}}(Y)_{\mathbb{R}_{\geq 0}}$. Define $\Delta = D - \epsilon W$, $0 < \epsilon \ll 1$. Then (Y, Δ) is KLT (Kawamata log terminal). We claim $K_Y + \Delta \sim -\epsilon W$ is negative on $F' \setminus \{0\}$. By construction $(K_Y + \Delta) \cdot D_j < 0$ for $[D_j] \in F'$ and $(K_Y + \Delta) \cdot C < 0$ for $C \not\subset D$. Let N be a nef divisor such that $F' = \overline{\mathrm{NE}}(Y)_{\mathbb{R}_{\geq 0}} \cap N^{\perp}$. Then $aN - (K_Y + \Delta)$ is nef and big for $a \gg 0$, and thus some multiple defines a birational morphism g by the basepoint-free theorem [KM98], Theorem 3.3. Thus $(K_Y + \Delta)^{\perp} \cap F'$ is generated by exceptional curves of g. We deduce that $(K_Y + \Delta)^{\perp} \cap F' = \{0\}$ and $(K_Y + \Delta)$ is negative on $F' \setminus \{0\}$ as claimed. Now by the cone theorem [KM98], Theorem 3.7, $\overline{\mathrm{NE}}(Y)_{\mathbb{R}_{\geq 0}}$ is rational polyhedral near F' and there is a contraction $p \colon Y \to \bar{Y}$ such that F' is generated by the classes of curves contracted by p. It follows that we can find $\mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}} \subset \sigma_{P'} \subset \sigma_P$ such that F' is a face of $\sigma_{P'}$. Now the algebra structure for P comes from P' by base extension, so (replacing P by P') we can assume $F = F'$, and thus that W is positive on $F \setminus \{0\}$.

Now let J be a monomial ideal with $\sqrt{J} = P \setminus F$. Consider condition (1'). By the T^D-equivariance of Theorem 3.43, any $z^C \vartheta_s$ that appears in $\vartheta_p \cdot \vartheta_q$ has the same weight for T^D. Thus it is enough to show that the map

$$w\colon B(\mathbb{Z}) \times (P \setminus J) \to \chi(T^D), \quad (q, C) \mapsto w(q) + w(C)$$

has finite fibres. It is enough to consider fibres of $\sigma(\mathbb{Z}) \times (P \setminus J) \to \chi(T^D)$ for each $\sigma \in \Sigma_{\max}$. Note that $\sigma(\mathbb{Z}) \times P$ is the set of integral points of a rational polyhedral cone, and w is linear on this set. Thus it is enough to check that $\ker(w) \cap F = 0$. So suppose we have $q \in \sigma(\mathbb{Z})$, $C \in F$ with $w(q) + w(C) = 0$. Say $\sigma = \sigma_{i,i+1}$. Then $q = av_i + bv_{i+1}$, for $a, b \in \mathbb{Z}_{\geq 0}$. We have

$$w(q) + w(C) = ae_{D_i} + be_{D_{i+1}} + \sum_j (C \cdot D_j) e_{D_j};$$

thus if this is zero, we have $C \cdot D_j \leq 0$ for all j. In particular, $W \cdot C \leq 0$. Since W is positive on $F \setminus \{0\}$, $C = 0$. Now necessarily $a = b = q = 0$. This proves ($1'$).

For (2), let $A'_J \subset A_J$ be the subalgebra generated by the $\vartheta_q, q \in G$. Fix a weight $w \in \chi(T)$. To show $A'_J = A_J$ it is enough to show that each of the finitely many $z^C \vartheta_q \in A_J$, with $(q, C) \in B(\mathbb{Z}) \times (P \setminus J)$ of weight w lies in A'_J (since the $z^C \vartheta_q$ give a $\Bbbk$-basis of A_J). We argue by decreasing induction on $\operatorname{ord}_{\mathfrak{m}}(C)$ (see Definition 2.26). Since the set of possible (q, C) is finite, there is an upper bound on the possible $\operatorname{ord}_{\mathfrak{m}}$'s. so the claim is vacuously true for large $\operatorname{ord}_{\mathfrak{m}}$. Consider $z^C \cdot \vartheta_p$, with $\operatorname{ord}_{\mathfrak{m}}(C) = h$. Since the ϑ_q generate A_J modulo $\mathfrak{m}$, we can find $a \in A'_J$ such that

$$\vartheta_p = a + m$$

with $m \in \mathfrak{m} \cdot A_J$. Moreover, we can assume a, and thus m, is homogeneous for the T^D action. Now

$$z^C \vartheta_p = z^C a + z^C m.$$

Clearly $z^C m$ is a sum of terms $z^D \vartheta_q$ of weight w and $\operatorname{ord}_{\mathfrak{m}}(D) > h$, so $z^C m \in A'_J$ by induction. □

Remark 5.7. Suppose $p\colon Y \to \bar{Y}$ is a contraction such that some component of D is not contracted by p. Let F be the face of $\operatorname{NE}(Y)_{\mathbb{R}_{\geq 0}}$ generated by classes of curves contracted by p. Then $\operatorname{NE}(Y)_{\mathbb{R}_{\geq 0}}$ is rational polyhedral near F. (This follows from the cone theorem, cf. the proof of Proposition 5.6.) In particular there exists a rational polyhedral cone $\sigma_P \subset A_1(Y, \mathbb{R})$ such that $\operatorname{NE}(Y)_{\mathbb{R}_{\geq 0}} \subset \sigma_P$ and σ_P coincides with $\operatorname{NE}(Y)_{\mathbb{R}_{\geq 0}}$ near F.

Corollary 5.8. *Theorem 0.2 holds for $n \geq 3$.*

Proof. Immediate from Proposition 5.6. □

5.1. **Proof of Theorems 0.1 and 0.2 for** $n < 3$**.** If $n \leq 2$, let $p : (Y', D') \to (Y, D)$ be a toric blowup with $n' \geq 3$. We can find a strictly convex rational polyhedral cone $\sigma_{P'}$ with

$$\mathrm{NE}(Y')_{\mathbb{R}_{\geq 0}} \subset \sigma_{P'} \subset A_1(Y')_{\mathbb{R}}$$

which has a face F spanned by the p-exceptional curves, and which surjects onto $\sigma_P \subset A_1(Y)_{\mathbb{R}}$. For any monomial ideal $I \subset P$ with $\sqrt{I} = \mathfrak{m}$, let $J \subset P'$ be the inverse image. Then $\sqrt{J}$ is the prime monomial ideal associated to the face F. Since the exceptional curves are a proper subset of D' we have $\sqrt{J} \in \mathcal{A}(Y')$ by Proposition 5.6. Now restrict the family $\mathcal{X}_J \to \operatorname{Spec} \Bbbk[P']/J$ to $\operatorname{Spec} \Bbbk[P]/I \subset \operatorname{Spec} \Bbbk[P']/J$ (induced by the surjection $P' \to P$). This gives an algebra structure on

$$A_I := \bigoplus_{q \in B(\mathbb{Z})} (\Bbbk[P]/I)\vartheta_q.$$

We claim this is precisely the algebra described by Theorem 2.38. The argument is just as in the proof of Proposition 3.12: We have $B_{(Y',D')} = B_{(Y,D)}$ and for $\eta : \Bbbk[\mathrm{NE}(Y')] \to \Bbbk[\mathrm{NE}(Y)]$ the natural surjection (induced by $p_* : A_1(Y') \to A_1(Y)$), $\eta(\mathfrak{D}^{\mathrm{can}}) = \mathfrak{D}^{\mathrm{can}}$ (i.e. the rays are the same, and we apply p_* to the decoration function). This does not literally give a bijection on broken lines (because different monomials in the decoration of a ray in $\mathfrak{D}^{\mathrm{can}}_{(Y',D')}$ could map to the same monomial under p_*). However, by Equation (3.4), with z a point close to q,

$$\begin{aligned}\sum_{\substack{(\gamma_1,\gamma_2)\\ \mathrm{Limits}(\gamma_i)=(q_i,z)\\ s(\gamma_1)+s(\gamma_2)=q}} c(\gamma_1)c(\gamma_2) &= \sum_{\substack{(\gamma_1,\gamma_2)\\ \mathrm{Limits}(\gamma_i)=(q_i,z)\\ s(\gamma_1)+s(\gamma_2)=q}} \eta\left(\sum_{\eta(\gamma_1')=\gamma_1} c(\gamma_1')\right)\eta\left(\sum_{\eta(\gamma_2')=\gamma_2} c(\gamma_2')\right)\\ &= \eta\left(\sum_{\substack{(\gamma_1',\gamma_2')\\ \mathrm{Limits}(\gamma_i')=(q_i,z)\\ s(\gamma_1')+s(\gamma_2')=q}} c(\gamma_1')c(\gamma_2')\right)\end{aligned}$$

which implies the claim.

Now to complete the proof we need to check that the the fibre over the zero stratum of $\operatorname{Spec} \Bbbk[P]$ is $\mathbb{V}_n$. We check this directly using the algebra structure. We will do the case of $n = 1$, as $n = 2$ is similar (and simpler). We cut $B = B_{(Y,D)}$ along the unique ray $\rho \in \Sigma$, and consider the image under the developing map of the complement. This is a strictly convex rational cone in $\mathbb{R}^2$. Let w, w' be the primitive generators of the two boundary ray. Modulo $\mathfrak{m}$ the decoration on every scattering ray is trivial, so every broken line is straight. Moreover, no line can cross ρ (or the attached monomial becomes trivial by the strict convexity of φ). Now it follows for any $x \in B(\mathbb{R}) \setminus \rho$ and any $q \in (B \setminus \rho)(\mathbb{Z})$ there is a unique (straight) broken line with Limits $= (q, x)$, while there are exactly two (straight) broken lines with Limits $= (v, x)$, $v = v_1$ — under the developing map these become two distinct straight lines with directions w, w'. Performing a toric blowup of (Y, D) to get $n' = 3$ can be accomplished by subdividing the cone generated by w and w' along the rays generated by $w + w'$ and $2w + w'$. Then by Theorem 0.2 in the case $n = 3$, we see that $A_{\mathfrak{m}}$ is generated over $\Bbbk$ by

$$\vartheta_v = \vartheta_w = \vartheta_{w'}, \vartheta_{w+w'}, \vartheta_{2w+w'}$$

where we abuse notation and use the same symbol for an integer point in the convex cone generated by w and w', and the corresponding point in $B(\mathbb{Z})$. Now applying the multiplication rule of Theorem 2.38 one checks easily the equalities:

$$\begin{aligned} \vartheta_v^2 &= \vartheta_{w+w'} + \vartheta_{2v} \\ \vartheta_v \cdot \vartheta_{w+w'} &= \vartheta_{2w+w'} + \vartheta_{w+2w'} \\ \vartheta_{2w+w'} \cdot \vartheta_{w+2w'} &= \vartheta_{3w+3w'} = \vartheta_{w+w'}^3. \end{aligned}$$

It follows that

$$\vartheta_{2w+w'} \cdot \vartheta_v \cdot \vartheta_{w+w'} = \vartheta_{2w+w'}^2 + \vartheta_{w+w'}^3$$

and thus $A_{\mathfrak{m}} = \Bbbk[x, y, z]/(xyz - x^2 - z^3)$, which is isomorphic to the ring of sections

$$\bigoplus_{m \geq 0} H^0(C, \mathcal{O}(m))$$

for a line bundle $\mathcal{O}(1)$ of degree one on an irreducible rational nodal curve C of arithmetic genus 1. Thus $\operatorname{Spec} A_{\mathfrak{m}} = \mathbb{V}_1$. ☐

Now Theorems 0.1 and 0.2 holds for all n by Proposition 5.6.

5.2. D **positive.**

Lemma 5.9. *The following are equivalent for a Looijenga pair* (Y, D)*:*

- (1.1) *There exist integers* $a_1, \ldots, a_n$ *such that* $(\sum a_i D_i)^2 > 0$.
- (1.2) *There exist positive integers* $b_1, \ldots, b_n$ *such that* $(\sum b_i D_i) \cdot D_j > 0$ *for all* j.
- (1.3) $Y \setminus D$ *is the minimal resolution of an affine surface with (at worst) Du Val singularities.*
- (1.4) *There exist* $0 < c_i < 1$ *such that* $-(K_Y + \sum c_i D_i)$ *is nef and big.*

If any of the above equivalent conditions hold, then so do the following:

- (2.1) *The Mori cone* $\mathrm{NE}(Y)_{\mathbb{R}_{\geq 0}}$ *is rational polyhedral, generated by finitely many classes of rational curves. Every nef line bundle on* Y *is semi-ample.*
- (2.2) *The subgroup* G *of* $\mathrm{Aut}(\mathrm{Pic}(Y), \langle \cdot, \cdot \rangle)$ *fixing the classes* $[D_i]$ *is finite.*
- (2.3) *The union* $R \subset Y$ *of all curves disjoint from* D *is contractible.*

Proof. We have

$$K_Y + \sum c_i D_i = (K_Y + D) - \sum (1 - c_i) D_i = - \sum (1 - c_i) D_i$$

so (1.2) and (1.4) are equivalent, and (1.2) obviously implies (1.1).

If (1.1) holds then $(D^{\perp}, \langle \cdot, \cdot \rangle)$ is negative definite, by the Hodge Index Theorem, and this implies (2.2) and (2.3).

Suppose (1.4) holds. By the basepoint-free theorem [KM98], 3.3, the linear system

$$|m(\sum b_i D_i)| = |-m(K_Y + \sum c_i D_i)|$$

defines a birational morphism for $m \in \mathbb{N}$ sufficiently large, with exceptional locus the union R of curves disjoint from D. Adjunction shows R is a contractible configuration of (-2)-curves, which gives (1.3). (2.1) follows from the cone theorem [KM98], 3.7.

We show (1.1) implies (1.2). By the Riemann–Roch theorem, if W is a Weil divisor (on any smooth surface) and $W^2 > 0$ then either W or $-W$ is big (i.e., the rational map given by $|nW|$ is birational for sufficiently large n). So, possibly replacing the divisor by its negative, we may assume $W = \sum a_i D_i$ is big. Write

$$W' = \sum_{a_i > 0} a_i D_i = W + \sum_{-a_i > 0} (-a_i) D_i.$$

Thus W' is big, and replacing W by nW', we may assume all $a_i \geq 0$ and $|W|$ defines a birational (rational) map. Subtracting off the divisorial base-locus (which does not affect the rational map) we may further assume the base locus is at most zero dimensional. Now $W = \sum b_i D_i$ is effective, nef and big, and supported on D. We show we may assume that in addition $b_i > 0$ and $W \cdot D_i > 0$ for each i. If $W \cdot D_i > 0$, then we may assume $b_i > 0$ (by adding ϵD_i to W if necessary). Now consider the set $S \subset \{1, \ldots, n\}$ of components D_i of D such that $W \cdot D_i = 0$. By connectedness of D we find $b_i > 0$ for each $i \in S$. Thus $\mathrm{Supp}(W) = D$. By the Hodge index theorem the intersection matrix $(D_i \cdot D_j)_{i,j \in S}$ is negative definite. Hence there exists a linear combination $E = \sum_{i \in S} \alpha_i D_i$, with $\alpha_i \in \mathbb{Z}_{>0}$ for each $i \in S$, such that $E \cdot D_i < 0$ for each $i \in S$. Now replacing W by $W - \epsilon E$, we obtain $W \cdot D_i > 0$ for each $i = 1, \ldots, n$.

Finally we show (1.3) implies (1.1). Since U is affine, we have $U = Y' \setminus D'$ where Y' is a normal projective surface and D' is a Weil divisor such that D' is the support of an ample divisor A. Let $\pi \colon \tilde{Y} \to Y'$ be a resolution of singularities such that π is an isomorphism over U and the inclusion $U \subset Y$ extends to a birational morphism $f \colon \tilde{Y} \to Y$. Let $\tilde{D}$ be the inverse image of D' under π, so $\tilde{Y} \setminus \tilde{D} = Y' \setminus D' = U$. The divisor $\pi^* A$ has support $\tilde{D}$. So we can write $\pi^* A = f^*(\sum a_i D_i) + \sum \mu_j E_j$ where the E_j are the f-exceptional curves and $a_i, \mu_j \in \mathbb{Z}$. Then $(\sum a_i D_i)^2 \geq (\pi^* A)^2 = A^2 > 0$. $\square$

Corollary 5.10. *Let (Y, D) be a Looijenga pair satisfying any of the equivalent conditions of Lemma 5.9. Let $P = \mathrm{NE}(Y)$. The multiplication rule Theorem 2.38 determines a finitely generated T^D-equivariant $R = \Bbbk[P]$-algebra structure on the free R-module*

$$A = \bigoplus_{q \in B(\mathbb{Z})} R \cdot \vartheta_q.$$

Furthermore, $\operatorname{Spec} A \to \operatorname{Spec} R$ *is a flat affine family of Gorenstein SLC surfaces with central fibre $\mathbb{V}_n$, and smooth generic fibre. Any collection of ϑ_q whose restrictions generate $A/\mathfrak{m} = H^0(\mathbb{V}_n, \mathcal{O}_{\mathbb{V}_n})$ generate A as an R-algebra. In particular the ϑ_{v_i} generate for $n \geq 3$.*

Proof. Everything but the singularity statement follows from Theorem 0.2. The Gorenstein SLC locus in the base is open, and T^D-equivariant. By the positivity there is a one parameter subgroup $\mathbb{G}_m \subset T^D$ so that the torus fixed point $0 \in \operatorname{Spec} R$ is in the closure of every orbit. Thus the locus is all of $\operatorname{Spec} R$. □

In Part II we will prove that when D is positive, our mirror family admits a canonical fibrewise T^D-equivariant compactification $\mathcal{X} \subset (\mathcal{Z}, \mathcal{D})$. The restriction $(\mathcal{Z}, \mathcal{D}) \to T_Y$ comes with a trivialization $\mathcal{D} \xrightarrow{\sim} D_* \times T_Y$. We will show that $(\mathcal{Z}, \mathcal{D})$ is the universal family of Looijenga pairs (Z, D_Z) deformation equivalent to (Y, D) together with a choice of isomorphism $D_Z \xrightarrow{\sim} D_*$. Now for any positive pair (Z, D_Z) together with a choice of isomorphism $\phi\colon D_Z \xrightarrow{\sim} D_*$, our construction equips the complement $U = Z \setminus D_Z$ with canonical theta functions ϑ_q, $q \in B_{(Z,D)}(\mathbb{Z})$. We will give a characterisation in terms of the intrinsic geometry of (Z, D_Z). Changing the choice of isomorphism ϕ changes ϑ_q by a character of $T^D = \mathrm{Aut}^0(D_*)$, the identity component of $\mathrm{Aut}(D_*)$. Here we illustrate with two examples:

Example 5.11. Consider first the case (Y, D) a 5-cycle of (-1)-curves on the (unique) degree 5 del Pezzo surface, Example 3.7. In this case $T^D = T_Y = \mathrm{Pic}(Y) \otimes_{\mathbb{Z}} \mathbb{G}_m$, and thus by the T^D-equivariance, all fibres of the restriction $\mathcal{X} \to T_Y$ are isomorphic. We consider the fibre over the identity $e \in T_Y$, thus specializing the equations of Example 3.7 by setting all $z^{D_i} = 1$. It's well known that these equations define an embedding of the original $Y \setminus D$ into $\mathbb{A}^5$ — if we take the closure in $\mathbb{P}^5$ (for the standard compactification $\mathbb{A}^5 \subset \mathbb{P}^5$) one checks easily we obtain Y with D the hyperplane section at infinity.

Now it is easy to compute the zeroes and poles:

$$(\vartheta_{v_i}) = E_i + D_i - D_{i+2} - D_{i-2}$$

(indices mod 5). In particular $\{\vartheta_{v_i} = 0\} = E_i \cap U \subset U$, which characterizes ϑ_{v_i} up to scaling.

Example 5.12. Now let $(Y, D = D_1 + D_2 + D_3)$ be (the deformation type of) a cubic surface together with a triangle of lines. Let $\mathcal{X} \subset \operatorname{Spec}(\Bbbk[\mathrm{NE}(Y)]) \times \mathbb{A}^3$ be the canonical embedding given by $\vartheta_i := \vartheta_{v_i}$, $i = 1, 2, 3$. We will show in Part II that this is given by the equation

$$\vartheta_1\vartheta_2\vartheta_3 = \sum_i z^{D_i}\vartheta_i^2 + \sum_i \left(\sum_j z^{E_{ij}}\right) z^{D_i}\vartheta_i + \sum_\pi z^{\pi^* H} + 4z^{D_1+D_2+D_3}.$$

Here the E_{ij} are the interior (-1)-curves meeting D_i, and the sum over π is the sum over all possible toric models $\pi\colon Y \to \bar{Y}$ of (Y, D) to a pair $(\bar{Y}, \bar{D})$ isomorphic to $\mathbb{P}^2$ with its toric boundary. (Such π are permuted simply transitively by the Weyl group $W(D_4)$ by [L81], Prop. 4.5, p. 283.) The same family, in the same canonical coordinates, was discovered by Oblomkov [Ob04]. As we learned from Dolgachev, after a change of variables (in $\mathbb{A}^3$), and restricting to T_Y (the locus over which the fibers have at worst Du Val singularities) this is identified with the universal family of affine cubic surfaces (the complement to a triangle of lines on projective cubic surface) constructed by Cayley in [C1869]. The universal family of cubic surfaces with triangle is obtained as the closure in $\mathbb{A}^3 \subset \mathbb{P}^3$. In particular, as in the first example, our mirror family compactifies naturally to the universal family of Looijenga pairs deformation equivalent to the original (Y, D). There is again a geometric characterisation of ϑ_i (up to scaling): The linear system $|-K_Y - D_i| = |D_j + D_k|$ (here $\{i, j, k\} = \{1, 2, 3\}$) is a basepoint free pencil. It defines a ruling $\pi : Y \to \mathbb{P}^1$ which restricts to a double cover $D_i \to \mathbb{P}^1$. Let $\{a, b\} \subset \mathbb{P}^1$ be the branch points of $\pi|_{D_i}$. Let $p = \pi(D_j + D_k) \in \mathbb{P}^1$. There is a unique point $q \in \mathbb{P}^1 \setminus \{a, b, p\}$ fixed by the unique involution of $\mathbb{P}^1$ interchanging a and b and fixing p. Let $Q = \pi^*(q) \in |D_j + D_k|$ be the corresponding divisor. The curve $Q \subset \mathbb{P}^3$ is a smooth conic. In Part II we will show

Proposition 5.13. $(\vartheta_i) = Q - D_j - D_k$.

5.3. **Relation to cluster varieties.** Fock and Goncharov define a (rational) birational automorphism $h : T \dashrightarrow T$ of a split algebraic torus to be *positive* if the pullback of any character is a ratio of Laurent polynomials with positive integer coefficients. A *positive atlas* $\mathcal{W}$ is a collection of (rational) birational isomorphisms $T \dashrightarrow W$ such that the transition functions $h : T \dashrightarrow T$ are positive. They then point out two flavors $\mathcal{X}, \mathcal{A}$ of cluster varieties which are examples. We refer to [FG09] for the definition of cluster variety and precise statements. We write $U(\mathcal{W})$ for the affine closure (that is, the spectrum of the global sections of the structure sheaf) of the scheme over $\mathbb{Z}$ obtained by glueing together the tori in the atlas.

Note that a split torus has a nowhere vanishing top degree form Ω canonical up to sign, namely $\operatorname{dlog}(x_1) \wedge \cdots \wedge \operatorname{dlog}(x_n)$ for a basis $x_1, \ldots, x_n$ of characters. We observe:

Lemma 5.14. *$U(\mathcal{W})$ is an affine log Calabi Yau variety, that is, has a nowhere vanishing top degree form with log poles at infinity, unique up to scaling.*

Proof. The tori cover U up to a codimension two subset. It is immediate from the explicit formulae for cluster transformation, [FG09],(13),(14) that the canonical forms Ω agree up to sign. It follows that they extend to a nowhere vanishing form on U. Uniqueness follows from the same statement on a torus. $\square$

Fock and Goncharov define the tropical points $\mathcal{W}(\mathbb{Z}^t)$ of any positive atlas and observe that each chart canonically $g : T \dashrightarrow W$ identifies $\mathcal{W}(\mathbb{Z}^t)$ with the lattice of co-characters of T. They associate to each *seed* i (a certain combinatorial object introduced by Fomin and Zelevinsky) a pair of positive atlases $\mathcal{A}_i, \mathcal{X}_i$. Each seed has a natural Langlands dual $i^\vee$. They conjecture a remarkable duality. A universally positive Laurent polynomial means a regular function on U which in each toric chart is a Laurent polynomial with positive integer coefficients. These form a semi-group under addition. Let $\mathbb{E}(\mathcal{W})$ be the set of minimal elements. They conjecture there is a natural bijection $\mathcal{A}_i(\mathbb{Z}^t) \to \mathbb{E}(\mathcal{X}_{i^\vee})$, and moreover that $\mathbb{E}(\mathcal{X}_{i^\vee})$ gives a $\mathbb{Z}$-basis of $H^0(U(\mathcal{X}_{i^\vee}), \mathcal{O})$. We believe this is a special case of our Conjecture 0.8.

We note first that their tropical points are canonically identified with our (weighted) log canonical centers $B(\mathbb{Z})$ of §0.4. This is immediate from the definitions: $B(\mathbb{Z})$ is a birational invariant of the form Ω and so can be computed on any of the toric charts; for a torus $B(\mathbb{Z})$ is the lattice of co-characters. We believe that the $U(\mathcal{A}_i)$ and $U(\mathcal{X}_{i^\vee})$ are mirror Calabi-Yau varieties and under the identification $\mathcal{A}_i(\mathbb{Z}^t) = B(\mathbb{Z})$ our theta functions ϑ_q give the minimal generators of the semi-group $\mathbb{E}(\mathcal{X}_{i^\vee})$.

Thus we believe that our theta functions give a natural generalisation of the Fock–Goncharov minimal universally positive Laurent polynomials to any affine Calabi-Yau with maximal boundary (recall this means that B has the maximal possible dimension $d = \dim(U)$), moreover that the multiplication rule for the generators is governed by counts of rational curves, according to the formula of Conjecture 0.8. We note this will include non-rational examples, which of course have no torus charts.

We have checked that this is true for the three cluster surfaces of finite type – associated to A_2, B_2 and G_2. The A_2 case is Example 5.11.

Example 5.12 is also an instance. In this case the total space of our deformation is the Fock-Goncharov cluster variety $\mathcal{X}_{\mathrm{PGL}_2,S}$, parameterizing PGL_2-local systems on the surface S given by the 2-sphere with 4 punctures (more precisely, $\mathcal{X}_{\mathrm{PGL}_2,S} = \mathcal{X}/T^D$, the quotient of our total space by the relative torus T^D). In this case (by Fock–Goncharov) $A(\mathbb{Z}^t) = B(\mathbb{Z})$ is the space of integral $\mathcal{A}$-laminations of S, which are (weighted) isotopy classes of simple loops in S. We have checked that under the correspondence our ϑ_q gives the trace of the monodromy transformation for the corresponding loop. Note we now have several quite different interpretations for ϑ_q: The current view as a trace function on a moduli space of local systems, the interpretation in terms of classical geometry of a cubic surface, Example 5.12, as functions arising from areas of holomorphic discs in the SYZ picture of mirror symmetry as in (0.6), or the tropical version of discs via broken lines, Theorem 0.7, and finally a (speculative) Floer theoretic incarnation via our Conjecture 0.8.

6. DEFORMATIONS OF CYCLIC QUOTIENT SINGULARITIES

In this section we describe the deformation theory of surface cyclic quotient singularities using a variant of our construction. The deformation theory of these singularities is now completely understood and has a beautiful and intricate structure. A key advance was obtained by Kollár and Shepherd-Barron [KSB88]. They showed that the irreducible components of the deformation space are in bijective correspondence with certain partial resolutions of the singularity, dubbed P-resolutions. J. Stevens obtained an explicit description of all P-resolutions and the deformation associated to each P-resolution [S91].

Our construction in this case is much easier than in the proof of Theorem 0.1 because, roughly speaking, there is no "scattering", see Remark 6.2. So we are able to prove versality results for our deformations. We also obtain a geometric interpretation of Stevens' combinatorial description of P-resolutions given by a real 2-torus fibration of the Milnor fiber.

Remark 6.1. More recently, Lisca obtained a classification of the Stein symplectic fillings of the link of the cyclic quotient singularity (a lens space) up to diffeomorphism [L08]. He conjectured that all such fillings arise as the Milnor fiber of a smoothing of the singularity. Lisca's conjecture was verified by Némethi and Popescu-Pampu [NPP10]. This establishes a bijective correspondence between components of the deformation space of the singularity and Stein symplectic fillings of its link up to diffeomorphism. It would be very interesting to establish an analogous correspondence for cusp singularities.

In this section, we use the notation $\mathbb{A}^m/\frac{1}{r}(a_1,\ldots,a_m)$ or just $\frac{1}{r}(a_1,\ldots,a_m)$ to denote the cyclic quotient singularity

$$\mathbb{A}^m/\mu_r, \quad \mu_r \ni \zeta\colon (x_1,\ldots,x_m) \mapsto (\zeta^{a_1}x_1,\ldots,\zeta^{a_m}x_m).$$

6.1. **Construction of the deformation.** Let $\bar{Y}$ be a smooth quasiprojective toric surface admitting a toric proper morphism $\bar{f}\colon \bar{Y} \to \mathbb{A}^1$. Thus the fan $\bar{\Sigma}$ of $\bar{Y}$ has support a half-plane and the morphism $\bar{f}$ corresponds to the linear projection $|\bar{\Sigma}| \to \mathbb{R}_{\geq 0}$. Let $\bar{D}$ be the toric boundary of $\bar{Y}$. Then $\bar{D}$ is a chain of curves $\bar{D} = \bar{D}_0 + \bar{D}_1 +$

$\cdots + \bar{D}_n + \bar{D}_{n+1}$ where $\bar{D}_0, \bar{D}_{n+1}$ are sections of $\bar{f}$ and $\bar{D}_1 + \cdots + \bar{D}_n$ is the reduction of the fiber of $\bar{f}$ over $0 \in \mathbb{A}^1$. Fix integers $l_1, \ldots, l_n \in \mathbb{Z}_{\geq 0}$. Let $\pi \colon Y \to \bar{Y}$ be the blowup of $\bar{Y}$ at distinct points x_{ij} in the interior of the boundary divisor $\bar{D}_i$ for each $1 \leq i \leq n$ and $1 \leq j \leq l_i$. Let E_{ij} denote the exceptional curve of π over x_{ij} and $f \colon Y \to \mathbb{A}^1$ the composition $f = \bar{f} \circ \pi$. Let $D = D_0 + \cdots + D_{n+1}$ denote the strict transform of the boundary $\bar{D}$ of $\bar{Y}$. Then $K_Y + D = 0$.

Let $P = \mathrm{NE}(Y/\mathbb{A}^1)$ be the monoid of classes of curves contracted by $f \colon Y \to \mathbb{A}^1$. Explicitly, $P = \mathbb{N}^{n+\sum l_i}$, generated by $D_1, \ldots, D_n$ and the π-exceptional curves E_{ij}. Thus $S := \operatorname{Spec} \Bbbk[P] = \mathbb{A}^{n+\sum l_i}$ with coordinates $z^{D_i}, 1 \leq i \leq n$ and the $z^{E_{ij}}$.

Define

$$\mathbb{S}_n := \mathbb{A}^2_{x_0,x_1} \cup \cdots \cup \mathbb{A}^2_{x_n,x_{n+1}} \subset \mathbb{A}^{n+2}_{x_0,\ldots,x_{n+1}},$$

the union of a chain of $n+1$ coordinate planes in $\mathbb{A}^{n+2}$. Let $\mathbb{B}_n \subset \mathbb{S}_n$ denote the union $\mathbb{A}^1_{x_0} \cup \mathbb{A}^1_{x_{n+1}}$ of coordinate lines, a closed subscheme of codimension 1. We construct a deformation of the pair $(\mathbb{S}_n, \mathbb{B}_n)$ over $\operatorname{Spec} \Bbbk[P]$ associated to (Y, D).

We define a singular integral affine manifold with boundary $(B, \partial B)$ together with a subdivision Σ into cones associated to (Y, D) as in §1.1. Thus B is a union of 2-dimensional convex cones $\sigma_{0,1}, \ldots, \sigma_{n,n+1}$ corresponding to the nodes of D, glued along rays $\rho_1, \ldots, \rho_n$ corresponding to the proper components $D_1, \ldots, D_n$ of D, and B has boundary ∂B the union of the two rays ρ_0, ρ_{n+1} corresponding to the affine components D_0, D_{n+1} of D. The integral affine structure on B is defined by declaring each cone $\sigma_{i,i+1}$ to be integral affine isomorphic to the positive quadrant in $\mathbb{R}^2$, and defining the integral affine structure on the union $\sigma_{i-1,i} \cup \sigma_{i,i+1}$ of two adjacent cones using the self-intersection number D_i^2 as in §1.1. Writing $B_0 := B \setminus \{0\}$ and $\partial B_0 := \partial B \setminus \{0\}$, the pair $(B_0, \partial B_0)$ is an *integral affine manifold with boundary* in the obvious sense. We write Σ for the polyhedral subdivision of B given by the cones $\{0\}$, ρ_i, and $\sigma_{i,i+1}$.

The developing map identifies B with a convex cone σ in $M_{\mathbb{R}} \simeq \mathbb{R}^2$. If π is an isomorphism then σ is a half-plane and Σ can be identified with $\bar{\Sigma}$, otherwise σ is strictly convex. Let $v_i \in M$ denote the primitive generator of ρ_i for $i = 0, \ldots, n+1$.

As explained in §0.6.2, the toric model $\pi \colon Y \to \bar{Y}$ corresponds to a deformation of the singular integral affine manifold with boundary $(B, \partial B)$ to the integral affine manifold with boundary $(\bar{B}, \partial \bar{B})$ given by the support $|\bar{\Sigma}|$ of the fan of $\bar{Y}$. This deformation is obtained by moving l_i singularities of type I_1 with invariant direction ρ_i from $0 \in B$ to infinity along the ray ρ_i, for each $i = 1, \ldots, n$. Let $\nu \colon B \to \bar{B}$ denote the induced piecewise linear identification.

Let $\varphi \colon B \to P^{\mathrm{gp}}$ denote the P-convex Σ-piecewise linear function defined (up to a linear function) by $p_{\rho_i} = D_i$ for $i = 1, \ldots, n$. Let $\mathfrak{D}$ denote the scattering diagram on

B with rays

$$\mathfrak{d}_i = (\rho_i, f_i = \prod_{j=1}^{l_i} (1 + z^{E_{ij}} z^{(-v_i, -\varphi(v_i))})), \quad i = 1, \ldots, n.$$

Remark 6.2. Compare with the similar diagram $\bar{\mathfrak{D}}_0$ of (3.6) used in the proof of Theorem 0.1. In that case $\bar{\mathfrak{D}}_0$ is only the *input* to the Kontsevich-Soibelman lemma, Theorem 3.24, while the diagram we use for the construction, $\mathfrak{D}^{\mathrm{can}}$, is obtained using the complicated scattering process via $\nu(\mathfrak{D}^{\mathrm{can}}) = \bar{\mathfrak{D}} = \mathrm{Scatter}(\bar{\mathfrak{D}}_0)$. The present case is vastly simpler: $\bar{B} \subset M_{\mathbb{R}}$ is a half-space, and so any outgoing ray obtained by the scattering process applied to $\bar{\mathfrak{D}}_0 := \nu(\mathfrak{D})$ lies in the opposite half-space. (In general, if the incoming rays are contained in a convex cone $C \subset \bar{B} \subset M_{\mathbb{R}}$, then the outgoing rays generated by the scattering process lie in the opposite cone $-C \subset M_{\mathbb{R}}$. This follows easily from the proof of the Kontsevich Soibelman lemma.) So restricting to $\bar{B}$ we obtain $\bar{\mathfrak{D}} := \mathrm{Scatter}(\bar{\mathfrak{D}}_0)|_{\bar{B}} = \bar{\mathfrak{D}}_0$. Alternatively, because Y is proper over $\mathbb{A}^1$, the only $\mathbb{A}^1$-classes are the E_{ij}, and thus $\mathfrak{D}^{\mathrm{can}} = \mathfrak{D}$.

Let $\hat{S}$ denote $\mathrm{Spf}\, \widehat{\Bbbk[P]}$, the formal spectrum of the completion of $\Bbbk[P]$ at the maximal monomial ideal $\mathfrak{m} = P \setminus \{0\}$. We construct a formal deformation $\hat{X}/\hat{S}$ of $\mathbb{S}_n$ over $\hat{S}$ by modifying the Mumford deformation determined by φ using the scattering diagram $\mathfrak{D}$ as in §2. We now spell this out explicitly.

Let $U_i = (x_i \neq 0) \subset \mathbb{S}_n$ for $i = 0, \ldots, n+1$. Thus

$$U_i = V(x_{i-1}x_{i+1}) \subset \mathbb{A}^2_{x_{i-1},x_{i+1}} \times \mathbb{G}_{m,x_i} \text{ for } i = 1, \ldots, n$$

and

$$U_0 = \mathbb{A}^1_{x_1} \times \mathbb{G}_{m,x_0}, \quad U_{n+1} = \mathbb{A}^1_{x_n} \times \mathbb{G}_{m,x_{n+1}}.$$

Note that $\bigcup U_i = \mathbb{S}_n^o := \mathbb{S}_n \setminus \{0\}$ and

$$U_i \cap U_{i+1} = \mathbb{G}^2_{m,x_i,x_{i+1}},$$

$$U_i \cap U_j = \emptyset \text{ for } |i-j| > 1.$$

We define deformations $\mathcal{U}_i/S$ of U_i over S as follows:

$$\mathcal{U}_i = V\left(x_{i-1}x_{i+1} - z^{D_i} x_i^{-D_i^2} \prod (1 + z^{E_{ij}} x_i^{-1})\right) \subset \mathbb{A}^2_{x_{i-1},x_{i+1}} \times \mathbb{G}_{m,x_i} \times S \quad \text{for } 1 \leq i \leq n,$$

$$\mathcal{U}_i = U_i \times S \quad \text{for } i = 0, n+1.$$

Let $\hat{\mathcal{U}}_i/\hat{S}$ denote the restrictions to $\hat{S} := \mathrm{Spf}\, \widehat{\Bbbk[P]}$. We glue the deformations $\mathcal{U}_i$ by identifying the open sets

$$\hat{\mathcal{U}}_i \supset (x_{i+1} \neq 0) = \mathbb{G}^2_{m,x_i,x_{i+1}} \times \hat{S}$$

and

$$\hat{\mathcal{U}}_{i+1} \supset (x_i \neq 0) = \mathbb{G}^2_{m,x_i,x_{i+1}} \times \hat{S}$$

via the identity map on $\mathbb{G}^2_{m,x_i,x_{i+1}} \times \hat{S}$, for $i = 0, \ldots, n$. Note that the gluings are purely toric because the only scattering rays are the rays ρ_i, $i = 1, \ldots, n$, of Σ, and the associated automorphisms have been incorporated into the definition of the open sets $\mathcal{U}_i$ as in Construction 2.18. As before, there is no compatibility condition for the gluing because there are no triple overlaps of the open sets U_i. Write

$$\hat{\mathcal{U}}_{i,i+1} := \hat{\mathcal{U}}_i \cap \hat{\mathcal{U}}_{i+1} = \mathbb{G}^2_{m,x_i,x_{i+1}} \times \hat{S}.$$

In the coordinate free notation of §2, letting $I \subset P$ be a monomial ideal such that $\sqrt{I} = \mathfrak{m}$ and using the subscript I to denote restriction to $\operatorname{Spec}(\Bbbk[P]/I)$, we have

$$\mathcal{U}_{i,i+1,I} = \operatorname{Spec} R_{\rho_i,\sigma_{i,i+1},I} = \operatorname{Spec} R_{\rho_{i+1},\sigma_{i,i+1},I} = \operatorname{Spec} R_{\sigma_{i,i+1},\sigma_{i,i+1},I},$$

$$x_j = z^{(v_j,\varphi(v_j))} \in R_{\sigma_{i,i+1},\sigma_{i,i+1},I} \quad \text{for } j = i, i+1,$$

$$\mathcal{U}_{i,I} = \operatorname{Spec}(R_{\rho_i,I}) = \operatorname{Spec}(R_{\rho_i,\sigma_{i-1,i},I} \times_{(R_{\rho_i,\rho_i,I})_{f_i}} R_{\rho_i,\sigma_{i,i+1},I}) \quad \text{for } i = 1, \ldots, n,$$

and

$$\mathcal{U}_{0,I} = \operatorname{Spec}(R_{\rho_0,\sigma_{0,1},I}), \quad \mathcal{U}_{n,I} = \operatorname{Spec}(R_{\rho_{n+1},\sigma_{n,n+1},I}).$$

We thus obtain a formal deformation $\hat{X}^o/\hat{S}$ of $\mathbb{S}_n^o$ over $\hat{S}$. We define a formal scheme $\hat{X}/\hat{S}$ by $\mathcal{O}_{\hat{X}} := i_*\mathcal{O}_{\hat{X}^o}$ where i denotes the inclusion $\mathbb{S}_n^o \subset \mathbb{S}_n$. We explain below that the coordinates x_i lift to global sections $\vartheta_{v_i} \in \Gamma(\hat{X}^o, \mathcal{O}_{\hat{X}^o})$. Thus by Lemma 2.34, $\hat{X}/\hat{S}$ is a formal deformation of $\mathbb{S}_n$ over $\hat{S}$.

Let $q \in B_0(\mathbb{Z})$ and $Q \in B \setminus \partial B$. We define broken lines γ in B_0 for q with endpoint Q as before (see Definition 2.22). For Q lying in the interior of $\sigma_{i,i+1}$ we have $\operatorname{Lift}_Q(q) := \sum_\gamma \operatorname{Mono}(\gamma) \in \Gamma(\hat{\mathcal{U}}_{i,i+1}, \mathcal{O}_{\hat{X}^o})$. We assert that the $\operatorname{Lift}_Q(q)$ for varying Q patch to define a lift $\vartheta_q \in \Gamma(\hat{X}^o, \mathcal{O}_{\hat{X}^o})$ of $z^q \in \Gamma(X, \mathcal{O}_X) = \Bbbk[\Sigma]$. In §3.2 the analogous statement was established by analyzing the behaviour of broken lines as the endpoint Q varies. The difficulty is entirely due to limits of broken lines passing through the singular point $0 \in B$. In the present case this cannot happen by Lemma 6.3 below. Indeed, if γ is a broken line in B_0 with endpoint Q and $\zeta\colon [0,1] \to B \setminus \partial B$ is a path from Q to Q' which does not pass through a ray at which γ bends, then we can deform γ to a family $\Gamma = \{\Gamma_s\}$ of broken lines in B_0 with endpoints $\zeta(s)$ obtained by applying a homothety and either truncating γ or extending the final segment linearly.

Lemma 6.3. *Let γ be a broken line in B_0 for q with endpoint Q. Then the ray $\rho \subset M_\mathbb{R}$ in the direction of the final segment of γ is not contained in the interior of σ, and if ρ coincides with one of the edges ρ_0, ρ_{n+1} of σ then $\mathbb{R}_{\geq 0}q$ coincides with the other edge.*

Proof. The broken line γ is convex when viewed from the origin $0 \in B$ (because all rays in the scattering diagram $\mathfrak{D}$ are outgoing rays). Consider the piecewise linear path $\hat{\gamma} \subset B$ obtained by extending the final segment of γ as far as possible. We must show that $\hat{\gamma}$ either meets the boundary of B or its final segment is parallel to one of the edges ρ_0, ρ_{n+1} of B. We may assume that the bend in γ at each ray ρ_i is as large as possible, that is, the change in the primitive integral tangent vector v is given by $v \mapsto v + l_i|v_i \wedge v|v_i$. Then $\nu(\hat{\gamma}) \subset \bar{B}$ is a straight line in the half-space $\bar{B} \subset M_{\mathbb{R}}$. Thus either $\hat{\gamma}$ meets the boundary of B or both its incoming and outgoing unbounded segments are parallel to an edge of B. $\square$

The global function ϑ_q for $q = av_i + bv_{i+1} \in \sigma_{i,i+1} \cap B_0(\mathbb{Z})$ is given by $z^{(q,\varphi(q))} = x_i^a x_{i+1}^b$ on $\mathcal{U}_{i,i+1}$ (because the only broken line for q with endpoint Q in the interior of $\sigma_{i,i+1}$ is given by the ray $Q + \mathbb{R}_{\geq 0} \cdot q$). (We remark that the analogous statement is *not* true in general in the case (Y, D) proper, because broken lines can "wrap around" the origin $0 \in B$.) Write $x_i := \vartheta_{v_i}$.

Next we verify that the formal deformation $\hat{X}/\hat{S}$ is algebraic. As in §5 this follows because the formal deformation admits a $\mathbb{G}_m$-action with positive weights on $\Gamma(\hat{X}, \mathcal{O}_{\hat{X}})$: The torus $T^D = \mathbb{G}_m^{n+2}$ acts on $\hat{X} \subset \mathbb{A}^{n+2}_{x_0,\ldots,x_{n+1}} \times \hat{S}$ via

$$T^D \ni t = (t_0, \ldots, t_{n+1})\colon \quad x_i \mapsto t_i \cdot x_i, \quad z^\beta \mapsto \left(\prod t_i^{D_i \cdot \beta}\right) \cdot z^\beta.$$

There exists a linear combination $A = \sum \alpha_i D_i$, such that $\alpha_i > 0$ for all i and A is f-ample. Write $\hat{R} := \Gamma(\mathcal{O}_{\hat{X}})$ and let $R_m \subset \hat{R}$ denote the homogeneous elements of weight m for the one-parameter subgroup $\mathbb{G}_m \to T^D$ corresponding to A. Then $R_0 = \Bbbk$, $R := \bigoplus_{m \geq 0} R_m \subset \hat{R}$ is the $\Bbbk[P]$-subalgebra generated by $x_0, \ldots, x_{n+1}$, and $\hat{R}$ is the completion of R with respect to the torus invariant maximal ideal $\mathfrak{m} \subset \Bbbk[P]$. Thus the formal deformation $\hat{X}/\hat{S}$ extends (uniquely) to an T^D-equivariant algebraic deformation X/S, given by $X = \operatorname{Spec} R$.

In fact, due to the finiteness of broken lines on B for $\mathfrak{D}$, the lifts ϑ_q are algebraic functions on the open patches $\mathcal{U}_i$ and we have the following more precise statement. The open embeddings $\hat{\mathcal{U}}_i \subset \hat{X}$ over $\hat{S}$ extend uniquely to T^D-equivariant open embeddings $\mathcal{U}_i \subset X$ over S. Moreover $\mathcal{U}_i = (x_i \neq 0) \subset X \subset \mathbb{A}^n_{x_0,\ldots,x_{n+1}} \times S$, so $\bigcup \mathcal{U}_i = X^o := X \setminus \{0\} \times S$.

The coordinate ring $R = \Gamma(X, \mathcal{O}_X)$ is a free $\Bbbk[P]$-module with basis ϑ_q for $q \in B(\mathbb{Z})$ (as usual we define $\vartheta_0 = 1$). Let $I \subset R$ be the (free) $\Bbbk[P]$-submodule generated by ϑ_q for $q \in B(\mathbb{Z}) \setminus \partial B$. Using the last clause of Lemma 6.3, we see that I is an ideal of R. The ideal I defines a closed subscheme $C \subset X/S$ of codimension one, flat over S, with special fiber $C_0 = \mathbb{B}_n \subset X_0 = \mathbb{S}_n$ the union of the coordinate lines $\mathbb{A}^1_{x_0}$, $\mathbb{A}^1_{x_{n+1}}$

corresponding to the edges ρ_0, ρ_{n+1} of B. We remark that $K_X + C = 0$, but neither C nor K_X is $\mathbb{Q}$-Cartier for $n > 1$.

The product of permutation groups

$$W := S_{l_1} \times \cdots \times S_{l_n}$$

acts on S by permuting the $z^{E_{ij}}$, and this action lifts to $(X, C)/S$.

Now suppose that π is not an isomorphism, so that $D_1 + \cdots + D_n$ is negative definite. Let $p\colon Y \to Y'$ denote the contraction of $D_1 + \cdots + D_n$. Then $D_1 + \cdots + D_n$ contracts to a cyclic quotient singularity $q \in Y'$ isomorphic to $0 \in \operatorname{Spec} \Bbbk[\sigma^\vee \cap N]$.

Let $S_p \subset S$ be the toric stratum of S associated to the contraction p. The restriction of the family $(X, C)/S$ to S_p is a purely toric deformation with general fiber $Z := \operatorname{Spec} \Bbbk[\sigma \cap M]$, the dual cyclic quotient singularity to $q \in Y'$, together with its toric boundary.

Let S' denote a W-invariant transverse slice to S_p. Explicitly, $S_p \subset S$ is the coordinate subspace $\mathbb{A}^n_{z^{D_1},\dots,z^{D_n}}$ and we may take

$$S' = V(z^{D_1} - 1, \cdots, z^{D_n} - 1) = \mathbb{A}^{\sum l_i}_{\{z^{E_{ij}}\}}.$$

Let $(X', C')/S'$ denote the restriction of $(X, C)/S$ to S'. Thus $(X', C')/(0 \in S')$ is a deformation of the cyclic quotient singularity Z together with its toric boundary.

6.2. ***P*-resolutions and deformations.** Let $p \in V$ be a normal surface singularity such that K_V is $\mathbb{Q}$-Cartier. Let N be the index of $p \in V$, that is, NK_V is Cartier at p and $N \in \mathbb{Z}_{>0}$ is minimal with this property. Then the *index one cover* of $p \in V$ is the μ_N Galois cover $q\colon W \to V$ given by

$$W = \underline{\operatorname{Spec}}_V(\mathcal{O}_V \oplus \mathcal{O}_V(K_V) \oplus \cdots \oplus \mathcal{O}_V((N-1)K_V)),$$

where the multiplication on $q_*\mathcal{O}_W$ is defined by choosing an isomorphism $\mathcal{O}_V(NK_V) \xrightarrow{\sim} \mathcal{O}_V$. We say a deformation $\mathcal{V}/(0 \in S)$ of V is $\mathbb{Q}$-*Gorenstein* if it is induced by an equivariant deformation of the index one cover. (If $p \in V$ is a quotient singularity and $0 \in S$ is a smooth curve germ this is equivalent to requiring that $K_{\mathcal{V}}$ is $\mathbb{Q}$-Cartier.)

A quotient singularity is said to be of *class T* if it admits a $\mathbb{Q}$-Gorenstein smoothing. The quotient singularities of class T are the Du Val singularities and the cyclic quotient singularities of the form $\frac{1}{dn^2}(1, dna - 1)$ for some $d, n, a \in \mathbb{Z}_{>0}$, $(a, n) = 1$.

If Z is a quotient singularity then a *P-resolution* $g\colon \tilde{Z} \to Z$ is a proper birational morphism such that $K_{\tilde{Z}}$ is relatively ample over Z and $\tilde{Z}$ has quotient singularities of class T. We can now state the main result of Kollár and Shepherd-Barron on deformations of cyclic quotient singularities (together with some improvements obtained subsequently by Stevens):

Theorem 6.4. [KSB88],[S91] *Let Z be a cyclic quotient singularity. Let $g\colon \tilde{Z} \to Z$ be a P-resolution of Z. The versal $\mathbb{Q}$-Gorenstein deformation space* $\operatorname{Def}^{\mathrm{QG}}(\tilde{Z})$ *of $\tilde{Z}$ is smooth. There is a morphism*

$$F\colon \operatorname{Def}^{\mathrm{QG}}(\tilde{Z}) \to \operatorname{Def}(Z)$$

to the versal deformation space of Z given by blowing down deformations (using $R^1 g_* \mathcal{O}_{\tilde{Z}} = 0$). The morphism F is an isomorphism onto an irreducible component of $\operatorname{Def}(Z)$*, and this establishes a bijective correspondence between P-resolutions of Z and irreducible components of* $\operatorname{Def}(Z)$.

We now describe a P-resolution of the cyclic quotient singularity Z determined by the pair (Y, D). Let $n \in N$ be the primitive integral inward normal of the half-plane $\bar{B} \subset M_{\mathbb{R}}$, so $B = (n \geq 0) \subset M_{\mathbb{R}}$. Let $\bar{\Xi} \subset \bar{B}$ be the half-plane $(n \geq 1) \subset M_{\mathbb{R}}$. Let Ξ be the convex polytope $\nu^{-1}(\bar{\Xi}) \subset B$, where we recall that $\nu\colon B \to \bar{B}$ denotes the piecewise linear identification of B and $\bar{B}$. Let $\tilde{Z}$ denote the quasiprojective toric variety associated to the polytope $\Xi \subset M_{\mathbb{R}}$. Explicitly, let Q denote the closure of $\mathbb{R}_{\geq 0} \cdot (\Xi \times \{1\}) \subset M_{\mathbb{R}} \oplus \mathbb{R}$, then $\tilde{Z} = \operatorname{Proj} \Bbbk[Q \cap (M \oplus \mathbb{Z})]$, where the grading is defined by the projection $M \oplus \mathbb{Z} \to \mathbb{Z}$. The surface $\tilde{Z}$ admits a proper birational map $g\colon \tilde{Z} \to Z$ to the cyclic quotient singularity $Z = \operatorname{Spec} \Bbbk[\sigma \cap M]$ given by the equality $\sigma \cap M = Q \cap (M \times \{0\})$.

Proposition 6.5. *The morphism $g\colon \tilde{Z} \to Z$ is a P-resolution of the cyclic quotient singularity Z.*

Proof. The exceptional locus of g is the union of the proper toric boundary divisors of $\tilde{Z}$, which correspond to the bounded edges of the polytope Ξ. The vertices of Ξ are the intersection points of Ξ with the rays ρ_i such that $1 \leq i \leq n$ and $l_i > 0$.

Write $\bar{v}_i := \nu(v_i) \in \bar{B} \subset M_{\mathbb{R}}$. Choose a basis of M such that $\bar{B} = \mathbb{R}_{\geq 0} \times \mathbb{R} \subset \mathbb{R}^2$. Then $\bar{v}_i = \begin{pmatrix} n_i \\ a_i \end{pmatrix}$ for some $n_i, a_i \in \mathbb{Z}_{>0}$, $(n_i, a_i) = 1$. (Note that n_i is independent of the choice of basis and a_i is determined modulo n_i.) Then it is an exercise in toric geometry to show that if $l_i > 0$ then the singularity of $\tilde{Z}$ corresponding to the vertex of Ξ on ρ_i is a cyclic quotient singularity of class T of type $\frac{1}{l_i n_i^2}(1, l_i n_i a_i - 1)$.

The canonical divisor $K_{\tilde{Z}}$ is relatively ample over Z by Lemma 4.3. Indeed, the lines considered in the statement of that lemma are the lines through the origin and the vertices of Ξ, by the construction of Ξ. $\square$

6.3. Main Theorem.

Theorem 6.6. *Let Z denote the cyclic quotient singularity dual to the singularity obtained by contracting $D_1, \ldots, D_n \subset Y$, and $B \subset Z$ its toric boundary.*

1. *The deformation* $(X', C')/(0 \in S')$ *induces an isomorphism*
$$(0 \in S'/W) \xrightarrow{\sim} D$$
from the quotient S'/W *to an irreducible component* D *of the versal deformation space of the pair* (Z, B).
2. *The group* W *is the monodromy group of the restriction of the versal deformation to* D.
3. *The irreducible component* D *corresponds to the* P*-resolution of* Z *described in Proposition 6.5 under the bijection of Theorem 6.4 and Proposition 6.7.*
4. *Every irreducible component of the versal deformation of a cyclic quotient singularity together with its toric boundary arises uniquely in this way for a pair* (Y, D) *as above such that the contraction* $p\colon Y \to Y'$ *is the minimal resolution of the dual cyclic quotient singularity* $q \in Y'$.

Proof. (1) Note of course that taking the quotient by W amounts to replacing the parameters $z^{E_{ij}}$, $j = 1, \ldots, l_i$ by the elementary symmetric functions $a_{i1}, \ldots, a_{il_i}$ in these parameters, for each $i = 1, \ldots, n$. Then the equation of the open patch $\mathcal{U}_i$ of X yields an equation

$$x_{i-1}x_{i+1} = x_i^{-D_i^2 - l_i}(x_i^{l_i} + a_{i1}x_i^{l_i-1} + \cdots + a_{il_i}) \tag{6.1}$$

for X', for each $i = 1, \ldots, n$. Let $D' \subset \mathrm{Def}(Z)$ be the irreducible component associated to the P-resolution $g\colon \tilde{Z} \to Z$ of Proposition 6.5. By (3) below the deformation X'/S' induces a morphism $S'/W \to D'$. If Z is Du Val of type A then $n = 1$ and the statement is easy to check. Now assume that Z is not Du Val of type A. By comparing the description of the tangent space of $\mathrm{Def}(Z)$ in [S91], §2.2 with the equations (6.1) above we deduce that the morphism $S'/W \to D'$ is a closed embedding. Let $F\colon \mathrm{Def}(Z, B) \to \mathrm{Def}(Z)$ be the forgetful map. By Proposition 6.7 the inverse image $D := F^{-1}(D')$ is an irreducible component of $\mathrm{Def}(Z, B)$ and $D \to D'$ is a closed embedding. The embedding $S'/W \to D'$ factors through D, and we observe that the dimension of D is equal to the dimension $\sum l_i$ of S'/W by Remark 6.8, hence S'/W maps isomorphically to D.

(2) This follows from [BC94].

(3) We need to exhibit a simultaneous P-resolution $\tilde{X}' \to X'/S'$ such that $K_{\tilde{X}'}$ is $\mathbb{Q}$-Cartier and the special fiber is the P-resolution $\tilde{Z} \to Z$ of Proposition 6.5. In fact we describe a proper birational morphism $\tilde{X} \to X/S$ whose restriction to S' is the desired simultaneous P-resolution. Let $\underline{\Xi}$ denote the polyhedral subdivision of the polytope $\Xi \subset B \subset M_{\mathbb{R}}$ induced by the decomposition of B. For $i = 1, \ldots, n$, let α_i be the multiplicity of $\bar{D}_i$ in the fiber of $\bar{f}\colon \bar{Y} \to \mathbb{A}^1$ over $0 \in \mathbb{A}^1$, and define $\alpha_0 = \alpha_{n+1} = 0$.

Then the vertices of Ξ are the points v_i/α_i for $i = 1, \ldots, n$. We remark that $(\alpha_i, \alpha_j) = 1$ for $|i - j| = 1$ (proof by induction on the number of blowups required to obtain $\bar{Y}/\mathbb{A}^1$ from a $\mathbb{P}^1$-bundle). Then $\tilde{X}$ is glued from patches

$$\tilde{\mathcal{U}_i} = V\left(x'_{i-1}x'_{i+1} - z^{D_i}\prod_{j=1}^{l_i}(u_i^{\alpha_i} + z^{E_{ij}})\right) \subset \left(\mathbb{A}^3_{x'_{i-1},u_i,x'_{i+1}} \Big/ \frac{1}{\alpha_i}(\alpha_{i-1}, -1, \alpha_{i+1})\right) \times S$$

for $i = 1, \ldots, n$. The morphism $\tilde{X} \to X$ can be described as follows: we have identifications

$$\tilde{\mathcal{U}_i} \setminus (u_i = 0) \xrightarrow{\sim} \mathcal{U}_i, \quad (x'_{i-1}, u_i, x'_{i+1}) \mapsto (x_{i-1}, x_i, x_{i+1}) = (u_i^{\alpha_{i-1}}x'_{i-1}, u_i^{\alpha_i}, u_i^{\alpha_{i+1}}x'_{i+1})$$

and

$$\tilde{\mathcal{U}_1} \setminus (x'_0 \neq 0) \xrightarrow{\sim} \mathcal{U}_0, \quad (x'_0, u_1, x'_2) \mapsto (x_0, x_1) = (x'_0, u_1),$$

$$\tilde{\mathcal{U}_n} \setminus (x'_{n+1} \neq 0) \xrightarrow{\sim} \mathcal{U}_{n+1}, \quad (x'_{n-1}, u_n, x'_{n+1}) \mapsto (x_n, x_{n+1}) = (u_n, x'_{n+1}).$$

These patch to give an identification $\tilde{X}^o \xrightarrow{\sim} X^o$ where $X^o = X \setminus \{0\} \times S$ and $\tilde{X}^o \subset \tilde{X}$ has complement of codimension at least 2. This identification extends to the desired birational morphism $\tilde{X} \to X$ because X is affine and both $\tilde{X}$ and X are normal.

(4) By Theorem 6.4 and Proposition 6.7 there is a bijective correspondence between P-resolutions of Z and irreducible components of $\operatorname{Def}(Z, B)$. So it suffices to show that P-resolutions arise uniquely by the procedure described in Proposition 6.5 for a pair (Y, D) such that $Y \to Y'$ is the minimal resolution of the dual singularity. This follows from [A98], §4. $\square$

6.4. **Deformations of pairs.** The deformations we construct are deformations of the *pair* (Z, B) consisting of a cyclic quotient singularity together with its toric boundary. The following result explains the relation between deformations of the pair and deformations of the singularity.

We remark that, in the theory of compact moduli of surfaces [KSB88], the deformation theory of the pair (Z, B) is more important than that of the singularity Z. This is because for a deformation $(\mathcal{Z}, \mathcal{B})/S$ of the pair (Z, B) the sheaf $\omega_{\mathcal{Z}/S}(\mathcal{B})$ is a line bundle (see Lemma 6.9 below), whereas for a deformation $\mathcal{Z}/S$ of a cyclic quotient singularity Z it is usually *not* the case that $\omega_{\mathcal{Z}/S}$ is a $\mathbb{Q}$-line bundle.

If Z is a normal surface and $B \subset Z$ is a reduced closed subscheme of codimension 1, a *deformation* $(\mathcal{Z}, \mathcal{B})/(0 \in S)$ of (Z, B) over a base $(0 \in S)$ is a deformation $\mathcal{Z}/S$ together with a closed subscheme $\mathcal{B} \subset \mathcal{Z}$, flat over S, such that $\mathcal{B}_0 = B$. If in addition Z is a $\mathbb{Q}$-Gorenstein surface, we say a deformation $(\mathcal{Z}, \mathcal{B})/(0 \in S)$ of (Z, B) is *$\mathbb{Q}$-Gorenstein* if it is locally induced by a deformation of (Z', B') where $Z' \to Z$ is a local index one cover and B' is the reduced inverse image of B.

Proposition 6.7. *Let (Z, B) be a cyclic quotient singularity together with its toric boundary.*

(1) *Let*

$$F\colon \operatorname{Def}(Z, B) \to \operatorname{Def}(Z)$$

be the forgetful map from the deformation space of the pair (Z, B) to the deformation space of Z. Then F induces a bijection between the irreducible components of $\operatorname{Def}(Z)$ and $\operatorname{Def}(Z, B)$ given by $D \mapsto F^{-1}(D)$. Moreover $F^{-1}(D) \to D$ is a closed embedding unless Z is a Du Val singularity of type A, in which case it is smooth of relative dimension 1.

(2) *Let $g\colon \tilde{Z} \to Z$ be a P-resolution of Z and $D \subset \operatorname{Def}(Z)$ the associated irreducible component. Let $\tilde{B} \subset \tilde{Z}$ be the reduced inverse image of B. Let $\operatorname{Def}^{QG}(\tilde{Z}, \tilde{B})$ be the $\mathbb{Q}$-Gorenstein deformation space of the pair $(\tilde{Z}, \tilde{B})$. Then $\operatorname{Def}^{QG}(\tilde{Z}, \tilde{B})$ is smooth and there is a morphism*

$$G\colon \operatorname{Def}^{QG}(\tilde{Z}, \tilde{B}) \to \operatorname{Def}(Z, B)$$

to the deformation space of the pair (Z, B) given by blowing down deformations. The morphism G is an isomorphism onto the irreducible component $F^{-1}(D)$ of $\operatorname{Def}(Z, B)$.

Remark 6.8. The dimension of D and of $F^{-1}(D)$ may be computed as follows. Let $g\colon \tilde{Z} \to Z$ be the P-resolution associated to the irreducible component $D \subset \operatorname{Def}(Z)$. Let $\{Q_i\}_{i=1}^{r}$ be the nodes of $\tilde{B} \subset \tilde{Z}$. Then $Q_i \in \tilde{Z}$ is a singularity of type $\frac{1}{d_i n_i^2}(1, d_i n_i a_i - 1)$ for some d_i, n_i, a_i (the smooth case $d_i = n_i = 1$ is not excluded). Then $F^{-1}(D)$ has dimension $\sum d_i$. Let $\tilde{Z}' \to \tilde{Z}$ be the minimal resolution of $\tilde{Z}$ and let $E'_1, \ldots, E'_{r-1} \subset \tilde{Z}'$ denote the strict transforms of the exceptional cuves of g. Let s denote the number of indices i for which $n_i = 1$. Then D has dimension $\sum_{i=1}^{r} d_i - s + \sum_{j=1}^{r-1}(-{E'_j}^2 - 1)$, see [KSB88], Corollary 3.20.

Proof. Let $g\colon \tilde{Z} \to Z$ be a P-resolution and $\tilde{B}$ the inverse image of B with its reduced structure. Then g is toric because Z is a cyclic quotient, and the divisor $\tilde{B} \subset \tilde{Z}$ is the toric boundary of $\tilde{Z}$. In particular the singularities of $\tilde{Z}$ are of type $\frac{1}{dn^2}(1, dna - 1)$.

Let $\tilde{\mathfrak{Z}} \to \tilde{Z}$ be the Deligne–Mumford stack with coarse moduli space $\tilde{Z}$ defined by the local index one covers $\frac{1}{dn}(1, -1) \to \frac{1}{dn^2}(1, dna - 1)$ of the singularities of $\tilde{Z}$. Thus, locally over $\tilde{Z}$, the morphism $\tilde{\mathfrak{Z}} \to \tilde{Z}$ is of the form $[U/\mu_n] \to U/\mu_n$ where

$$U = \mathbb{A}^2_{u,v} \Big/ \frac{1}{dn}(1, -1) = V(xy - z^{dn}) \subset \mathbb{A}^3_{x,y,z}$$

and the μ_n action has weights $\frac{1}{n}(1,-1,a)$ on x, y, z. The deformations of $\tilde{\mathfrak{Z}}$ are the deformations of the coarse moduli space $\tilde{Z}$ which are locally induced by an equivariant deformation of the index one cover, that is, the $\mathbb{Q}$-Gorenstein deformations of $\tilde{Z}$.

Let $\tilde{\mathfrak{B}} \subset \tilde{\mathfrak{Z}}$ denote the closed substack corresponding to $\tilde{B} \subset \tilde{Z}$. Let $\{P_i\}_{i=1}^r$ denote the nodes of $\tilde{\mathfrak{B}}$. Let $T_{\tilde{\mathfrak{Z}}}(-\log \tilde{\mathfrak{B}})$ be the sheaf of vector fields with logarithmic zeroes along $\tilde{\mathfrak{B}}$. This is the sheaf of first order infinitesimal automorphisms of the pair $(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}})$. We have $T_{\tilde{\mathfrak{Z}}}(-\log \tilde{\mathfrak{B}}) \simeq \mathcal{O}_{\tilde{\mathfrak{Z}}}^{\oplus 2}$ because the pair $(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}})$ is toric. So $H^i(T_{\tilde{\mathfrak{Z}}}(-\log \tilde{\mathfrak{B}})) = 0$ for $i > 0$. It follows that the local-to-global map

$$\mathrm{Def}(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}}) \to \prod_i \mathrm{Def}(P_i \in \tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}})$$

is an isomorphism. Let $Q_i \in \tilde{Z}$ be the image of $P_i \in \tilde{\mathfrak{Z}}$. Let the singularity $Q_i \in \tilde{Z}$ be of type $\frac{1}{d_i n_i^2}(1, d_i n_i a_i - 1)$. The $\mathbb{Q}$-Gorenstein deformation space of the pair

$$\left(0 \in \left(\mathbb{A}^2_{u,v} \Big/ \frac{1}{dn^2}(1, dna - 1)\right), V(uv)\right)$$

is smooth of dimension d (this includes the smooth case $d = n = 1$), with versal deformation

$$((xy = z^{dn} + a_{d-1} z^{(d-1)n} + \cdots + a_0), V(z)) \subset \left(\mathbb{A}^3_{x,y,z} \Big/ \frac{1}{n}(1,-1,a)\right) \times \mathbb{A}^d_{a_0,\ldots,a_{d-1}}.$$

Thus $\mathrm{Def}(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}})$ is smooth of dimension $\sum_i d_i$.

We note that if Z is a Du Val singularity of type A, then the Proposition is trivially true: $\mathrm{Def}(Z)$ is smooth, the map F is smooth of relative dimension 1, Z is Gorenstein, and the only P-resolution is the identity. We exclude this case in what follows.

We assert that the forgetful map $\mathrm{Def}(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}}) \to \mathrm{Def}(\tilde{\mathfrak{Z}})$ is a closed embedding. To prove this, we need to show that all first order embedded deformations of $\tilde{\mathfrak{B}} \subset \tilde{\mathfrak{Z}}$ are induced by infinitesimal automorphisms of $\tilde{\mathfrak{Z}}$, that is, the map $H^0(T_{\tilde{\mathfrak{Z}}}) \to H^0(\mathcal{N})$ is surjective, where $\mathcal{N}$ is the normal bundle of $\tilde{\mathfrak{B}} \subset \tilde{\mathfrak{Z}}$. Note that $\tilde{\mathfrak{B}}$ is a Cartier divisor on $\tilde{\mathfrak{Z}}$ (because $K_{\tilde{Z}} + \tilde{B} = 0$ and $\tilde{\mathfrak{Z}} \to \tilde{Z}$ is the index one cover). So $\mathcal{N}$ is a line bundle on $\tilde{\mathfrak{B}}$. Let $\mathcal{N}'$ denote the kernel of the surjection

$$\mathcal{N} \to \bigoplus_i \mathcal{N} \otimes k(P_i).$$

Then we have an exact sequence

$$0 \to T_{\tilde{\mathfrak{Z}}}(-\log \tilde{\mathfrak{B}}) \to T_{\tilde{\mathfrak{Z}}} \to \mathcal{N}' \to 0.$$

Hence $H^0(T_{\tilde{\mathfrak{Z}}}) \to H^0(\mathcal{N}')$ is surjective (because $H^1(T_{\tilde{\mathfrak{Z}}}(-\log \tilde{\mathfrak{B}})) = 0$, see above). We claim that $H^0(\mathcal{N}') = H^0(\mathcal{N})$ (here we use that Z is *not* a Du Val singularity of type A): Indeed, on $\tilde{Z}$ we have $K_{\tilde{Z}} + \tilde{B} = 0$ and $K_{\tilde{Z}}$ is relatively ample, so $\mathcal{N}$ is negative on proper components of $\tilde{\mathfrak{B}}$. It follows that the map $H^0(\mathcal{N}) \to \bigoplus H^0(\mathcal{N} \otimes k(P_i))$ is zero unless $\tilde{Z} = Z$. If $\tilde{Z} = Z$ then the action of μ_n on the fiber $\mathcal{N} \otimes k(P)$ of $\mathcal{N}$ at the singular point $P \in \tilde{\mathfrak{Z}}$ is nontrivial because $n > 1$ by assumption, so $H^0(\mathcal{N} \otimes k(P)) = 0$. Hence $H^0(\mathcal{N}') = H^0(\mathcal{N})$ as claimed, and combining we obtain that $H^0(T_{\tilde{\mathfrak{Z}}}) \to H^0(\mathcal{N})$ is surjective as required.

We have a map $\mathrm{Def}(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}}) \to \mathrm{Def}(Z, B)$ given by blowing down deformations. Indeed, by [W76], Thm. 1.4(c), deformations $\tilde{\mathcal{Z}}$ and $\tilde{\mathcal{B}}$ of $\tilde{\mathfrak{Z}}$ and $\tilde{\mathfrak{B}}$ over an Artinian base S induce deformations $\mathcal{Z}$ and $\mathcal{B}$ of Z and B, respectively. We just need to check that the closed embedding $\tilde{\mathcal{B}} \subset \tilde{\mathcal{Z}}$ induces a closed embedding $\mathcal{B} \subset \mathcal{Z}$. Pushing forward the exact sequence

$$0 \to \mathcal{O}_{\tilde{\mathcal{Z}}}(-\tilde{\mathcal{B}}) \to \mathcal{O}_{\tilde{\mathcal{Z}}} \to \mathcal{O}_{\tilde{\mathcal{B}}} \to 0$$

gives

$$0 \to g_*\mathcal{O}_{\tilde{\mathcal{Z}}}(-\tilde{\mathcal{B}}) \to \mathcal{O}_{\mathcal{Z}} \to \mathcal{O}_{\mathcal{B}} \to R^1 g_*\mathcal{O}_{\tilde{\mathcal{Z}}}(-\tilde{\mathcal{B}}) \to 0.$$

We have $K_{\tilde{Z}} + \tilde{B} = 0$, and $\omega_{\tilde{\mathcal{Z}}/S}(\tilde{\mathcal{B}})$ is invertible by Lemma 6.9, so $\omega_{\tilde{\mathcal{Z}}/S}(\tilde{\mathcal{B}}) \simeq \mathcal{O}_{\tilde{\mathcal{Z}}}$ (using $R^1 g_*\mathcal{O}_{\tilde{Z}} = 0$). Thus $R^1 g_*\mathcal{O}_{\tilde{\mathcal{Z}}}(-\tilde{\mathcal{B}}) = R^1 g_*\omega_{\tilde{\mathcal{Z}}/S} = 0$ by Grauert-Riemenschneider vanishing [KM98], p. 73, Cor. 2.68. Thus $\mathcal{B} \subset \mathcal{Z}$ is a closed subscheme as required.

We claim that $\mathrm{Def}(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}}) \to \mathrm{Def}(Z, B)$ is an isomorphism onto an irreducible component of $\mathrm{Def}(Z, B)$, which is the inverse image $F^{-1}(D)$ of the irreducible component $D \subset \mathrm{Def}(Z)$ corresponding to the P-resolution g under the forgetful map $F\colon \mathrm{Def}(Z, B) \to \mathrm{Def}(Z)$. The map $\mathrm{Def}(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}}) \to \mathrm{Def}(Z, B)$ is a closed embedding with image contained in $F^{-1}(D)$ because $\mathrm{Def}(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}}) \to \mathrm{Def}(\tilde{\mathfrak{Z}})$ is a closed embedding and $\mathrm{Def}(\tilde{\mathfrak{Z}}) \to \mathrm{Def}(Z)$ is an isomorphism onto D. So it suffices to show that a map $(0 \in S) \to F^{-1}(D)$ from a smooth curve germ $(0 \in S)$ lifts to $\mathrm{Def}(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}})$. Let $(\mathcal{Z}, \mathcal{B})/S$ be the induced deformation of the pair (Z, B). The divisor $K_{\mathcal{Z}} + \mathcal{B}$ is Cartier by Lemma 6.9. Thus $(\mathcal{Z}, Z + \mathcal{B})$ is log canonical by inversion of adjunction [K07]. By Theorem 6.4 the deformation $\mathcal{Z}/S$ is induced by a $\mathbb{Q}$-Gorenstein deformation $f\colon \tilde{\mathcal{Z}} \to \mathcal{Z}/S$ of the P-resolution $g\colon \tilde{Z} \to Z$. Let $\tilde{\mathcal{B}}$ be the inverse image of $\mathcal{B}$ with its reduced structure. Then $K_{\tilde{\mathcal{Z}}} + \tilde{Z} + \tilde{\mathcal{B}} = f^*(K_{\mathcal{Z}} + Z + \mathcal{B})$ because $K_{\tilde{Z}} + \tilde{B} = g^*(K_Z + B)$. It follows that $(\tilde{\mathcal{Z}}, \tilde{Z} + \tilde{B})$ is log canonical. Now by [A08], Thm. 0.1, $\tilde{\mathcal{B}}_0$ is reduced, so $(\tilde{\mathcal{Z}}, \tilde{\mathcal{B}})/S$ is a $\mathbb{Q}$-Gorenstein deformation of $(\tilde{Z}, \tilde{B})$. We deduce that $\mathrm{Def}(\tilde{\mathfrak{Z}}, \tilde{\mathfrak{B}}) \to \mathrm{Def}(Z, B)$ is an isomorphism onto $F^{-1}(D)$ (with its reduced structure) and $F^{-1}(D) \to D$ is a closed embedding. ☐

Lemma 6.9. *Let (Z, B) be a surface cyclic quotient singularity together with its toric boundary. Let $(\mathcal{Z}, \mathcal{B})/(0 \in S)$ be a deformation of (X, D). Then $\omega_{\mathcal{Z}/S}(\mathcal{B})$ is invertible.*

Proof. We glue on another component to obtain a deformation of a reducible surface without boundary, and obtain the result by restriction. Let $\mathcal{W} = (0 \in \mathbb{A}^2_{x,y}) \times S$

be a smooth surface germ over S. Observe that the curve singularity B is a node. Choose an embedding $\mathcal{B} \subset \mathcal{W}$ over S, and glue $\mathcal{Z}$ and $\mathcal{W}$ along $\mathcal{B}$. We obtain a flat family $\mathcal{X}/S$. The special fiber X is a Gorenstein semi log canonical singularity — a so-called degenerate cusp. In particular $\omega_{\mathcal{X}/S}$ is invertible (because ω_X is invertible, and the relative dualising sheaf commutes with base change for a flat family of Cohen-Macaulay schemes). Write $\mathcal{U} := \mathcal{X} \setminus \mathcal{W} = \mathcal{Z} \setminus \mathcal{B}$. We have a natural identification

$$\omega_{\mathcal{X}/S}|_{\mathcal{U}} = \omega_{\mathcal{U}/S} = \omega_{\mathcal{Z}/S}|_{\mathcal{U}}.$$

We claim that this extends to an isomorphism

$$\omega_{\mathcal{X}/S}|_{\mathcal{Z}} \xrightarrow{\sim} \omega_{\mathcal{Z}/S}(\mathcal{B})$$

Both $\omega_{\mathcal{X}/S}|_{\mathcal{Z}}$ and $\omega_{\mathcal{Z}/S}(\mathcal{B})$ satisfy the relative S_2 property $i_*i^*\mathcal{F} = \mathcal{F}$, where $i\colon \mathcal{Z}^o \subset \mathcal{Z}$ is the inclusion of a open subset whose complement has finite fibers. So it suffices to check the claim in relative codimension 1, where it is obvious. Hence $\omega_{\mathcal{Z}/S}(B)$ is invertible as required. $\square$

REFERENCES

[AS10] M. Abouzaid, P. Seidel, An open string analogue of Viterbo functoriality, Geom. Topol. 14 (2010), no. 2, 627–718.

[AC11] D. Abramovich, Q. Chen, Stable logarithmic maps to Deligne-Faltings pairs II, arXiv:1102.4531.

[A02] V. Alexeev, Complete moduli in the presence of semiabelian group action, Ann. of Math. (2) 155 (2002), no. 3, 611–708.

[A08] V. Alexeev, Limits of stable pairs, Pure Appl. Math. Q. 4 (2008), no. 3, 767–783.

[A98] K. Altmann, P-resolutions of cyclic quotients from the toric viewpoint, in Singularities (Oberwolfach, 1996), 241–250, Progr. Math. 162, Birkhäuser, 1998.

[A76] M. Artin, Lectures on deformations of singularities, Lectures on Mathematics and Physics 54, Tata Inst. Fund. Res., 1976.

[A07] D. Auroux, Mirror symmetry and T-duality in the complement of an anticanonical divisor, J. Gökova Geom. Topol. 1 (2007), 51-91, and preprint arXiv:0706.3207 (2007).

[A09] D. Auroux, Special Lagrangian fibrations, wall-crossing, and mirror symmetry, in Geometry, analysis, and algebraic geometry, Surv. Differ. Geom. 13, 1–47, Int. Press, 2009.

[AKO06] D. Auroux, L. Katzarkov, D. Orlov, Mirror symmetry for del Pezzo surfaces: vanishing cycles and coherent sheaves, Invent. Math. 166 (2006), no. 3, 537-582.

[AMRT75] A. Ash, D. Mumford, M. Rapoport, Y. Tai, Smooth compactification of locally symmetric varieties, Math. Sci. Press, 1975.

[B93] V. Batyrev, Quantum cohomology rings of toric manifolds, in Journées de Géométrie Algébrique d'Orsay, Astérisque No. 218 (1993), 9-34.

[BC94] K. Behnke, J. Christophersen, M-resolutions and deformations of quotient singularities, Amer. J. Math. 116 (1994), no. 4, 881–903.

[B78] J. Bingener, Über formale komplexe Räume, Manuscripta Math. 24 (1978), no. 3, 253–293.

[BKR01] T. Bridgeland, A. King, M. Reid, The McKay correspondence as an equivalence of derived categories, J. Amer. Math. Soc. 14 (2001), no. 3, 535-554.

[C1869] A. Cayley, A Memoir on Cubic Surfaces, Philosophical Transactions of the Royal Society of London 159 (1869), 231–326.

[CPS] M. Carl, M. Pumperla, and B. Siebert, A tropical view of Landau-Ginzburg models, in preparation.

[CO06] C.-H. Cho, Y.-G. Oh, Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds, Asian J. Math. **10**, (2006), 773-814.

[D09] T. Dyckerhoff, Compact generators in categories of matrix factorizations, preprint arXiv:0904.4713 .

[E95] D. Eisenbud, *Commutative algebra with a view toward Algebraic Geometry,* Graduate Texts in Mathematics 150, Springer-Verlag, New York, 1995.

[FG09] V. Fock and A. Goncharov, Cluster ensembles, quantization and the dilogarithm, Invent. Math. 175 (2009), no. 2, 223286.

[F83a] R. Friedman, Global smoothings of varieties with normal crossings, Ann. of Math. (2) 118 (1983), no. 1, 75–114.

[F83b] R. Friedman, The mixed Hodge structure of an open variety, unpublished manuscript, 1983.

[FM83] R. Friedman and R. Miranda, Smoothing cusp singularities of small length, Math. Ann. 263 (1983), no. 2, 185–212.

[FP84] R. Friedman and H. Pinkham, Smoothings of cusp singularities via triangle singularities. With an appendix by H. Pinkham, Compositio Math. 53 (1984), no. 3, 303–324.

[F75] A. Fujiki, On the blowing down of analytic spaces, Publ. Res. Inst. Math. Sci. 10 (1974/75), 473–507.

[GMN1] D. Gaiotto, G. Moore, and A. Neitzke, Four- dimensional wall-crossing via three-dimensional field theory," Commun. Math. Phys. 299 (2010) 163–224.

[GMN2] D. Gaiotto, G. Moore, and A. Neitzke, Wall-crossing, Hitchin Systems, and the WKB Approximation, arXiv:0907.3987.

[GMN3] D. Gaiotto, G. Moore, and A. Neitzke, Framed BPS States, arXiv:1006.014.

[Giv] A. Givental, Homological geometry. I. Projective hypersurfaces. Selecta Math. 1 (1995), 325–345.

[G07] A. Goncharov, Pentagon relation for the quantum dilogarithm and quantized $M_{0,5}^{\text{cyc}}$. Geometry and dynamics of groups and spaces, 415428, Progr. Math., 265, Birkhuser, Basel, 2008.

[G09] M. Gross, Mirror symmetry for $\mathbb{P}^2$ and tropical geometry, arXiv:0903.1378v1 [math.AG] .

[G11] M. Gross, Tropical geometry and mirror symmetry, CBMS Regional Conf. Ser. in Math. 114, A.M.S., 2011.

[K3] M. Gross, P Hacking, S. Keel, Theta functions for K3 surfaces, in preparation.

[GPS09] M. Gross, R. Pandharipande, B. Siebert, The tropical vertex, arXiv:0902.0779v1 [math.AG].

[GS06] M. Gross, B. Siebert, Mirror symmetry via logarithmic degeneration data, I. J. Differential Geom., **72**, (2006) 169–338.

[GS07] M. Gross, B. Siebert, From real affine geometry to complex geometry, arXiv:math/0703822v2 [math.AG] .

[GS08] M. Gross, B. Siebert, An invitation to toric degenerations, arXiv:0808.2749v1 [math.AG] .

[GS11] M. Gross, B. Siebert, Logarithmic Gromov-Witten invariants, arXiv:1102.4322v1 [math.AG] .

[G60] A. Grothendieck, Éléments de géométrie algébrique I. Le langage des schémas, Inst. Hautes Études Sci. Publ. Math. No. 4.

[G61] A. Grothendieck, Éléments de géométrie algébrique III, Étude cohomologique des faisceaux cohérents I, Inst. Hautes Études Sci. Publ. Math. No. 11 (1961).

[H04] P. Hacking, Compact moduli of plane curves, Duke Math. J. 124 (2004), no. 2, 213–257.

[Hi73] F. Hirzebruch, Hilbert modular surfaces, Enseignement Math., **19**, (1973), 183-281.

[H77] R. Hartshorne, Algebraic geometry, Grad. Texts in Math. 52, Springer, 1977.

[IP03] E.-N. Ionel, T. Parker, Relative Gromov-Witten invariants, Ann. of Math. **157** (2003), 45–96.

[KKP08] L. Katzarkov, M. Kontsevich, T. Pantev, Hodge theoretic aspects of mirror symmetry, in From Hodge theory to integrability and TQFT tt^*-geometry, 87-174, Proc. Sympos. Pure Math. 78, A.M.S., 2008.

[K07] M. Kawakita, *Inversion of adjunction on log canonicity*, Invent. Math. **167** (2007), no. 1, 129–133.

[KM98] J. Kollár, S. Mori, Birational geometry of algebraic varieties, Cambridge Tracts in Math. 134. C.U.P., 1998.

[KSB88] J. Kollár, N. Shepherd-Barron, Threefolds and deformations of surface singularities, Invent. Math. 91 (1988), no. 2, 299–338.

[K11] J. Kollár et al., Moduli of varieties of general type, book in preparation, preliminary version available at http://www.math.princeton.edu/ kollar/

[K95] M. Kontsevich, Homological algebra of mirror symmetry, Proceedings of the International Congress of Mathematicians, Vol. 1, (Zürich, 1994), 120-139, Birkhäuser, 1995.

[KS06] M. Kontsevich, Y. Soibelman: *Affine structures and non-Archimedean analytic spaces*, in: *The unity of mathematics* (P. Etingof, V. Retakh, I.M. Singer, eds.), 321–385, Progr. Math. 244, Birkhäuser 2006.

[L02] S. Lang, Algebra, 3rd ed., Grad. Texts in Math. 211, Springer, 2002.

[L73] H. Laufer, Taut two-dimensional singularities, Math. Ann. 205 (1973), 131–164.

[Li00] J. Li, Stable morphisms to singular schemes and relative stable morphisms, J. Diff. Geom. **57**, (2000), 509–578.

[Li02] J. Li, A degeneration formula of GW-invariants, J. Diff. Geom., **60**, (2002) 199-293.

[LR01] A.-M. Li, Y. Ruan, Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds, I, Invent. Math. **145** (2001), 151–218.

[L08] P. Lisca, On symplectic fillings of lens spaces, Trans. Amer. Math. Soc. 360 (2008), no. 2, 765-799.

[L76] E. Looijenga, Root systems and elliptic curves, Invent. Math. 38 (1976), 17–32.

[L81] E. Looijenga, Rational surfaces with an anticanonical cycle. Ann. of Math. (2) 114 (1981), no. 2, 267–322.

[M82] J. Mérindol, Les singularités simples elliptiques, leurs déformations, les surfaces de del Pezzo et les transformations quadratiques. Ann. Sci. École Norm. Sup. (4) 15 (1982), no. 1, 17–44

[Ma89] H. Matsumura, Commutative ring theory, C.U.P., 1989.

[Mk05] G. Mikhalkin, Enumerative tropical algebraic geometry in $\mathbb{R}^2$. J. Amer. Math. Soc. 18 (2005), no. 2, 313–377.

[Mum] D. Mumford, An analytic construction of degenerating abelian varieties over complete rings, Compositio Math. **24**, (1972), 239-272.

[N76] Y. Namikawa, A new compactification of the Siegel space and degeneration of Abelian varieties. I. Math. Ann. 221 (1976), no. 2, 97–141.

[N80] I. Nakamura, Inoue-Hirzebruch surfaces and a duality of hyperbolic unimodular singularities, Math. Ann. 252 (1980), no. 3, 221–235.

[NPP10] A. Némethi, P. Popescu-Pampu, On the Milnor fibres of cyclic quotient singularities, Proc. Lond. Math. Soc. (3) 101 (2010), no. 2, 554-588.

[Oa04] A. Oancea, A survey of Floer homology for manifolds with contact type boundary or Symplectic homology, in Symplectic geometry and Floer homology, 51-91, Ensaios Mat. 7, Soc. Brasil. Mat., Rio de Janeiro, 2004.

[Ob04] A. Oblomkov, Double affine Hecke algebras of rank 1 and affine cubic surfaces, Int. Math. Res. Not. 2004, no. 18, 877–912, and arXiv:math/0306393v4 [math.RT] .

[O09] M. Olsson, Compactifying moduli spaces for abelian varieties. Lecture Notes in Mathematics, 1958. Springer-Verlag, Berlin, 2008

[Or04] D. Orlov, Triangulated categories of singularities and D-branes in Landau-Ginzburg models, Proc. Steklov Inst. Math. 246, no. 3 (2004), 227-248.

[P09] T. Pantev, Notes from a lecture "Homological mirror symmetry for del Pezzo surfaces" at Princeton University, November 2009.

[P74] H. Pinkham, Deformations of algebraic varieties with $\mathbb{G}_m$ action, Astérisque 20, S.M.F., 1974.

[R83] M. Reid, Decomposition of toric morphisms, in Arithmetic and geometry, Vol. II, 395-418, Progr. Math. 36, Birkhäuser, 1983.

[R09] A. Ritter, The Novikov theory for symplectic cohomology and exact Lagrangian embeddings, PhD thesis, MIT, 2009.

[R10] A. Ritter, Topological quantum field theory structure on symplectic cohomology, preprint arXiv:1003.1781 (2010).

[S02] P. Seidel, Fukaya categories and deformations, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 351-360, Higher Ed. Press, Beijing, 2002. in Current developments in mathematics, 2006, 211–253, Int. Press, 2008.

[S08b] P. Seidel, A_∞-subalgebras and natural transformations, Homology, Homotopy Appl. 10 (2008), no. 2, 83-114.

[S08a] P. Seidel, A biased view of symplectic cohomology, Homology, Homotopy Appl. 10 (2008), no. 2, 83-114.

[S09a] P. Seidel, Symplectic homology as Hochschild homology, in Algebraic geometry–Seattle 2005, Proc. Sympos. Pure Math. 80, Part 1, 415-434, A.M.S., 2009.

[S09b] P. Seidel, Fukaya categories and Picard-Lefschetz theory, Zürich Lectures in Advanced Mathematics, E.M.S, 2008.

[S91] J. Stevens, On the versal deformation of cyclic quotient singularities, in Singularity theory and its applications, Part I (Coventry, 1988/1989), 302–319, Lecture Notes in Math. 1462, Springer, 1991.

[SYZ96] A. Strominger, S.-T. Yau, and E. Zaslow, Mirror Symmetry is T-duality, Nucl. Phys. **B479**, (1996) 243–259.

[S03] M. Symington, Four dimensions from two in symplectic topology, in Topology and geometry of manifolds (Athens, GA, 2001), 153–208, Proc. Sympos. Pure Math. 71, A.M.S., 2003.

[W76] J. Wahl, Equisingular deformations of normal surface singularities I, Ann. of Math. (2) 104 (1976), no. 2, 325–356.

UCSD Mathematics, 9500 Gilman Drive, La Jolla, CA 92093-0112, USA
E-mail address: `mgross@math.ucsd.edu`

Department of Mathematics and Statistics, Lederle Graduate Research Tower, University of Massachusetts, Amherst, MA 01003-9305
E-mail address: `hacking@math.umass.edu`

Department of Mathematics, 1 University Station C1200, Austin, TX 78712-0257
E-mail address: `keel@math.utexas.edu`